



# HHS Public Access

Author manuscript

*Rehabil Nurs.* Author manuscript; available in PMC 2015 November 02.

Published in final edited form as:

*Rehabil Nurs.* 2014 ; 39(1): 16–25. doi:10.1002/rnj.107.

## Negative Impacts of Shiftwork and Long Work Hours

Claire C. Caruso, PhD, RN

National Institute for Occupational Safety and Health, Cincinnati, Ohio, USA

### Abstract

**Purpose**—Healthcare organizations often have to provide patient care around the clock. Shift work (any shift outside of 7 a.m. to 6 p.m) and long work hours increase the risk for short sleep duration and sleep disturbances. Thirty-two percent of healthcare workers report they do not get enough sleep. The purpose of the article is to give an overview of the wide range of risks to nurses, patients, and employers that are linked to shift work, long work hours, and poor sleep from other sources.

**Findings**—Shift work and long work hours increase the risk for reduced performance on the job, obesity, injuries, and a wide range of chronic diseases. In addition, fatigue-related errors could harm patients. Fatigued nurses also endanger others during their commute to and from work.

**Conclusion and Clinical Relevance**—The key strategy to reduce these risks is making sleep a priority in the employer’s systems for organizing work and in the nurse’s personal life.

### Keywords

Shift work; occupational diseases; occupational injury; occupational exposure; work schedule tolerance; circadian rhythms; job stress; overtime work; extended work shifts; sleepiness

### Introduction

Scientific evidence is growing that adequate sleep is a biological need for life similar to the need to eat and drink and is critical to maintain life and health and to work safely (Everson, 2009). Sleep duration of 7–8 hours a night is associated with a lower risk of obesity, diabetes, high blood pressure, myocardial infarction, and cerebral vascular accidents as well as reduced risk for injuries and errors (Colten & Altevogt, 2006). Recent studies warn that a growing number of Americans are not getting enough sleep (CDC, 2008, 2011). Between the periods of 1985–1990 and 2004–2007, the percentage of healthcare workers reporting 6 or less hours sleep each day (a level considered too short by sleep experts) increased from 28% to 32% (Luckhaupt, Tak, & Calvert, 2010).

This trend for shorter sleep is likely linked to several factors. Nursing staff in facilities that operate around the clock are commonly exposed to shift work and long work hours. These

---

**Correspondence** Claire C. Caruso, Research Health Scientist, National Institute, for Occupational Safety and Health, 4676, Columbia Parkway MS C-24, Cincinnati, OH, 45226-1998., ZHL1@cdc.gov.

The findings and conclusions in this report are those of the author and do not necessarily represent the views of the National Institute for Occupational Safety and Health.

demanding schedules can lead to difficulties with sleep because of the need to sleep at irregular times and at times that are out of phase with circadian rhythms. Misalignment of sleep with circadian rhythms leads to difficulties with falling asleep, more arousals during sleep, and early awakenings leading to poorer sleep quality and shorter sleep duration. Also, sleep duration may be shortened by insufficient time between work shifts and the competing demands of work and personal life. In addition, economic pressures could push healthcare workers to take on second jobs or work longer hours. Another factor is lack of knowledge about the importance of sleep which can lead people to cut their sleep time for other activities (Colten & Altevogt, 2006). Of all industrial sectors, health care is the second highest sector for the number of workers reporting short sleep duration (5 million workers which is 32% of the total healthcare and social assistance workforce of 16 million Americans) (Luckhaupt et al., 2010). The purpose of this article is to give an overview of the wide range of risks to nurses, their patients, and employers that are linked to shift work, long work hours, and poor sleep from other sources.

### **Inadequate sleep linked to shift work, long work hours, and sleep disorders**

Scientific evidence provides strong support for the link between sleep problems and shift work. A study of a representative sample from the Midwest found 32% of night-shift workers and 26% of rotating-shift workers reported long-term insomnia and excessive sleepiness (Drake, Roehrs, Richardson, Walsh, & Roth, 2004). Ingre & Åkerstedt (2004) studied 169 sets of monozygotic twins who were discordant on exposure to night-shift work. History of night shift significantly increased risk for sleep disturbances into retirement years. A meta-analysis by Pilcher, Lambert, & Huffcutt (2000) of 36 studies found that permanent day-shift workers slept an average of 7.0 hours per day, permanent and rotating evening-shift workers slept 7.6–8.1 hours, permanent night-shift workers slept 6.6 hours, and rotating night-shift workers slept the least, 5.9 hours.

In a review of studies examining extended work shifts, Knauth (2007) reported 13 studies showed shifts longer than 8 hours had negative effects on sleep, eight studies showed mixed results, and four studies showed positive effects. Long weekly work hours and overtime have been associated with shorter sleep duration or sleep disturbances in several studies (Hayashi, Kobayashi, Yamaoka, & Yano, 1996; Sasaki et al., 1999). Geiger-Brown et al. (2012) examined sleep duration in nurses across a series of work shifts and found a mean sleep duration of 5.5 hours between 12-hour shifts. Basner et al. (2007) conducted a study of a nationally representative sample of Americans and reported one additional hour of work a day was associated with one half-hour less sleep.

Other sources of poor sleep and sleepiness are sleep disorders, other chronic diseases and certain medications (Smolensky, Di Milia, Ohayon, & Philip, 2011). Sleep disorders are common, afflicting 50–70 million Americans, but are often undiagnosed and therefore not treated (Colten & Altevogt, 2006). Insomnia increases with age. According to a National Sleep Foundation (2007) poll, sleep problems a few nights a week increased from 33% of women aged 18–24 to 48% of women aged 55–64. Sleepiness is a side effect of certain commonly used medications, such as benzodiazepines, narcotic analgesics, some antihistamines, some antidepressants, and prescription and over-the-counter medications

used in the treatment of insomnia (Smolensky et al., 2011). Several chronic diseases can promote daytime sleepiness and comprise cognitive functioning such as, arthritis, asthma, chronic fatigue syndrome, chronic obstructive pulmonary disease, and rhinitis. See Smolensky et al. (2011) for more information.

## Declines in neurocognitive functioning and performance

Laboratory studies provide evidence for a wide range of cognitive declines associated with sleep deprivation (Goel, Rao, Durmer, & Dinges, 2009). Without adequate sleep, people feel sleepy, are less alert, and can fall asleep involuntarily. Sleep deprivation increases pressure for the brain to fall asleep. The transition from wakefulness to sleep is abrupt and rapid and occurs in an on-off-type switch in the brain stem (Schwartz & Roth, 2008). This can lead to dangerous situations if the brain is under high pressure for sleep and the person falls asleep involuntarily during critical times, such as when driving or providing patient care.

Involuntary episodes of microsleep, which are short periods of sleep a few seconds long, can also occur. During microsleeps, the person may appear to be awake (eyes open), but his or her brain will not process information and lapses in attention occur (Boyle, Tippin, Paul, & Rizzo, 2008).

A sleep-deprived person cannot reliably force him or herself to stay awake and alert. Studies of medical residents suggest that motivation, training, and longer experience living with restricted sleep do not make a person more resistant to the negative performance effects of sleep deprivation (Arnedt, Owens, Crouch, Stahl, & Carskadon, 2005; Van Dongen, Maislin, Mullington, & Dinges, 2003).

Sleep deprivation impairs many types of performance: reduces the ability to concentrate; slows reaction time; and reduces the ability to remember and learn new facts and motor skills (Goel et al., 2009). Sleep deprivation also leads to irritability, bad mood, reduced communication skills and ability to cope with the emotional demands of the workplace. Researchers report that decision-making ability deteriorates and risk-taking behavior increases (Killgore, Grugle, & Balkin, 2012). In addition, reduced situation awareness can impair a nurse's ability to respond to patient care needs. Importantly, researchers found that sleep-deprived participants did not recognize how poorly they were performing: they tended to think they were doing better than they were (Van Dongen & Belenky, 2009).

## Worker injury and errors

Declines in neurocognitive performance lead to higher rates of fatigue-related injury and worker errors. Certain work scheduling patterns are associated with higher risks for errors and incidents (Folkard & Lombardi, 2006). Compared with day shifts, risks are 15% higher for evening shifts and 28% higher for night shifts. When compared with 8-hour shifts, 10-hour shifts increased the risk by 13%, and 12-hour shifts increased risk by 28%. Risk increased by 17% for the third consecutive night shift and 36% for the fourth. Dembe, Erickson, Delbos, & Banks (2005) reported a possible "dose response": as weekly work hours increased, injury rates to workers increased. Folkard and Lombardi estimate the highest risk is for shifts that are both long and during the night.

Only a few studies have examined mandatory overtime but the evidence available raises concern. Studies of nurses linked mandatory overtime with higher risk for needlestick and other work-related injury, work-related illness, and missing more than 2 days of work because of these (de Castro et al., 2010; Gershon et al., 2009).

### **Poor health behaviors and metabolic syndrome**

Research links shift work and long work hours to poor health behaviors. Bushnell, Colombi, Caruso, & Tak (2010) found that of nine work schedule patterns, the pattern predicted to be the most difficult (12-hour rotation shifts) was associated with the high rates of all five poor health behaviors examined: short sleep duration, smoking, obesity, low physical activity, and higher alcohol use. A systematic review by Frost, Kolstad, & Bonde (2009) concluded that smoking was generally more frequent in shift workers. Workers may be drawn to smoking as a countermeasure for the fatigue, sleepiness, and stress linked to demanding work schedules. A review of obesity and shift work by Antunes, Levandovski, Dantas, Caumo, & Hidalgo (2010) concludes that there is considerable epidemiological evidence for a link between shift work and obesity. A few studies have examined long work hours and obesity, and some report a significant positive relationship (Nakamura et al., 1998; Shields, 1999). The mechanism could be through poor-quality sleep and short sleep duration promoting an imbalance in appetite hormones that increase feelings of hunger and metabolic changes leading to obesity, insulin resistance, and reduced lipid tolerance (Spiegel, Tasali, Leproult, & Van Cauter, 2009).

### **Diseases associated with shift work**

Along with sleepiness, disturbances to sleep also promote mood disturbances and gastrointestinal complaints (Sack, 2010). Shift workers commonly complain of gastrointestinal symptoms including abdominal pain, gas, diarrhea, constipation, nausea, vomiting, change in appetite, indigestion, and heartburn (Caruso, Lusk, & Gillespie, 2004; Knutsson & Boggild, 2010). A variety of stomach and digestive disorders have been reported in shift workers. Schernhammer et al. (2003) reported an increased risk of colon cancer in nurses working three or more nights per month for 15 or more years. Possible mechanisms for gastrointestinal disturbances include disturbances to circadian rhythms involved in eating and digesting food, sleep deprivation leading to stress response, changes in immune function, or the types of foods that are available during these shifts (Caruso et al., 2004; Knutsson & Boggild, 2010). Shift workers commonly report psychological complaints including bad mood, depression, irritability, anxiety, personality changes, and difficulty with personal relationships (Rohr, Von Essen, & Farr, 2003).

Shift workers are at somewhat higher risk for cardiovascular disorders. A review by Puttonen, Harma, & Hublin (2010) reported that epidemiologic data support a possible link with cardiovascular disease, including myocardial infarction, chest pain, and high blood pressure. Brown et al. (2010) recently reported an increased risk for ischemic strokes. The mechanism could be through metabolic changes and smoking. Frost et al. (2009) concluded there is evidence that workers are at higher risk for metabolic disturbances and increases in smoking after starting shift work.

In 2007, the International Agency for Research on Cancer of the World Health Organization announced sufficient evidence supports that shift work with circadian rhythm disruption is a probable carcinogen (Straif et al., 2007). They concluded data from human studies suggest the link is possible, while the evidence from animal studies is sufficient. The most studied cancer in shift workers is breast cancer. A meta-analysis of 13 studies reported night work increased cancer risk by 48% (Megdal, Kroenke, Laden, Pukkala, & Schernhammer, 2005). According to Bonde et al. (2012), statistically significant increases in risk for breast cancer were seen in people working night shift for 20 years or more, but it is not clear from existing studies whether or not there is risk for shorter durations. They also suggest that “women with previous or current breast cancer be advised not to work night shifts because of strong experimental evidence demonstrating accelerated tumor growth by suppression of melatonin secretion.” Two of three studies found an increase in prostate cancer (Conlon, Lightfoot, & Kreiger, 2007; Kubo et al., 2006; Schwartzbaum, Ahlbom, & Feychting, 2007). Costa, Haus, & Stevens (2010) discuss mechanisms that could possibly lead to cancer: repeated shifting of the times of sleep and activity; depression of immune surveillance; and exposure to light at night reducing melatonin levels.

Shift work is associated with adverse reproductive outcomes for women. A review by Frazier & Grainger (2003) reported that shift work was associated with a modest increase in spontaneous abortion, preterm birth, and reduced ability for women to become pregnant. Shechter, James, & Boivin (2008) reported increased medical office visits for menstrual cycle symptoms and cycles that were shorter, longer, or irregular.

People with chronic diseases may have more difficulty managing their symptoms and disease progression when working shift work. Some medications have changes in effectiveness across the circadian cycle which can lead to more difficulty in determining the appropriate dosage and time for administration when working rotating or night shifts. Shift work can also interfere with treatment regimens that involve maintaining regular sleep times, avoiding sleep deprivation, and controlling amounts and times of meals and exercise. Sood (2003) warned that the following conditions could get worse with shift work: heart disease, high blood pressure, stomach and intestinal disorders, sleep disorders, insulin-dependent diabetes, seizure disorders, asthma requiring medication, psychiatric diseases, and alcohol/drug abuse.

Some people have many symptoms, do not get relief from medical therapies, and appear unable to adjust to shift work. Harrington (2001) estimated 20% of workers who tried shift work were unable to adjust. The reason is not fully understood, but it may be due to an inherited tendency to be more sensitive to sleep loss and disruption of circadian rhythms (Van Dongen & Belenky, 2009).

## **Diseases associated with long work hours**

The number of studies examining the influence of long work hours on health has been growing recently but is substantially less than the number examining shift work. Several studies report long work hours are associated with increased fatigue, poor mood, poor recovery from work, and reduced perceived health (Caruso, Hitchcock, Dick, Russo, &

Schmit, 2004; Ettner & Grzywacz, 2001; van der Hulst, van Veldhoven, & Beckers, 2006; Siu & Donald, 1995). Men and women working long hours showed higher prevalence of depression and anxiety disorders (Kleppa, Sanne, & Tell, 2008). Similarly, Suwazono et al. (2007) reported better scores for depression and fatigue when work hours were less than 12 per day and 58 per week. Long work hours increase the risk for coronary artery disease by 40% according to a recent systematic review and meta-analysis by Virtanen et al. (2012). Long hours are associated with high blood pressure and diabetes, although not all studies consistently report a significant increase (Caruso et al., 2004). A systematic review identified eight studies with controls for physical demands that linked long work hours to adverse musculoskeletal disorders (Caruso & Waters, 2008). This may be due to longer exposure to physical demands and less recovery time between work periods. Possible adverse reproductive outcomes include increased time to pregnancy (Tuntiseranee, Olsen, Geater, & Kor-anantakul, 1998). A meta-analysis of six studies suggested a possible weak relationship with preterm birth (Mozurkewich, Luke, Avni, & Wolf, 2000).

### **Impacts to healthcare organizations and community**

Fatigue-related impairments can lead to reduced job performance which in turn negatively impacts the employer and the community. Evidence is growing that long work hours and shift work are linked to errors in the delivery of patient care and vehicle crashes on the commute.

Stimpfel, Sloane, & Aiken (2012) examined data from four states of nurses, their hospitals, and their patients. They reported that shifts greater than 13 hours were linked to patient dissatisfaction. Specifically, patients reported that nurses were not communicating well, their pain was not controlled, they did not get help as soon as they wanted, and they would not recommend the hospital. In this study, nurses with shifts greater than 13 hours had over double the risk for burnout and job dissatisfaction.

Rogers, Hwang, Scott, Aiken, & Dinges (2004) measured the working hours of nurses and their errors or near-errors during work. They found nurses had over three times the odds of making an error when working 12 or more hours, compared with 8.5-hour shifts. Similarly, Scott, Rogers, Hwang, & Zhang (2006) found the risk for patient care errors almost doubled when critical care nursing shifts lasted longer than 12.5 hours. In addition, more than 40 hours per week increased care errors by 46%.

Looking at fatal patient care outcomes, Trinkoff et al. (2011) reported hospitals with higher patient mortality rates had higher rates of nurses working long hours, having lack of time away from work, working while sick, and having higher weekly work hour burden. Staffing and hospital characteristics were controlled in the analysis indicating these schedule features had an independent impact on patient outcomes. Also, Landrigan et al. (2004) and Barger et al. (2006) found that interns working long hours made more serious medical errors.

Circadian, a shift work consulting company, estimates fatigue-related accidents, lost productivity, and health care expenses costs U.S. employers \$116.5 billion per year (Sirois, 2007). Similarly, Rosekind et al. (2010) estimated annual fatigue-related productivity losses for four employers were almost \$2000 per employee. As discussed above, shift work and

long work hours are linked to a wide range of health risks which can lead to more sickness absence. Employers are also at risk for loss of nurses because they become disabled and unable to work (Tüchsen, Christensen, Lund, & Feveile, 2008). This can lead to higher insurance and workers compensation rates. Failure to retain nurses increases costs for healthcare organizations: shift work and long work hours were major reasons for leaving the nursing profession in a study by Peter D. Hart Research Associates (2001). This remains a top concern according to a recent study by the American Nurses Association (2011): 74% of the nurses said their number one concern was acute or chronic effects of stress and overwork.

Fatigued nurses can endanger others during their commute home. Drowsy driving experts identify shift work and long work hours in the top five factors that increase risk for a motor vehicle crash (National Sleep Foundation, n.d.). An estimated 20% of all roadway crashes have been linked to drowsy driving (Connor et al., 2002; Horne & Reyner, 1995). A survey by the American Nurses Association (2011) found one in ten nurses reported a motor vehicle crash which they believed was related to fatigue or shift work.

## **U.S. National Public Health Goals and The Joint Commission**

Recognizing the widespread negative impacts of poor sleep on the person as well as the workplace and community, a growing number of organizations in the U.S. Department of Health and Human Services are targeting sleep health. These include the National Center for Sleep Disorders Research of the National Institutes of Health, Healthy People 2020, the Centers for Disease Control and Prevention (CDC), and National Institute for Occupational Safety and Health (NIOSH, an institute within the CDC). For decades, NIOSH has recognized shift work and work-related sleep loss to be a hazard in the workplace and has carried out an active research program to address this hazard. A goal of NIOSH's National Occupational Research Agenda (NORA, 2009) for Healthcare and Social Assistance is that healthcare organizations adopt best practices of scheduling and staffing that minimize excessive workload and other factors associated with fatigue.

Healthy People 2020 (2010) launched a new chapter for Sleep Health which includes three objectives for adults: (1) increase the proportion of adults who get 7 or more hours sleep per day; (2) increase the proportion of adults with sleep apnea who seek medical evaluation; (3) reduce the rate of vehicular crashes attributed to drowsy driving. Healthy People 2020 (2010) recognized that strong scientific evidence links sleep health to reduced rates of disease, injury, disability, and premature death.

Frieden (2010), the director of the CDC, identified "winnable battles" for meaningful and measurable progress over the next 1 to 4 years. Two are relevant to sleep: reducing obesity and motor vehicle crashes.

In December 2011, The Joint Commission issued a Sentinel Event Alert about healthcare worker fatigue and patient safety. In the alert, The Joint Commission gave suggestions for healthcare organizations to reduce risks from fatigue: examine and improve work schedules, staffing, hand-off process, and consider fatigue in all adverse events; involve staff in the design of their schedules; create and implement a fatigue management plan; educate staff

about sleep and fatigue; support staff who work long shifts; and for organizations who allow sleep breaks during the work shift, provide a good sleep environment and adequate release from work responsibilities.

## Concluding remarks

Substantial scientific evidence supports that insufficient sleep can lead to broad-reaching negative health and safety risks to the sleep-deprived person and can also endanger others around them. Employers, nurse managers, and staff nurses all share in the responsibility of adopting strategies to reduce these risks. The key strategy to reduce these risks is making sleep a priority in the employer's systems for organizing work and in the nurse's personal life.

To promote progress, education and training programs for managers and staff nurses would help increase their appreciation and knowledge about the demands of shift work, long work hours, and related sleep and fatigue issues. Strategies for managers and employers include improving the design of their work schedules, promoting frequent breaks during the work shift, fostering good co-worker and supervisor relationships, and establishing policies and systems that reduce the risk for fatigue and related health problems. In addition, workplaces can conduct periodic assessments to examine the influence of work schedules on factors on- and off-the-job including performance, alertness, sleep, unintentional injury, worker errors, near misses, illnesses, and off-the-job responsibilities. An anonymous, no-blame reporting system can collect reports on incident and near misses and assess if fatigue was a contributing factor. Strategies for staff nurses include allowing enough time for sleep and adopting good practices and behaviors to maximize sleep and alertness. Nurses with sleep problems or excessive work time sleepiness can seek an assessment and treatment from a sleep disorders specialist. Staff nurses also will benefit from educating their family about their special needs to gain their support and reduce conflicts between work and home demands.

NIOSH's topic page for work schedules has a number of resources to help nurses and healthcare organizations become better prepared to reduce risks linked to work-related sleep loss (<http://www.cdc.gov/niosh/topics/workschedules/>). NIOSH has drafted online training materials to help educate managers and nursing staff. When these are finalized, they will be available on this website.

## References

- American Nurses Association. 2011 ANA health & safety survey report. 2011 from <http://nursingworld.org/FunctionalMenuCategories/MediaResources/MediaBackgrounders/The-Nurse-Work-Environment-2011-Health-Safety-Survey.pdf>.
- Antunes LC, Levandovski R, Dantas G, Caumo W, Hidalgo MP. Obesity and shift work: chronobiological aspects. *Nutrition Research Reviews*. 2010; 23:155–168. [PubMed: 20122305]
- Arnedt JT, Owens J, Crouch M, Stahl J, Carskadon MA. Neurobehavioral performance of residents after heavy night call vs after alcohol ingestion. *Journal of the American Medical Association*. 2005; 294:1025–1033. [PubMed: 16145022]



- Barger LK, Ayas NT, Cade BE, Cronin JW, Rosner B, Speizer FE, et al. Impact of extended-duration shifts on medical errors, adverse events, and attentional failures. *PLoS Medicine*. 2006; 3:e487. [PubMed: 17194188]
- Basner M, Fomberstein K, Razavi FM, William J, Simpson N, Rosa RR, et al. American time use survey: sleep time and its relationship to waking activities. *Sleep*. 2007; 30:1085–1095. [PubMed: 17910380]
- Bonde JP, Hansen J, Kolstad HA, Mikkelsen S, Olsen JH, Blask DE, et al. Work at night and breast cancer – report on evidence-based options for preventive actions. *Scandinavian Journal of Work Environment and Health*. 2012; 38:380–390.
- Boyle LN, Tippin J, Paul A, Rizzo M. Driver performance in the moments surrounding a microsleep. *Transportation Research Part F: Traffic Psychology and Behaviour*. 2008; 11:126–136.
- Brown DL, Feskanich D, Sanchez BN, Rexrode KM, Schernhammer ES, Lisabeth LD. Rotating night shift work and the risk of ischemic stroke. *American Journal of Epidemiology*. 2010; 169:11370–1377.
- Bushnell PT, Colombi A, Caruso CC, Tak S. Work schedules and health behavior outcomes at a large manufacturer. *Industrial Health*. 2010; 48:395–405. [PubMed: 20720331]
- Caruso, CC.; Hitchcock, EM.; Dick, RB.; Russo, JM.; Schmit, JM. Cincinnati, OH: Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health; 2004. Overtime and extended work shifts: recent findings on illnesses, injuries, and health behaviors. DHHS (NIOSH) Publication No. 2004-143.
- Caruso CC, Lusk SL, Gillespie BW. Relationship of work schedules to gastrointestinal diagnoses, symptoms, and medication use in auto factory workers. *American Journal of Industrial Medicine*. 2004; 46:586–598. [PubMed: 15551368]
- Caruso CC, Waters TR. A review of work schedule issues and musculoskeletal disorders with an emphasis on the healthcare sector. *Industrial Health*. 2008; 46:523–534. [PubMed: 19088404]
- de Castro AB, Fujishiro K, Rue T, Tagalog EA, Samaco-Paquiz LP, Gee GC. Associations between work schedule characteristics and occupational injury and illness. *International Nursing Review*. 2010; 57:188–194. [PubMed: 20579153]
- Centers for Disease Control. Perceived insufficient rest or sleep among adults – United States, 2009. *Morbidity and Mortality Weekly Report*. 2008; 58(42):1175–1179.
- Centers for Disease Control. Effect of short sleep duration on daily activities – United States, 2005 – 2008. *Morbidity and Mortality Weekly Report*. 2011; 60:239–242. [PubMed: 21368739]
- Colten, HR.; Altevogt, BM. Sleep disorders and sleep deprivation an unmet public health problem. Washington, DC: National Academies; 2006.
- Conlon M, Lightfoot N, Kreiger N. Rotating shift work and risk of prostate cancer. *Epidemiology*. 2007; 18:182–183. [PubMed: 17179764]
- Connor J, Norton R, Ameratunga S, Robinson E, Civil I, Dunn R, et al. Driver sleepiness and risk of serious injury to car occupants: population based case control study. *BMJ*. 2002; 324(7346):1125. [PubMed: 12003884]
- Costa G, Haus E, Stevens R. Shift work and cancer – considerations on rationale, mechanisms, and epidemiology. *Scandinavian Journal of Work Environment and Health*. 2010; 36:163–179.
- Dembe AE, Erickson JB, Delbos RG, Banks SM. The impact of overtime and long work hours on occupational injuries and illnesses: new evidence from the United States. *Occupational and Environmental Medicine*. 2005; 62:588–597. [PubMed: 16109814]
- Drake CL, Roehrs T, Richardson G, Walsh JK, Roth T. Shift work sleep disorder: prevalence and consequences beyond that of symptomatic day workers. *Sleep*. 2004; 27:1453–1462. [PubMed: 15683134]
- Ettner SL, Grzywacz JG. Workers' perceptions of how jobs affect health: a social ecological perspective. *Journal of Occupational Health Psychology*. 2001; 6:101–113. [PubMed: 11326723]
- Everson, CA. Comparative research approaches to discovering the biomedical implications of sleep loss and sleep recovery. In: Amlaner, CJ.; Phil, D.; Fuller, PM., editors. *Basics of Sleep Guide*. 2nd ed.. Westchester, IL: Sleep Research Society; 2009. p. 237-248.

- Folkard S, Lombardi DA. Modeling the impact of the components of long work hours on injuries and “accidents”. *American Journal of Industrial Medicine*. 2006; 49:953–963. [PubMed: 16570251]
- Frazier LM, Grainger DA. Shift work and adverse reproductive outcomes among men and women. *Clinics in Occupational and Environmental Medicine*. 2003; 3:279–292.
- Frieden, TR. [Retrieved April 9, 2013] Winnable battles. 2010. from <http://www.cdc.gov/winnablebattles/FocusAreas.html>
- Frost P, Kolstad HA, Bonde JP. Shift work and the risk of ischemic heart disease – a systematic review of the epidemiologic evidence. *Scandinavian Journal of Work Environment and Health*. 2009; 35:163–179.
- Geiger-Brown J, Rogers VE, Trinkoff AM, Kane RL, Bausell RB, Scharf SM. Sleep, sleepiness, fatigue, and performance of 12-hour-shift nurses. *Chronobiology International*. 2012; 29(2):211–219. [PubMed: 22324559]
- Gershon RRM, Pearson JM, Sherman MF, Samar SM, Canton AN, Stone PW. The prevalence and risk factors for percutaneous injuries in registered nurses in the home health care sector. *American Journal of Infection Control*. 2009; 37:525–533. [PubMed: 19216006]
- Goel N, Rao H, Durmer JS, Dinges DF. Neurocognitive consequences of sleep deprivation. *Seminars in Neurology*. 2009; 29:320–339. [PubMed: 19742409]
- Harrington JM. Health effects of shift work and extended hours of work. *Occupational and Environmental Medicine*. 2001; 58:68–72.
- Hayashi T, Kobayashi Y, Yamaoka K, Yano E. Effect of overtime work on 24-h ambulatory blood pressure. *Journal of Occupational and Environmental Medicine*. 1996; 38:1007–1011. [PubMed: 8899576]
- Healthy People 2020. [Retrieved April 9, 2013] Sleep Health. 2010. from <http://www.healthypeople.gov/2020/topicsobjectives2020/objectiveslist.aspx?topicid=38>
- Horne JA, Reyner LA. Sleep-related vehicle accidents. *BMJ*. 1995; 310:565–567. [PubMed: 7888930]
- van der Hulst M, van Veldhoven M, Beckers D. Overtime and need for recovery in relation to job demands and job control. *Journal of Occupational Health*. 2006; 48:11–19. [PubMed: 16484758]
- Ingre M, Åkerstedt T. Effect of accumulated night work during the working lifetime, on subjective health and sleep in monozygotic twins. *Journal of Sleep Research*. 2004; 13:45–48. [PubMed: 14996034]
- Killgore WD, Grugle NL, Balkin TJ. Gambling when sleep deprived: don’t bet on stimulants. *Chronobiology International*. 2012; 29:43–54. [PubMed: 22217100]
- Kleppa E, Sanne B, Tell GS. Working overtime is associated with anxiety and depression: the Hordaland health study. *Journal of Occupational and Environmental Medicine*. 2008; 50:658–666. [PubMed: 18545093]
- Knauth P. Extended work periods. *Industrial Health*. 2007; 45:125–136. [PubMed: 17284884]
- Knutsson A, Boggild H. Gastrointestinal disorders among shift workers. *Scandinavian Journal of Work Environment and Health*. 2010; 36:85–95.
- Kubo T, Ozasa K, Mikami K, Wakai K, Fujino Y, Watanabe Y, et al. Prospective cohort study of the risk of prostate cancer among rotating-shift workers: findings from the Japan collaborative cohort study. *American Journal of Epidemiology*. 2006; 164:549–555. [PubMed: 16829554]
- Landrigan CP, Rothschild JM, Cronin JW, Kaushal R, Burdick E, Katz JT, et al. Effect of reducing interns’ work hours on serious medical errors in intensive care units. *New England Journal of Medicine*. 2004; 351:1838–1848. [PubMed: 15509817]
- Luckhaupt SE, Tak SWZ, Calvert GM. The prevalence of short sleep duration by industry and occupation in the National Health Interview Survey. *Sleep*. 2010; 33:149–159. [PubMed: 20175398]
- Megdal SP, Kroenke CH, Laden F, Pukkala E, Schernhammer ES. Night work and breast cancer risk: a systematic review and meta-analysis. *European Journal of Cancer*. 2005; 41:2023–2032. [PubMed: 16084719]
- Mozurkewich EL, Luke B, Avni M, Wolf FM. Working conditions and adverse pregnancy outcome: a meta-analysis. *Obstetrics and Gynecology*. 2000; 95:623–635. [PubMed: 10725502]

- Nakamura K, Shimai S, Kikuchi S, Takahashi H, Tanaka M, Nakano S, et al. Increases in body mass index and waist circumference as outcomes of working overtime. *Occupational Medicine (London)*. 1998; 48:169–173.
- National Sleep Foundation. Summary of findings of 2007 sleep in America poll. 2007 from [http://www.sleepfoundation.org/sites/default/files/Summary\\_Of\\_Findings%20-%20FINAL.pdf](http://www.sleepfoundation.org/sites/default/files/Summary_Of_Findings%20-%20FINAL.pdf).
- National Sleep Foundation. (n.d.). Who is at risk? from <http://drowsydriving.org/about/%20whos-at-risk/>.
- NORA Healthcare and Social Assistance Sector Council. [Retrieved April 9, 2013] State of the sector healthcare and social assistance. Department of Health and Human Services, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health Publication No. 2009-139. 2009. from <http://www.cdc.gov/niosh/docs/2009-139/pdfs/2009-139.pdf>
- Peter D. Hart Research Associates. The nurse shortage: perspectives from current direct care nurses and former direct care nurses. 2001 from [http://www.aft.org/pubs-reports/healthcare/Hart\\_Report.pdf](http://www.aft.org/pubs-reports/healthcare/Hart_Report.pdf).
- Pilcher JJ, Lambert BJ, Huffcutt AI. Differential effects of permanent and rotating shifts on self-report sleep length: a meta-analytic review. *Sleep*. 2000; 23:155–163. [PubMed: 10737332]
- Puttonen S, Harma M, Hublin C. Shift work and cardiovascular disease – pathways from circadian stress to morbidity. *Scandinavian Journal of Work Environment and Health*. 2010; 36:96–108.
- Rogers AE, Hwang W, Scott LD, Aiken LH, Dinges DF. The working hours of hospital staff nurses and patient safety. *Health Affairs*. 2004; 23:202–212. [PubMed: 15318582]
- Rohr SM, Von Essen SG, Farr LA. Overview of the medical consequences of shift work. *Clinics in Occupational and Environmental Medicine*. 2003; 3:351–361.
- Rosekind MR, Gregory KB, Mallis MM, Brandt SL, Seal B, Lerner D. The cost of poor sleep: workplace productivity loss and associated costs. *Journal of Occupational & Environmental Medicine*. 2010; 52:91–98. [PubMed: 20042880]
- Sack RL. Jet lag. *New England Journal of Medicine*. 2010; 362:440–447. [PubMed: 20130253]
- Sasaki T, Iwasaki K, Oka T, Hisanaga N, Ueda T, Takada Y. Effect of working hours on cardiovascular-autonomic nervous functions in engineers in an electronics manufacturing company. *Industrial Health*. 1999; 37:55–61. [PubMed: 10052300]
- Schernhammer ES, Laden F, Speizer FE, Willett WC, Hunter DJ, Kawachi I, et al. Night-shift work and risk of colorectal cancer in the nurses' health study. *Journal of the National Cancer Institute*. 2003; 95:825–828. [PubMed: 12783938]
- Schwartz JR, Roth T. Neurophysiology of sleep and wakefulness: basic science and clinical implications. *Current Neuropharmacology*. 2008; 64:367–378. [PubMed: 19587857]
- Schwartzbaum J, Ahlbom A, Feychting M. Cohort study of cancer risk among male and female shift workers. *Scandinavian Journal of Work Environment and Health*. 2007; 33:336–343.
- Scott LD, Rogers AE, Hwang WT, Zhang Y. Effects of critical care nurses' work hours on vigilance and patients' safety. *American Journal of Critical Care*. 2006; 15:30–37. [PubMed: 16391312]
- Shechter A, James FO, Boivin DB. Circadian rhythms and shift working women. *Sleep Medicine Clinics*. 2008; 3:13–24.
- Shields M. Long working hours and health. *Health Reports*. 1999; 11:33–48. [PubMed: 10618741]
- Sirois WG. The secret cost of fatigue. 2007 from <http://www.taproot.com/archives/445>.
- Siu O-L, Donald I. Psychosocial factors at work and workers' health in Hong Kong: an exploratory study. *Bulletin of the Hong Kong Psychological Society*. 1995; 34/35:30–56.
- Smolensky MH, Di Milia L, Ohayon MM, Philip P. Sleep disorders, medical conditions, and road accident risk. *Accident Analysis and Prevention*. 2011; 43:533–548. [PubMed: 21130215]
- Sood A. Medical screening and surveillance of shift workers for health problems. *Clinics in Occupational and Environmental Medicine*. 2003; 3:339–349.
- Spiegel K, Tasali E, Leproult R, Van Cauter E. Effects of poor and short sleep on glucose metabolism and obesity risk. *Nature Reviews Endocrinology*. 2009; 5:253–261.
- Stimpfel AW, Sloane DM, Aiken LH. The longer the shifts for hospital nurses, the higher the levels of burnout and patient dissatisfaction. *Health Affairs (Millwood)*. 2012; 31:2501–2509.

- Straif K, Baan R, Grosse Y, Secretan B, Ghissassi FR, Bouvard V, et al. WHO International Agency for Research on Cancer Monograph Working Group. Carcinogenicity of shift-work, painting, and fire-fighting. *Lancet Oncology*. 2007; 8:1065–1066. [PubMed: 19271347]
- Suwazono Y, Nagashima S, Okubo Y, Uetani M, Kobayashi E, Kido T, et al. Estimation of the number of working hours critical for the development of mental and physical fatigue symptoms in Japanese male workers – Application of benchmark dose method. *American Journal Of Industrial Medicine*. 2007; 50:173–182. [PubMed: 17315176]
- The Joint Commission. Health care worker fatigue and patient safety. The Joint Commission Sentinel Event Alert. 2011; 48:1–4. from [http://www.jointcommission.org/assets/1/18/sea\\_48.pdf](http://www.jointcommission.org/assets/1/18/sea_48.pdf).
- Trinkoff AM, Johantgen M, Storr CL, Gurses AP, Liang Y, Han K. Nurses' work schedule characteristics, nurse staffing, and patient mortality. *Nursing Research*. 2011; 60:1–8. [PubMed: 21127449]
- Tüchsen F, Christensen KB, Lund T, Feveile H. A 15-year prospective study of shift work and disability pension. *Occupational & Environmental Medicine*. 2008; 65:283–285. [PubMed: 18198201]
- Tuntiseranee P, Olsen J, Geater A, Kor-anantakul O. Are long working hours and shiftwork risk factors for subfecundity? A study among couples from southern Thailand. *Occupational & Environmental Medicine*. 1998; 55:99–105. [PubMed: 9614393]
- Van Dongen HPA, Belenky G. Individual differences in vulnerability to sleep loss in the work environment. *Industrial Health*. 2009; 47:518–526. [PubMed: 19834261]
- Van Dongen HP, Maislin G, Mullington JM, Dinges DF. The cumulative cost of additional wakefulness: dose-response effects on neurobehavioral functions and sleep physiology from chronic sleep restriction and total sleep deprivation. *Sleep*. 2003; 26:117–126. [PubMed: 12683469]
- Virtanen M, Heikkilä K, Jokela M, Ferrie JE, Batty FG, Vahtera J, et al. Long working hours and coronary heart disease: a systematic review and meta-analysis. *American Journal of Epidemiology*. 2012; 176(7):586–596. [PubMed: 22952309]

### Key Practice Points

- Shift work and long work hours put nurses at risk for short sleep duration and sleep disturbances.
- Insufficient sleep increases the risk for reduced performance on the job, injuries, obesity, a wide range of chronic diseases, and fatigue-related errors which could harm patients and other people.
- Education and training programs for managers and staff nurses would help increase their appreciation and knowledge about the demands of shift work and long work hours and provide them with strategies to reduce these risks.
- Recognizing the widespread negative impacts of insufficient sleep on the person as well as the workplace and community, a growing number of U.S. public health organizations target improving sleep health including The Joint Commission and several organizations in the U.S. Department of Health and Human Services.