



## Archived Editions (COVID-19 Genomics and Precision Public Health Weekly Update)

Published on 01/14/2021

### COVID-19 Genomics and Precision Public Health Weekly Update Content

- Pathogen and Human Genomics Studies
- Non-Genomics Precision Health Studies
- News, Reviews and Commentaries

#### Pathogen and Human Genomics Studies

- Neutralization of N501Y mutant SARS-CoV-2 by BNT162b2 vaccine-elicited sera  
(/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=571)  
X Xie et al, BIORXIV, January 7, 2021

Rapidly spreading variants of SARS-CoV-2 that have arisen in the United Kingdom and South Africa share the spike N501Y substitution, which is of particular concern because it is located in the viral receptor binding site for cell entry and increases binding to the receptor (angiotensin converting enzyme 2). We generated isogenic N501 and Y501 SARS-CoV-2. Sera of 20 participants in a previously reported trial of the mRNA-based COVID-19 vaccine BNT162b2 had equivalent neutralizing titers to the N501 and Y501 viruses.

- Establishment and lineage dynamics of the SARS-CoV-2 epidemic in the UK  
(/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=576)  
L du Plessis et al, Science, January 8, 2021

Using large-scale phylogenetic analyses, combined with epidemiological and travel data, we quantify the size, spatio-temporal origins and persistence of genetically-distinct UK transmission lineages. Rapid fluctuations in virus importation rates resulted in >1000 lineages; those introduced prior to national lockdown tended to be larger and more dispersed. Lineage importation and regional lineage diversity declined after lockdown, while lineage elimination was size-dependent.

- Utilization of Whole Genome Sequencing to Understand SARS-CoV-2 Transmission Dynamics in Long-Term Care Facilities, Correctional Facilities and Meat Processing Plants in Minnesota, March – June 2020 (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=580)  
NB Lehnertz et al, MEDRXIV, January 8, 2021

Whole genome sequencing (WGS) was performed from 8 selected outbreaks in Minnesota from March–June, 2020. WGS and phylogenetic analysis was conducted on 319 samples, constituting 14.4% of the 2,222 total positive individuals associated with these outbreaks. Data suggest that a single introduction into a facility can result in a widespread outbreak, and early identification and

cohorting of cases, along with infection prevention and control measures is imperative.

- First genome sequencing of SARS-CoV-2 recovered from an infected cat and its owner in Latin America. (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=585)  
Carlos Renata Santiago Alberto et al. Transboundary and emerging diseases 2021 Jan
- Confirmed Reinfection with SARS-CoV-2 Variant VOC-202012/01. (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=586)  
Harrington David et al. Clinical infectious diseases : an official publication of the Infectious Diseases Society of America 2021 Jan
- Saliva viral load is a dynamic unifying correlate of COVID-19 severity and mortality (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=587)  
J Silva et al, MEDRXIV, January 10, 2021

We examine saliva and nasopharyngeal viral load over time and correlate them with patient demographics, and cellular and immune profiling. We found that saliva viral load was significantly higher in those with COVID-19 risk factors; that it correlated with increasing levels of disease severity and showed a superior ability over nasopharyngeal viral load as a predictor of mortality over time.

- The Sensitivity and Costs of Testing for SARS-CoV-2 Infection With Saliva Versus Nasopharyngeal Swabs- A Systematic Review and Meta-analysis (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=592)  
ML Bastos et al, Annals of Internal Medicine, January 12, 2021

Nasopharyngeal swabs are the primary sampling method used for detection of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), but they require a trained health care professional and extensive personal protective equipment. This systematic review shows that saliva sampling seems to be a similarly sensitive and less costly alternative that could replace nasopharyngeal swabs for collection of clinical samples for SARS-CoV-2 testing.

- Gut microbiota composition reflects disease severity and dysfunctional immune responses in patients with COVID-19 (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=593)  
YK Yeoh et al, BMJ Gut, January 11, 2021

Associations between gut microbiota composition, levels of cytokines and inflammatory markers in patients with COVID-19 suggest that the gut microbiome is involved in the magnitude of COVID-19 severity possibly via modulating host immune responses. Furthermore, the gut microbiota dysbiosis after disease resolution could contribute to persistent symptoms, highlighting a need to understand how gut microorganisms are involved in inflammation and COVID-19.

- Statistical power in COVID-19 case-control host genomic study design. (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=600)  
Lin Yu-Chung et al. Genome medicine 2020 12 (1) 115

" The field of infectious disease genetic epidemiology has long described the unique and complex implications that a requirement of exposure to the infectious agent imposes on study design. Here, we draw on current estimates of exposure, infectivity, and test accuracy of COVID-19 to demonstrate the feasibility of detecting host genetic factors associated with COVID-19 susceptibility and severity with current study designs. We demonstrate why studying susceptibility to SARS-CoV-2 infection could be futile at certain stages of the pandemic. Our insights can aid in the interpretation of genetic findings emerging from the literature and guide the design of future host genetic studies."

- Establishment and lineage dynamics of the SARS-CoV-2 epidemic in the UK.

(/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=601)

du Plessis Louis et al. Science (New York, N.Y.) 2021 Jan

Abstract: The UK's COVID-19 epidemic during early 2020 was one of world's largest and unusually well represented by virus genomic sampling. Here we reveal the fine-scale genetic lineage structure of this epidemic through analysis of 50,887 SARS-CoV-2 genomes, including 26,181 from the UK sampled throughout the country's first wave of infection. Using large-scale phylogenetic analyses, combined with epidemiological and travel data, we quantify the size, spatio-temporal origins and persistence of genetically-distinct UK transmission lineages. Rapid fluctuations in virus importation rates resulted in >1000 lineages; those introduced prior to national lockdown tended to be larger and more dispersed. Lineage importation and regional lineage diversity declined after lockdown, while lineage elimination was size-dependent. We discuss the implications of our genetic perspective on transmission dynamics for COVID-19 epidemiology and control.

## Non-Genomics Precision Health Studies

- Neutralization of N501Y mutant SARS-CoV-2 by BNT162b2 vaccine-elicited sera

(/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=571)

X Xie et al, BIORXIV, January 7, 2021

Rapidly spreading variants of SARS-CoV-2 that have arisen in the United Kingdom and South Africa share the spike N501Y substitution, which is of particular concern because it is located in the viral receptor binding site for cell entry and increases binding to the receptor (angiotensin converting enzyme 2). We generated isogenic N501 and Y501 SARS-CoV-2. Sera of 20 participants in a previously reported trial of the mRNA-based COVID-19 vaccine BNT162b2 had equivalent neutralizing titers to the N501 and Y501 viruses.

- Establishment and lineage dynamics of the SARS-CoV-2 epidemic in the UK

(/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=576)

L du Plessis et al, Science, January 8, 2021

Using large-scale phylogenetic analyses, combined with epidemiological and travel data, we quantify the size, spatio-temporal origins and persistence of genetically-distinct UK transmission lineages. Rapid fluctuations in virus importation rates resulted in >1000 lineages; those introduced prior to national lockdown tended to be larger and more dispersed. Lineage importation and regional lineage diversity declined after lockdown, while lineage elimination was size-dependent.

- Utilization of Whole Genome Sequencing to Understand SARS-CoV-2 Transmission Dynamics in Long-Term Care Facilities, Correctional Facilities and Meat Processing Plants in Minnesota, March – June 2020 (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=580)

NB Lehnertz et al, MEDRXIV, January 8, 2021

Whole genome sequencing (WGS) was performed from 8 selected outbreaks in Minnesota from March–June, 2020. WGS and phylogenetic analysis was conducted on 319 samples, constituting 14.4% of the 2,222 total positive individuals associated with these outbreaks. Data suggest that a single introduction into a facility can result in a widespread outbreak, and early identification and cohorting of cases, along with infection prevention and control measures is imperative.

- First genome sequencing of SARS-CoV-2 recovered from an infected cat and its owner in Latin America. (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=585)

Carlos Renata Santiago Alberto et al. Transboundary and emerging diseases 2021 Jan

- Confirmed Reinfection with SARS-CoV-2 Variant VOC-202012/01. (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=586)

Harrington David et al. Clinical infectious diseases : an official publication of the Infectious Diseases Society of America 2021 Jan

- Saliva viral load is a dynamic unifying correlate of COVID-19 severity and mortality (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=587)

J Silva et al, MEDRXIV, January 10, 2021

We examine saliva and nasopharyngeal viral load over time and correlate them with patient demographics, and cellular and immune profiling. We found that saliva viral load was significantly higher in those with COVID-19 risk factors; that it correlated with increasing levels of disease severity and showed a superior ability over nasopharyngeal viral load as a predictor of mortality over time.

- The Sensitivity and Costs of Testing for SARS-CoV-2 Infection With Saliva Versus Nasopharyngeal Swabs- A Systematic Review and Meta-analysis (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=592)

ML Bastos et al, Annals of Internal Medicine, January 12, 2021

Nasopharyngeal swabs are the primary sampling method used for detection of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), but they require a trained health care professional and extensive personal protective equipment. This systematic review shows that saliva sampling seems to be a similarly sensitive and less costly alternative that could replace nasopharyngeal swabs for collection of clinical samples for SARS-CoV-2 testing.

- Gut microbiota composition reflects disease severity and dysfunctional immune responses in patients with COVID-19 (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=593)

YK Yeoh et al, BMJ Gut, January 11, 2021

Associations between gut microbiota composition, levels of cytokines and inflammatory markers in patients with COVID-19 suggest that the gut microbiome is involved in the magnitude of COVID-19 severity possibly via modulating host immune responses. Furthermore, the gut microbiota dysbiosis after disease resolution could contribute to persistent symptoms, highlighting a need to understand how gut microorganisms are involved in inflammation and COVID-19.

- Statistical power in COVID-19 case-control host genomic study design. (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=600)

Lin Yu-Chung et al. Genome medicine 2020 12 (1) 115

" The field of infectious disease genetic epidemiology has long described the unique and complex implications that a requirement of exposure to the infectious agent imposes on study design. Here, we draw on current estimates of exposure, infectivity, and test accuracy of COVID-19 to demonstrate the feasibility of detecting host genetic factors associated with COVID-19 susceptibility and severity with current study designs. We demonstrate why studying susceptibility to SARS-CoV-2 infection could be futile at certain stages of the pandemic. Our insights can aid in the interpretation of genetic findings emerging from the literature and guide the

design of future host genetic studies."

- Establishment and lineage dynamics of the SARS-CoV-2 epidemic in the UK.

(/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=601)

du Plessis Louis et al. Science (New York, N.Y.) 2021 Jan

Abstract: The UK's COVID-19 epidemic during early 2020 was one of world's largest and unusually well represented by virus genomic sampling. Here we reveal the fine-scale genetic lineage structure of this epidemic through analysis of 50,887 SARS-CoV-2 genomes, including 26,181 from the UK sampled throughout the country's first wave of infection. Using large-scale phylogenetic analyses, combined with epidemiological and travel data, we quantify the size, spatio-temporal origins and persistence of genetically-distinct UK transmission lineages. Rapid fluctuations in virus importation rates resulted in >1000 lineages; those introduced prior to national lockdown tended to be larger and more dispersed. Lineage importation and regional lineage diversity declined after lockdown, while lineage elimination was size-dependent. We discuss the implications of our genetic perspective on transmission dynamics for COVID-19 epidemiology and control.

## News, Reviews and Commentaries

- Neutralization of N501Y mutant SARS-CoV-2 by BNT162b2 vaccine-elicited sera

(/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=571)

X Xie et al, BIORXIV, January 7, 2021

Rapidly spreading variants of SARS-CoV-2 that have arisen in the United Kingdom and South Africa share the spike N501Y substitution, which is of particular concern because it is located in the viral receptor binding site for cell entry and increases binding to the receptor (angiotensin converting enzyme 2). We generated isogenic N501 and Y501 SARS-CoV-2. Sera of 20 participants in a previously reported trial of the mRNA-based COVID-19 vaccine BNT162b2 had equivalent neutralizing titers to the N501 and Y501 viruses.

- Establishment and lineage dynamics of the SARS-CoV-2 epidemic in the UK

(/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=576)

L du Plessis et al, Science, January 8, 2021

Using large-scale phylogenetic analyses, combined with epidemiological and travel data, we quantify the size, spatio-temporal origins and persistence of genetically-distinct UK transmission lineages. Rapid fluctuations in virus importation rates resulted in >1000 lineages; those introduced prior to national lockdown tended to be larger and more dispersed. Lineage importation and regional lineage diversity declined after lockdown, while lineage elimination was size-dependent.

- Utilization of Whole Genome Sequencing to Understand SARS-CoV-2 Transmission Dynamics in Long-Term Care Facilities, Correctional Facilities and Meat Processing Plants in Minnesota, March – June 2020 (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=580)

NB Lehnertz et al, MEDRXIV, January 8, 2021

Whole genome sequencing (WGS) was performed from 8 selected outbreaks in Minnesota from March–June, 2020. WGS and phylogenetic analysis was conducted on 319 samples, constituting 14.4% of the 2,222 total positive individuals associated with these outbreaks. Data suggest that a single introduction into a facility can result in a widespread outbreak, and early identification and

cohorting of cases, along with infection prevention and control measures is imperative.

- First genome sequencing of SARS-CoV-2 recovered from an infected cat and its owner in Latin America. (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=585)

Carlos Renata Santiago Alberto et al. Transboundary and emerging diseases 2021 Jan

- Confirmed Reinfection with SARS-CoV-2 Variant VOC-202012/01. (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=586)

Harrington David et al. Clinical infectious diseases : an official publication of the Infectious Diseases Society of America 2021 Jan

- Saliva viral load is a dynamic unifying correlate of COVID-19 severity and mortality (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=587)

J Silva et al, MEDRXIV, January 10, 2021

We examine saliva and nasopharyngeal viral load over time and correlate them with patient demographics, and cellular and immune profiling. We found that saliva viral load was significantly higher in those with COVID-19 risk factors; that it correlated with increasing levels of disease severity and showed a superior ability over nasopharyngeal viral load as a predictor of mortality over time.

- The Sensitivity and Costs of Testing for SARS-CoV-2 Infection With Saliva Versus Nasopharyngeal Swabs- A Systematic Review and Meta-analysis (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=592)

ML Bastos et al, Annals of Internal Medicine, January 12, 2021

Nasopharyngeal swabs are the primary sampling method used for detection of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), but they require a trained health care professional and extensive personal protective equipment. This systematic review shows that saliva sampling seems to be a similarly sensitive and less costly alternative that could replace nasopharyngeal swabs for collection of clinical samples for SARS-CoV-2 testing.

- Gut microbiota composition reflects disease severity and dysfunctional immune responses in patients with COVID-19 (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=593)

YK Yeoh et al, BMJ Gut, January 11, 2021

Associations between gut microbiota composition, levels of cytokines and inflammatory markers in patients with COVID-19 suggest that the gut microbiome is involved in the magnitude of COVID-19 severity possibly via modulating host immune responses. Furthermore, the gut microbiota dysbiosis after disease resolution could contribute to persistent symptoms, highlighting a need to understand how gut microorganisms are involved in inflammation and COVID-19.

- Statistical power in COVID-19 case-control host genomic study design. (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=600)

Lin Yu-Chung et al. Genome medicine 2020 12 (1) 115

" The field of infectious disease genetic epidemiology has long described the unique and complex implications that a requirement of exposure to the infectious agent imposes on study design. Here, we draw on current estimates of exposure, infectivity, and test accuracy of COVID-19 to demonstrate the feasibility of detecting host genetic factors associated with COVID-19 susceptibility and severity with current study designs. We demonstrate why studying susceptibility to SARS-CoV-2 infection could be futile at certain stages of the pandemic. Our

insights can aid in the interpretation of genetic findings emerging from the literature and guide the design of future host genetic studies."

- Establishment and lineage dynamics of the SARS-CoV-2 epidemic in the UK.

(/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=601)

du Plessis Louis et al. Science (New York, N.Y.) 2021 Jan

Abstract: The UK's COVID-19 epidemic during early 2020 was one of world's largest and unusually well represented by virus genomic sampling. Here we reveal the fine-scale genetic lineage structure of this epidemic through analysis of 50,887 SARS-CoV-2 genomes, including 26,181 from the UK sampled throughout the country's first wave of infection. Using large-scale phylogenetic analyses, combined with epidemiological and travel data, we quantify the size, spatio-temporal origins and persistence of genetically-distinct UK transmission lineages. Rapid fluctuations in virus importation rates resulted in >1000 lineages; those introduced prior to national lockdown tended to be larger and more dispersed. Lineage importation and regional lineage diversity declined after lockdown, while lineage elimination was size-dependent. We discuss the implications of our genetic perspective on transmission dynamics for COVID-19 epidemiology and control.

**Disclaimer:** Articles listed in COVID-19 Genomics and Precision Public Health Weekly Update are selected by the CDC Office of Public Health Genomics to provide current awareness of the scientific literature and news. Inclusion in the update does not necessarily represent the views of the Centers for Disease Control and Prevention nor does it imply endorsement of the article's methods or findings. CDC and DHHS assume no responsibility for the factual accuracy of the items presented. The selection, omission, or content of items does not imply any endorsement or other position taken by CDC or DHHS. Opinion, findings and conclusions expressed by the original authors of items included in the Clips, or persons quoted therein, are strictly their own and are in no way meant to represent the opinion or views of CDC or DHHS. References to publications, news sources, and non-CDC Websites are provided solely for informational purposes and do not imply endorsement by CDC or DHHS.

Page last reviewed: Oct 1, 2020

Page last updated: Jan 20, 2021

Content source: Office of Genomics and Precision Public Health (<http://www.cdc.gov/genomics/>), CDC Office of Science (<https://www.cdc.gov/od/science/index.htm>)