



## COVID-19

WEAR A MASK

STAY 6 FEET APART

AVOID CROWDS

# Information About Johnson & Johnson's Janssen COVID-19 Vaccine

Updated Mar. 4, 2021

[Print](#)

## General information

**Name:** JNJ-78436735

**Manufacturer:** Janssen Pharmaceuticals Companies of Johnson & Johnson

**Type of vaccine:** Viral vector

Learn more about [how COVID-19 vaccines work](#) and get a better [understanding of COVID-19 viral vector vaccines](#).

**Number of shots:** 1 shot

**How given:** Shot in the muscle of the upper arm

### Does not contain:

- Eggs
- Preservatives
- Latex

For a full list of ingredients, see the [Fact Sheet for Recipients and Caregivers for the J&J/Janssen COVID-19 vaccine. \[124 KB, 6 pages\]](#)

## Who should get vaccinated

- The J&J/Janssen vaccine is recommended for people aged 18 years and older. Learn more about [how CDC is making COVID-19 vaccine recommendations](#) and [CDC's vaccine rollout recommendations](#)

## Who should not get vaccinated

- If you have had a severe allergic reaction (anaphylaxis) or an immediate allergic reaction—even if it was not severe—to any [ingredient](#) in the J&J/Janssen COVID-19 vaccine (such as polysorbate), you should not get the J&J/Janssen COVID-19 vaccine.
- An allergic reaction is considered severe when a person needs to be treated with epinephrine or EpiPen® or if they must go to the hospital. Experts refer to severe allergic reactions as anaphylaxis. Learn about [common side effects of COVID-19 vaccines](#) and when to call a doctor.
- An immediate allergic reaction means a reaction within 4 hours of getting vaccinated, including symptoms such as hives, swelling, or wheezing (respiratory distress).

If you aren't able to get the J&J/Janssen COVID-19 vaccine, you may still be able to get a different type of COVID-19 vaccine. Learn more [information for people with allergies](#).

# Side effects and safety information

## Most common side effects

In the arm where you got the shot:

- Pain
- Redness
- Swelling

Throughout the rest of your body:

- Tiredness
- Headache
- Muscle pain
- Chills
- Fever
- Nausea

These side effects usually start within a day or two of getting the vaccine. Side effects might affect your ability to do daily activities, but they should go away in a few days. [Get tips on what to expect after getting vaccinated.](#)

## Summary of safety data

- In clinical trials, side effects were common within 7 days of getting vaccinated but were mostly mild to moderate.
- Side effects were more common in people 18–59 years old compared to people 60 years and older.
- CDC will continue to provide updates as we learn more about the safety of the J&J/Janssen vaccine in real-world conditions. [Learn more about vaccine safety monitoring after a vaccine is authorized or approved for use.](#)

[Learn more about safety and reactogenicity data from the clinical trials.](#)

## Information on how well the vaccine works

- The J&J/Janssen vaccine was 66.3% effective in [clinical trials](#) (efficacy) at preventing laboratory-confirmed COVID-19 illness in people who had no evidence of prior infection 2 weeks after receiving the vaccine. People had the most protection 2 weeks after getting vaccinated.
- The vaccine had high efficacy at preventing hospitalization and death in people who did get sick. No one who got COVID-19 at least 4 weeks after receiving the J&J/Janssen vaccine had to be hospitalized.
- Early evidence suggests that the J&J/Janssen vaccine might provide protection against asymptomatic infection, which is when a person is infected by the virus that causes COVID-19 but does not get sick.
- CDC will continue to provide updates as we learn more about how well the J&J/Janssen vaccine works in real-world conditions.

# Demographic information from clinical trials

Clinical trials for the J&J/Janssen vaccine included people from the following racial and ethnic categories:

- 58.7% White
- 45.3% Hispanic or Latino
- 19.4% Black or African American
- 9.5% American Indian or Alaska Native
- 5.6% Multiple races
- 3.3% Asian
- 0.2% Native Hawaiian or other Pacific Islander

Sex breakdown:

- 54.9% Male
- 45.0% Female
- <0.1% Undifferentiated or unknown sex

Age breakdown:

- 66.5% 18–59 years
- 33.5% 60 years and older
  - 19.6% 65 years and older
  - 3.5% 75 years and older

[Learn more about demographic information for people who participated in the trials.](#) 



## For Healthcare Workers

[Johnson & Johnson's Janssen COVID-19 Vaccine:](#) General information, schedule, and administration overview.

## Related Links

[What to Expect After Getting a COVID-19 Vaccine](#)

[Learn more about how federal partners are ensuring the safety of COVID-19 vaccines](#)

[Learn more about the benefits of getting a COVID-19 vaccine](#)

