




COVID-19

ACT NOW!



WEAR A MASK



STAY 6 FEET APART



AVOID CROWDS

Myths and Facts about COVID-19 Vaccines

Updated Feb. 3, 2021

[Print](#)

Now that there are authorized and recommended COVID-19 vaccines in the United States, accurate vaccine information is critical.

How do I know which sources of COVID-19 vaccine information are accurate?

It can be difficult to know which sources of information you can trust. Learn more about [finding credible vaccine information](#).

What You Need to Know about Vaccines

[Frequently Asked Questions about Vaccination](#) >

[8 Things to Know About the Vaccination Program](#) >

Can a COVID-19 vaccine make me sick with COVID-19?

No. None of the authorized and recommended [COVID-19 vaccines](#) or [COVID-19 vaccines currently in development in the United States](#) contain the live virus that causes COVID-19. This means that a COVID-19 vaccine **cannot** make you sick with COVID-19.




There are several different types of vaccines in development. All of them teach our immune systems how to recognize and fight the virus that causes COVID-19. Sometimes this process can cause symptoms, such as fever. These symptoms are normal and are a sign that the body is building protection against the virus that causes COVID-19. Learn more about [how COVID-19 vaccines work](#).

It typically takes a few weeks for the body to build immunity (protection against the virus that causes COVID-19) after vaccination. That means it's possible a person could be infected with the virus that causes COVID-19 just before or just after vaccination and still get sick. This is because the vaccine has not had enough time to provide protection.

After getting a COVID-19 vaccine, will I test positive for COVID-19 on a viral test?

No. Neither the recently authorized and recommended vaccines nor the other COVID-19 vaccines currently in clinical trials in the United States can cause you to test positive on [viral tests](#), which are used to see if you have a **current infection**.



If your body develops an immune response—the goal of vaccination—there is a possibility you may test positive on some [antibody tests](#). Antibody tests indicate you had a **previous infection** and that you may have some level of protection against the virus. Experts are currently looking at how COVID-19 vaccination may affect antibody testing results.

If I have already had COVID-19 and recovered, do I still need to get vaccinated with a COVID-19 vaccine?

Yes, you should be vaccinated regardless of whether you already had COVID-19. That's because experts do not yet know how long you are protected from getting sick again after recovering from COVID-19. Even if you have already recovered from COVID-19, it is possible—although rare—that you could be infected with the virus that causes COVID-19 again.

Learn more about [why getting vaccinated is a safer way to build protection than getting infected](#).



If you were treated for COVID-19 with monoclonal antibodies or convalescent plasma, you should wait 90 days before getting a COVID-19 vaccine. Talk to your doctor if you are unsure what treatments you received or if you have more questions about getting a COVID-19 vaccine.

Experts are still learning more about how long vaccines protect against COVID-19 in real-world conditions. CDC will keep the public informed as new evidence becomes available.

Will a COVID-19 vaccination protect me from getting sick with COVID-19?

Yes. COVID-19 vaccination works by teaching your immune system how to recognize and fight the virus that causes COVID-19, and this protects you from getting sick with COVID-19.

Being protected from getting sick is important because even though many people with COVID-19 have only a mild illness, others may get a [severe illness](#), have [long-term health effects](#), or even die. There is no way to know how COVID-19 will affect you, even if you don't have an [increased risk of developing severe complications](#). Learn more about [how COVID-19 vaccines work](#).




Will a COVID-19 vaccine alter my DNA?

No. COVID-19 mRNA vaccines do not change or interact with your DNA in any way.

Messenger RNA vaccines—also called mRNA vaccines—are the first COVID-19 vaccines authorized for use in the United States. mRNA vaccines teach our cells how to make a protein that triggers an immune response. The mRNA

from a COVID-19 vaccine never enters the nucleus of the cell, which is where our DNA is kept. This means the mRNA cannot affect or interact with our DNA in any way. Instead, COVID-19 mRNA vaccines work with the body's natural defenses to safely develop immunity to disease. Learn more about [how COVID-19 mRNA vaccines work](#).

At the end of the process, our bodies have learned how to protect against future infection. That immune response and making antibodies is what protects us from getting infected if the real virus enters our bodies.



Is it safe for me to get a COVID-19 vaccine if I would like to have a baby one day?

Yes. People who want to get pregnant in the future may receive the COVID-19 vaccine.

Based on current knowledge, experts believe that COVID-19 vaccines are unlikely to pose a risk to a person trying to become pregnant in the short or long term. Scientists study every vaccine carefully for side effects immediately and for years afterward. The COVID-19 vaccines are being studied carefully now and will continue to be studied for many years, similar to other vaccines.



The COVID-19 vaccine, like other vaccines, works by training our bodies to develop antibodies to fight against the virus that causes COVID-19, to prevent future illness. There is currently no evidence that antibodies formed from COVID-19 vaccination cause any problems with pregnancy, including the development of the placenta. In addition, there is no evidence suggesting that fertility problems are a side effect of ANY vaccine. People who are trying to become pregnant now or who plan to try in the future may receive the COVID-19 vaccine when it becomes available to them.

Last Updated Feb. 3, 2021

Content source: [National Center for Immunization and Respiratory Diseases \(NCIRD\), Division of Viral Diseases](#)