



Published in final edited form as:

*J Agromedicine*. 2021 July ; 26(3): 313–322. doi:10.1080/1059924X.2020.1765931.

## Association of category of cattle exposure with tuberculosis knowledge among dairy workers in Bailey County, Texas

Anabel Rodriguez, PhD, MPH<sup>1</sup>, David I. Douphrate, PhD, MPT, MBA, CPE, CSP<sup>2</sup>, David Gimeno Ruiz de Porras, MSc, PhD<sup>3</sup>, Adriana Perez, PhD, MS<sup>4</sup>, Robert Hagevoort, PhD<sup>5</sup>, Matthew Nonnenmann, PhD, MS, CIH<sup>6</sup>, Leroy Cienega, MPH<sup>7</sup>

<sup>1</sup>University of Texas Health Science Center at Houston, School of Public Health in San Antonio, Texas USA, Department of Epidemiology, Human Genetics & Environmental Sciences, 7411 John Smith Drive, Suite 1100, San Antonio, Texas 78229

<sup>2</sup>University of Texas Health Science Center at Houston, School of Public Health in San Antonio, Texas USA, Department of Epidemiology, Human Genetics & Environmental Sciences, 7411 John Smith Drive, Suite 1100, San Antonio, Texas 78229

<sup>3</sup>University of Texas Health Science Center at Houston, School of Public Health in San Antonio, Texas USA, Department of Epidemiology, Human Genetics & Environmental Sciences, 7411 John Smith Drive, Suite 1100, San Antonio, Texas 78229

<sup>4</sup>University of Texas Health Science Center at Houston, School of Public Health in Austin, Department of Biostatistics and Data Science, 1616 Guadalupe Street, Suite 6.340, Austin, Texas 78701

<sup>5</sup>New Mexico State University, College of Agricultural, Consumer, and Environmental Sciences, Department of Agricultural Science Center at Clovis, 2346 State Road 288, Clovis, New Mexico 88101

<sup>6</sup>The University of Iowa, College of Public Health, Department of Occupational and Environmental Health, 145 N. Riverside Drive, 100 CPHB, Iowa City, Iowa 52242

<sup>7</sup>Los Alamos National Laboratory, Department of Industrial Hygiene and Safety, P.O. Box 1663, Los Alamos, New Mexico 87545

### Abstract

**Objective**—To determine the knowledge of TB among dairy workers in Bailey County, Texas.

**Methods**—A cross-sectional study design was used to collect 225 survey responses concerning knowledge of TB among dairy workers on ten dairy farms in Bailey County, Texas. iPad tablets were used to log responses to 15 demographic questions and 17-item TB knowledge quiz measured by: (1) TB characteristics, (2) TB transmission, (3) TB symptoms, (4) TB diagnosis, (5) TB treatment, and (6) bovine TB. A proxy for exposure, dairy workers were classified into categories of cattle exposure—high, medium, and low.

---

**Corresponding Author:** Correspondence to: Dr. Anabel Rodriguez, University of Texas Health Science Center at Houston, School of Public Health in San Antonio, Texas USA, Department of Epidemiology, Human Genetics & Environmental Sciences, 7411 John Smith Drive, Suite 1100, San Antonio, Texas 78229, Phone: 1-210-276-9030, Anabel.Rodriguez@uth.tmc.edu.

**Results**—Relative to the medium/low group, workers in the high group tended to be younger (32.6 (SD 11.0)), Guatemalan (52.6%), K'iche' speaking (37.8%), males (89.1%) with less years living in the U.S. (10.8 (SD 12.5)) and lower levels of formal education completed (59.6% with no formal/elementary level). Overall, the average score was 7.1 (SD 4.9) out of 17 (41.8% out of 100.0%). Relative to one another, the medium/low group (8.0 (SD 4.6) out of 17) scored better than high group (6.7 (SD 5.1) out of 17). No significant associations were found between category of exposure and TB knowledge score.

**Conclusion**—Deficiencies in TB knowledge were identified at all categories of exposure. TB training on dairy farms should include all measured tested in this study and should be administered to all workers regardless of category of exposure.

### Keywords

dairy; workers; knowledge; tuberculosis; Texas

## BACKGROUND

A health hazard on a dairy farm is the potential exposure to *Mycobacterium tuberculosis* (TB).<sup>1–6</sup> In general, TB affects one out of four individuals globally.<sup>7,8</sup> Most recently, the U.S. reported a rate of 2.9 TB cases per 100,000 persons—a record low.<sup>8</sup> However, 67.9% of confirmed TB cases in 2018 originated from foreign-born individuals residing in the U.S.<sup>8</sup> Besides the human version, there is also a *bovine* (cattle) version of the disease called *Mycobacterium bovis* (bTB) or bovine tuberculosis. Bovine TB is predominantly found among cattle and other grazing animals. However, bTB is also transmissible to humans due in part to its zoonotic characteristic.<sup>2,3,9</sup> Globally, an estimated 147,000 bTB cases were confirmed and 12,500 deaths recorded in 2016.<sup>10</sup> The overall impact of TB/bTB among U.S. dairy workers remains unknown.<sup>3</sup>

Large animal veterinarians undergo extensive bTB training during professional schooling.<sup>11</sup> Through this training, veterinarians learn the characteristics, transmission, symptoms, diagnostic tests, treatment, and prevention of bTB among cattle. They are also trained on the inherent health hazards while working with bTB suspected cattle and the potential health consequences.<sup>12</sup> However, milkers and all other job positions on a dairy farm do not undergo this type of extensive professional training education.<sup>13,14</sup> There is a small body of literature addressing bTB and TB knowledge among dairy workers. In addition, there is a limited understanding on the level of knowledge dairy workers have concerning the characteristics, transmission, symptoms, diagnostic tests, treatment, and prevention of TB as well as the potential exposure of bTB on a dairy farm. Currently, there are no standard TB knowledge questionnaires for dairy workers or other vulnerable high risk occupations.<sup>15</sup> Knowing levels of TB knowledge and awareness among dairy workers, can help guide the creation, delivery, and evaluation of a culturally, linguistically, and literacy conscious health and safety training focused on TB and bTB prevention.

Currently, the Department of Labor's Occupational Safety and Health Administration (DOL-OSHA) requires training on applicable hazards, but does not specifically address a form of safety training on TB and bTB as potential transmissible diseases on the farm.<sup>3</sup> The few

studies which have investigated the prevalence of bTB among dairy workers have suggested worker PPE and bTB education programs as an addition to the existing governmental eradication programs.<sup>2,4–6</sup> However, before a TB and bTB educational course/program can be created, delivered, and evaluated on dairy farms, a need exists to assess what content needs to be included and deficiencies in knowledge that need to be addressed.<sup>13,16</sup> The primary objective is to determine the awareness and knowledge of TB among dairy workers in Bailey County, Texas. This study was approved by the University of Texas Health Science Center at Houston Committee of the Protection of Human Subjects (CPHS) (HSC-SPH-18-0886).

## METHODS

### Study design.

A cross-sectional study design was used to collect survey responses concerning knowledge of TB among dairy workers in Bailey County, Texas. Bailey County has a total of 10 farms, employs approximately 225 workers, and milks an estimated 22,537 cows.<sup>17</sup> All ten dairy farm producers were called, personally visited, and invited to participate in this study. A total of 225 dairy workers were included in the study. Data collection took place between February and March 2019.

### Eligibility criteria.

A total of 225 consenting dairy workers were invited to participate and surveyed in Bailey County, Texas. Subject eligibility included being a male or female worker 18 years of age employed full-time/part-time/temporary on any of the ten dairy farms visited.

### Consent procedures.

Research staff read and explained the consent form to participants before the survey was completed. Participants were asked to consent and sign an electronic informed consent on an iPad® tablet in order to participate. All participants were given a hard copy of their consent form in English or Spanish. Once consent was collected, research staff proceeded to administer the survey. Surveys were administered in privacy in breakrooms, conference rooms, parlors, maintenance sheds, tractors, and other accessible dairy farm work spaces. Participants were compensated for their time with a \$10 gift card.

### Survey measures.

The survey included 15 demographic questions and a 17-item TB knowledge quiz. Currently, there are no standard TB knowledge questionnaires for dairy workers or other vulnerable high risk occupations.<sup>15</sup> Therefore, the help of a dairy extension specialist and previously published peer-reviewed literature was used to finalize the survey content. All TB knowledge questions were adopted from material publicly released by the CDC,<sup>18</sup> WHO,<sup>19</sup> University of Rochester Medical Center,<sup>20</sup> and a current study on workplace TB interventions by Eggerth et al. (2018). All questions were placed on the survey platform Qualtrics Mobile Survey Software® with both English and Spanish options. Subsequently, surveys were uploaded to iPad® tablet devices for offline use. Trained bilingual (English and Spanish proficient) research staff used these iPad® tablet devices to read questions to

participants and log, in real-time, their responses (Figure 1). Completed surveys were uploaded to our private and encrypted Qualtrics online account once internet services were available.

### Outcome variable.

TB knowledge, was measured via several questions on six different aspects of TB knowledge: (1) TB characteristics, (2) TB transmission, (3) TB symptoms, (4) TB diagnosis, (5) TB treatment, and (6) bovine TB. Questions testing knowledge were asked in both “True-False” and multiple choice format. In addition, administrators had the option of selecting “I don’t know” if the participant expressed not knowing the answer to a question or also had the option of selecting “Did not answer” if the participant did not choose an answer or did not want to answer the question. Each participant had a maximum of 17 points: one point for a correct answer and zero points for an incorrect answer (*Knowledge questions in Appendix A*).

### Exposure measure.

Following previous research by Torres et al. (2013), job position on a dairy farm was used as a proxy for categories of cattle exposure: (1) high exposure among workers with direct contact with cattle in confined spaces (e.g., milkers, pusher, veterinarians, supervisor/manager, hospital workers, and slaughter); (2) medium exposure among workers with direct contact with cattle in non-confined spaces (e.g., feeders, tractor operators, breeders, calf caretaker, maternity, hoof trimmer, maintenance technicians); and, (3) low exposure among workers with no direct contact with cattle in any type of space (e.g., owners, office staff).

### Data analyses.

A preliminary dataset check was completed in order to assess the percentage of missing data. Subsequently, a complete case analyses (CCA) was preformed because < 10% of the data was missing. Basic descriptive statistics (e.g., frequencies, proportions, means, and standard deviations) of all sociodemographic characteristics by category of exposure were estimated and reported in Table 1. Both chi-square and the nonparametric Kruskal-Wallis tests were conducted to explore potential confounders between category of exposure and sociodemographic variables. Corresponding p-values are also shown in Table 1. A type I error level of 0.05 was used to declare significance.

Table 2 shows individual TB knowledge questions by category of exposure (high, medium, low). The medium and low groups were collapsed due to a limited sample size. After analyzing TB knowledge scores separated on the basis of “Correct,” “Incorrect,” and “Don’t know,” by high, medium, and low category of exposure, we noticed the sample size for the low group was much lower (n=5) compared to high (n=156) and medium (n=64). The medium and low group frequencies for “Correct,” “Incorrect,” and “Don’t know” were statistically similar; therefore, we decided to collapse these groups in to one labeled medium/low. In this same preliminary analysis, we also collapsed the “Incorrect” and “Don’t know” categories of TB knowledge due to sample size and conceptual methods. A separate analysis found that several “Incorrect” cells had counts < 5. Conceptually, an incorrect answer indicates a gap in knowledge or a state of not knowing the answer to a question.

Correspondingly, previous studies assessing content knowledge, using similar methodological techniques, collapsed “Incorrect” and “Don’t know” categories.<sup>4,21,22</sup>

Frequencies are reported for correct and incorrect answer choices for each of the 17-questions by all, high, and medium/low groups (Table 2). In addition, multivariate models were fitted for individual knowledge questions utilizing Hosmer and Lemeshow’s model building methods. Crude prevalence odds ratio (POR), adjusted prevalence odds ratios (aPOR), and corresponding 95% confidence intervals (95%CI) were reported (Table 1). All statistical analyses were performed using Stata/SE v.14.0.<sup>23</sup>

## RESULTS

The mean age of workers was 34.4 (SD 12.0) with a range of 17-65 years of age and 89.3% of surveyed dairy workers were male. Almost all dairy workers (96.9%) were full-time employees with 7.4 (SD 8.2) years of dairy farm work experience in the U.S. and 4.8 (SD 8.8) years of experience working with cattle in their country of origin. The majority of dairy workers were Hispanic (88.0%) with 43.1% of participants reporting Mexico as their country of birth, 45.3% other Latin American countries, and 11.6% United States. On average, workers had 13.0 (SD 14.0) years residing in the U.S. with a range of 3 months to 64 years. Almost 60.0% of workers reported Spanish as their primary language, 9.8% English, and 30.7% spoke another language. In contrast, 36.9% reported Spanish as their secondary language, 20.9% as English, 6.7% spoke another language, and 35.6% claimed they had no proficient secondary language. The majority of workers reported no formal education/elementary (51.6%) as the highest level of education achieved, followed by high school/college/graduate (30.2%) levels and middle school (18.2%).

The sociodemographic characteristics of surveyed dairy workers by category of exposure: high and medium/low groups are found in Table 1. Workers in the high category of exposure group tended to be younger, Guatemalan, K’iche’ speaking males with less years living in the U.S. and lower levels of formal education completed. Relative to the medium/low group, high group had less dairy farm work experience in the U.S. and less work experience with cattle in their country of origin, but worked similar days per week and hours per day.

In this study, general awareness refers to consciousness that a condition (e.g., TB) exists and knowledge refers to understanding facts/information about a subject (e.g., TB).<sup>24</sup> Overall, 37.3% of surveyed workers had general awareness of TB—the high group was less aware of TB (34.0%) compared to the medium/low group (44.9%). The individual TB knowledge questions by category of TB exposure including high and medium/low groups are found in Table 2. Overall, the average score was 7.1 (SD 4.9) out of 17 (41.8% out of 100.0%). Relative to one another, the medium/low group (8.0 (SD 4.6) out of 17) scored better than high group (6.7 (SD 5.1) out of 17). Besides correct answers, most notable were the frequencies for incorrect answer choices. In general, dairy workers reported higher frequencies of incorrect answers than selecting the correct answers for 12 out of the 17 questions administered. In general, higher frequencies of incorrect answers than correct were found for all aspects of TB knowledge: (1) TB characteristics, (2) TB transmission, (3) TB symptoms, (4) TB diagnosis, (5) TB treatment, and (6) bovine TB.

Crude and adjusted POR and corresponding 95% CIs for the medium/low group with the high as the reference group can be found in Table 2. For crude POR, statistical TB knowledge score differences between the medium/low group relative to high occurred for three items (questions 2, 13, and 14) pertaining to TB characteristics and TB treatment. After adjustments, statistical TB knowledge score differences between the medium/low group relative to high occurred for two different items (questions 8 and 11).

## DISCUSSION

The authors observed TB knowledge deficiencies at all quizzed measures: (1) TB characteristics, (2) TB transmission, (3) TB symptoms, (4) TB diagnosis, (5) TB treatment, and (6) bovine TB. In addition, these deficiencies in TB knowledge were also found in both the high and medium/low exposure groups. Overall, the average score was 7.1 (SD 4.9) out of 17 (41.8% out of 100.0%). Relative to one another, the medium/low group (8.0 (SD 4.6) out of 17) scored better than high group (6.7 (SD 5.1) out of 17). This study also found that 37.3% of surveyed workers had general awareness of TB—the high group was less aware of TB (34.0%) compared to the medium/low group (44.9%).

Similarly, bTB knowledge assessments were conducted among 510 Nigerian dairy workers. Results indicated that 58.6% of herdsmen and 46.9% of abattoir workers were knowledgeable of bTB prevention.<sup>25</sup> Another study performed in Cameroon found that 73.9% out of 164 dairy farmers were aware of bTB. Despite this high bTB awareness, 55.9% were not able to correctly identify clinical signs and symptoms among themselves, coworkers, or cattle on farm.<sup>4</sup> A similar study from Malawi found that 74.3% out of 140 dairy farm workers were aware that bTB was a zoonotic disease; yet, only 15.7% were able to identify preventative measures (e.g., medical check-up) and only 7.9% identified contact with infect animals as a mode of transmission.<sup>26</sup> However, bTB is endemic in Nigeria, Cameroon, and Malawi; whereas, quality control standards have helped Texas manage sporadic bTB outbreaks among cattle.<sup>27</sup> This may be the reason for the difference in awareness levels between dairy workers in these countries and dairy workers in Texas.

Torres-Gonzalez et al. (2013) created three categories of cattle exposure groups based on activity, duration, and conditions of exposure to cattle—high, medium, low. These categories were used in this study to help categorize workers into proxy exposure groups.<sup>6</sup> Results indicated that there were no significant associations between category of exposure and TB knowledge score by question. An alternative to categories of cattle exposure could have been level of education. The no formal education/elementary group scored a 27.5% (out of 100%) compared to the middle school group at 48.4% and the high school/college/graduate with the highest percentage at 61.8%. The study previously described assessing knowledge of dairy workers in Nigeria found that dairy workers with post-primary education were 2.70 (95% CI: 1.68-4.33) more knowledgeable of bTB prevention compared to individuals with no formal education.<sup>25</sup>

### Study limitations

Study limitations included potential recall bias of information collect on the survey. A total of 15 demographic questions and 17 TB knowledge quiz questions were administered.



Asking workers to recall the exact number of years in the U.S., years of experience working on dairy farms, years of experience working with cattle in their country of origin, hours a day and days per week (which can vary in agriculture), among recalling TB knowledge information (if learned in past education) could have led to an underestimation or overestimation of these variables and/or the overall individual TB knowledge score. However, demographic characteristics obtained in this study resulted similar to previous studies indicating dairy workers are predominantly an immigrant,<sup>28</sup> Hispanic male,<sup>29</sup> of approximately 30 years of age<sup>16</sup> with limited English proficiency and formal education.<sup>14</sup> Another source of error could have come from respondent bias. There could have been a difference in reluctance to answer between individuals who had a personal experience or knew someone with TB or had an encounter/familiarity with bovine TB on the farm. In addition, participants could have felt the urgency to answer a question even if the attempt was wrong instead of electing to select “I don’t know.”

According to the USDA National Agricultural Statistics Service (NASS), as of July 1, 2018, Bailey County had a total of 10 licensed farms and milked an estimated 22,537 cows.<sup>17</sup> All ten dairy farm producers listed were called, personally visited, and invited to participate in this study. A total of 225 dairy workers were included in the study. When dairies were visited, producers were asked to provide the number of workers currently employed in order to best prepare for the day of survey administration and gift card compensations. As of March 23, 2019, a total of 293 dairy workers were totaled on ten dairy farms in Bailey County, Texas. This means that 77.0% (225/293) of available workers participated in this study. This information created a more accurate census of dairy workers in Bailey County, Texas. Unfortunately, this census will never be accurate because of high worker turnover rates.<sup>30,31</sup> This participation rate came about eight workers choosing not to participate. Despite the possibility of non-response bias and these eight workers being different compared to those who chose to participate in this study, the number is small enough to not affect overall results obtained. In addition, the remaining 60 workers not included were out on vacation, resting the days we visited the farms, or their work day was too busy for non-work related interruptions. Last, the methods of this study ensured interviewer bias remained low. A detailed script was created to guide research staff through survey administration and ensured quality assurance. A total of two researchers administered all 225 surveys. Both researchers were fluent and literate in English and Spanish. Several team meetings were held before and after data collection trips to train and guarantee consistency of survey administrations.

### Future plans and conclusions

This study found TB knowledge deficiencies at all assessment measures among all categories of TB exposure groups. The results found in this study have allowed us to conclude that a TB educational training could be beneficial for dairy workers at all job positions in Bailey County, Texas. Due to gaps identified in knowledge, the training should include content pertaining to: (1) TB characteristics, (2) TB transmission, (3) TB symptoms, (4) TB diagnosis, (5) TB treatment, and (6) bovine TB. Effective occupational health and safety trainings is a method that can be used to reduce fatal and nonfatal incidents on dairy farms.<sup>32</sup> Health and safety training can be delivered as class lecture, computer training, and

hands-on demonstration.<sup>16,33</sup> Mobile learning (m-learning) uses mobile devices for learning experiences.<sup>34</sup> M-learning has been used in occupational settings to provide learning experiences to individual workers or a group of workers.<sup>13,34</sup> Most recently, safety awareness training was delivered to 1,436 dairy workers in Texas, New Mexico, Colorado, Kansas, and New York using iPad® tablets. This safety training proved effective with a score change from 74.2% in the pre-test (baseline) to a 92.5% average in the post-test.<sup>13</sup> Similar methods can be used to create, deliver, and evaluate a TB educational course in Bailey County, Texas. Pre- and post-tests would be used to assess the change in knowledge gained from training. The majority of workers surveyed identified Spanish as their primary language (60.0%) and 51.6% stated they had no formal education/elementary level education; therefore, this training must be culturally, linguistically, and literacy conscious.<sup>31</sup> Despite the medium/low group scoring slightly higher than the high group, both groups showed low levels of TB knowledge and awareness. This training should be made available and required for all new employees and currently employed workers regardless of their years of experience on dairy farms.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

### Funding

Funding for this research was supported by Grant No. T42OH008421 09 from the National Institute for Occupational Safety and Health (NIOSH)/Centers for Disease Control and Prevention (CDC) to the Southwest Center for Occupational and Environmental Health (SWCOEH), a NIOSH Education and Research Center.

## REFERENCES

1. Adesokan H, Jenkins A, van Soolingen D, Cadmus S. Mycobacterium bovis infection in livestock workers in Ibadan, Nigeria: evidence of occupational exposure. *International Journal of Tuberculosis Lung Disease*. 2012;16(10):1388–1392.
2. Bekele M, Mamo G, Mulat S, Ameni G, Beyene G, Tekeba E. Epidemiology of Bovine Tuberculosis and Its Public Health Significance In Debre-Zeit Intensive Dairy Farms, Ethiopia. *Journal of Biomedicine and Nursing*. 2016;2(2):8–18.
3. Hlavsa M, Moonan P, Cowan LN, TR Kammerer, J S, Morlock GC, J T LoBue PA. Human tuberculosis due to Mycobacterium bovis in the United States, 1995–2005. *Clinical Infectious Diseases*. 2008;47(2):168–175. [PubMed: 18532886]
4. RF K, Hamman S, Morgan K, et al. Knowledge of Bovine Tuberculosis, Cattle Husbandry and Dairy Practices among Pastoralists and Small-Scale Dairy Farms in Cameroon. *PLoS ONE*. 2016;11(1):1–20.
5. Thakur A, Sharma M, Katoch V, Dhar P, Katoch R. A study on the prevalence of Bovine Tuberculosis in farmed dairy cattle in Himachal Pradesh. *Veterinary World*. 2010;3(9):409–414.
6. Torres-Gonzalez P, Soberanis-Ramos O, Martinez-Gamboa A, et al. Prevalence of Latent and Active Tuberculosis among Dairy Farm Workers Exposed to Cattle Infected by Mycobacterium bovis. *PLOS Neglected Tropical Diseases*. 2013;7(7):1–8.
7. (BLS) BoLS. 2016 Census of Fatal Occupational Injuries. 2017; <https://www.bls.gov/iif/oshwc/cfoi/cfch0015.pdf>.
8. CDC. Chapter 2: Transmission and pathogenesis of tuberculosis. 2016; <https://www.cdc.gov/tb/education/corecurr/pdf/chapter2.pdf>



9. Palmer M, Thacker T, Waters W, Gortazar C, Corner L. Mycobacterium bovis: A Model Pathogen at the Interface of Livestock, Wildlife, and Humans. *Veterinary Medicine International*. 2012;2012:1–17.
10. (WHO) WHO. The challenges of preventing bovine tuberculosis 2018.
11. Medicine ISUCoV. Tuberculosis. 2018; <https://vetmed.iastate.edu/vdpam/FSVD/swine/index-diseases/tuberculosis>.
12. Agriculture USDo. Tuberculosis. 2017; <https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/nvap/NVAP-Reference-Guide/Control-and-Eradication/Tuberculosis>.
13. Rodriguez A, Hagevoort G, Leal D, Pompeii L, Douphrate D. Using mobile technology to increase safety awareness among dairy workers in the United States. Under review. Under review.
14. Jenkins P, Stack S, May J, Earle-Richardson G. Growth of the Spanish-speaking workforce in the northeast dairy industry. *Journal of Agromedicine*. 2009;14(1):58–65. [PubMed: 19214856]
15. Eggerth D, Keller B, Flynn M. Exploring workplace TB interventions with foreign-born Latino workers. *American Journal of Industrial Medicine*. 2018:1–9.
16. Roman-Muniz IVM DC, Garry F, Reynolds S, Wailes W, Keefe T. Training methods and association with worker injury on Colorado dairies: a survey. *Journal of Agromedicine*. 2006;11(2):19–26. [PubMed: 17135139]
17. (NASS) NASS. Overview of the United States Dairy Industry: United States Department of Agriculture (USDA); 2010:1–13.
18. CDC. Test your TB knowledge 2009; <https://www2a.cdc.gov/tbquiz.asp>.
19. (WHO) WHO. Accelerating Advocacy for TB/HIV: Tuberculosis Quiz 2010; [http://www.who.int/tb/challenges/hiv/01\\_tb\\_quiz\\_eng.pdf](http://www.who.int/tb/challenges/hiv/01_tb_quiz_eng.pdf).
20. Fraser M, Lentnek A. What do you know about tuberculosis? In: Center UoRM, ed2018.
21. Eaden J, Abrams K, Mayberry J. The Crohn's and Colitis Knowledge Score: A Test for Measuring Patient Knowledge in Inflammatory Bowel Disease. *The Journal of Gastroenterology*. 1999;94(12):3561–3566.
22. Al-Maniri A, Al-Rawas O, Al-Ajmi F, Costa A, Eriksson B, Diwan V. Tuberculosis suspicion and knowledge among private and public general practitioners: Questionnaire Based Study in Oman. *BMC Public Health*. 2008;8(177):1–7. [PubMed: 18173844]
23. Stata Statistical Software: Release 14 [computer program]. College Station, TX 2015.
24. Trevethan R. Deconstructing and Assessing Knowledge and Awareness in Public Health Research. *Frontiers in Public Health*. 2017;5(194):1–6. [PubMed: 28229066]
25. Adesokan H, Akinseye V, Sulaimon M. Knowledge and practices about zoonotic tuberculosis prevention and associated determinants amongst livestock workers in Nigeria; 2015. *PLoS ONE*. 2018;13(6):e0198810. [PubMed: 29889870]
26. Tebug S, Njunga GC, M GG, Awah-Ndukum J, Widemann S. Risk, knowledge and preventive measures of smallholder dairy farmers in northern Malawi with regard to zoonotic brucellosis and bovine tuberculosis. *Journal of Veterinary Research*. 2014;81(1):6.
27. Winthrop K, Scott J, Brown D, et al. Investigating of human contacts: A Mycobacterium bovis outbreak among cattle at a California dairy. *International Journal of Tuberculosis Lung Disease*. 2004;9(7):809–8013.
28. Passel J, Cohn D, Rohal M. Unauthorized immigrant totals rise in 7 states, fall in 14: Decline in those from Mexico fuels most state decreases. Washington, DC: Pew Research Center, Project HT. 2014.
29. Adock F, Anderson D, Rosson P. The economic impacts of immigration labor on US dairy farms 2015.
30. Grey M, Immigrant migrants, and work turnover at the Hog Pride Packing Plant. *Human Organization*. 1999;58(1):16–27.
31. Arcury T, Estrada J, Quandt S. Overcoming language and literacy barriers in safety and health training of agricultural workers *Journal of Agromedicine*. 2010;15(3):236–248. [PubMed: 20665309]
32. OSHA. Recommended Practice for Safety and Health Programs. Washington, D.C.2016. 3885.

33. Burke M, Sarpy S, Smith-Crowe K, Chan-Serafin S, Salvador R, Islam G. Relative effectiveness of worker safety and health training methods. *American Journal of Public Health*. 2006;96(2):315–324. [PubMed: 16380566]
34. Park Y A pedagogical framework for mobile learning: categorizing educational applicaitons of mobile technologies into four types. *Int Rev Res Open Distributed Learn*. 2011;12(2):78–102.



**Figure 1.**  
Bilingual research staff member administering survey to dairy worker outside maintenance shed in Bailey County, Texas.

**Table 1.**

Demographic characteristics of surveyed dairy workers by category of TB exposure.

Characteristics	Category of TB exposure			<i>p</i> <sup>*</sup>
	All (n=225)	High (n=156)	Medium/Low (n=69)	
<b>Age</b>	34.4 (12.0)	32.6 (11.0)	38.3 (13.1)	0.0022
<b>Male</b>	201 (89.3)	139 (89.1)	62 (89.9)	0.8660
<b>Nationality</b>				
United States	26 (11.6)	11 (7.1)	15 (21.7)	<0.0001
Mexico	97 (43.1)	57 (36.5)	40 (58.0)	
Other Latin American Countries	102 (45.3)	88 (56.4)	14 (20.3)	
<b>Years in the US</b>	13.0 (14.0)	10.8 (12.5)	18.0 (16.0)	0.0002
<b>Primary Language</b>				<0.0001
English	22 (9.8)	11 (7.1)	11 (15.9)	
Spanish	134 (59.6)	83 (53.2)	51 (73.9)	
Other	69 (30.7)	62 (39.7)	7 (10.1)	
<b>Secondary Language</b>				0.0010
English	47 (20.9)	23 (14.7)	24 (34.8)	
Spanish	83 (36.9)	67 (43.0)	16 (23.2)	
Other	15 (6.7)	13 (8.3)	2 (2.9)	
None	80 (35.6)	53 (34.0)	27 (39.1)	
<b>Education</b>				0.0010
No Formal/Elementary school	116 (51.6)	93 (59.6)	23 (33.3)	
Middle school	41 (18.2)	24 (15.4)	17 (24.6)	
High school/College/Graduate	68 (30.2)	39 (25.0)	29 (42.0)	
<b>Years of experience on dairy farms</b>	7.4 (8.2)	7.0 (8.6)	8.3 (7.5)	0.0410
<b>Years working with cattle in origin country</b>	4.8 (8.8)	4.3 (8.6)	6.0 (9.3)	0.1482
<b>Full-time employment</b>	218 (96.9)	151 (96.8)	67 (97.1)	0.5800
Hours per day	10.6 (6.0)	10.7 (6.5)	10.4 (4.8)	0.0715
Days per week	6.0 (0.6)	6.0 (0.6)	6.0 (0.6)	0.8773

\* p-value from  $\chi^2$ ; p-value from Kruskal-Wallis

Table 2.

TB knowledge scores by categories of TB exposure (n=225)

TB knowledge evaluation content	Category of TB exposure						POR (95%CI) <sup>c</sup>	
	All (n=225)	Medium/Low (n=69)				Medium/Low (High as reference)		
		High (n=156)	Correct <sup>a</sup>	Incorrect <sup>b</sup>	n(%)			
TB CHARACTERISTICS								
1. TB is caused by germs called bacteria. <sup>d</sup>	123 (54.7)	102 (45.3)	79 (50.6)	77 (49.4)	44 (63.8)	25 (36.2)	1.7 (1.0-3.1)	0.9 (0.4-1.8)
2. The flu vaccine protects me from TB infections. <sup>d</sup>	77 (34.2)	148 (65.8)	47 (30.1)	109 (69.9)	30 (43.5)	39 (56.5)	1.8 (1.0-3.2)	0.8 (0.4-1.6)
3. TB affects the lungs and other organs. <sup>d</sup>	117 (52.0)	108 (48.0)	75 (48.1)	81 (51.9)	42 (60.9)	27 (39.1)	1.7 (0.9-3.0)	0.9 (0.4-1.7)
TB TRANSMISSION								
4. How do you get TB? <sup>d</sup>	64 (28.4)	161 (71.6)	44 (28.21)	112 (71.8)	20 (29.0)	49 (71.0)	1.0 (0.6-1.9)	0.8 (0.4-1.6)
5. TB can be transmitted from person-to-person through touching or sharing plates and cups. <sup>e</sup>	22 (9.8)	203 (90.2)	14 (9.0)	142 (91.0)	8 (11.6)	61 (88.4)	1.3 (0.5-3.3)	0.9 (0.3-2.4)
6. Who is at risk of developing TB in this country? <sup>d</sup>	82 (36.4)	143 (63.6)	54 (34.6)	102 (65.4)	28 (40.6)	41 (59.4)	1.3 (0.7-2.3)	0.5 (0.3-1.1)
TB SYMPTOMS								
7. What are the main symptoms of TB disease? <sup>d</sup>	74 (32.9)	151 (67.1)	51 (32.7)	105 (67.3)	23 (33.3)	46 (66.7)	1.0 (0.6-1.9)	0.5 (0.3-1.1)
8. You could have TB and not have symptoms. This is called latent tuberculosis. <sup>f</sup>	76 (33.8)	149 (66.2)	57 (36.5)	99 (63.5)	19 (27.5)	50 (72.5)	0.7 (0.4-0.2)	0.3 (0.2-0.7)
TB DIAGNOSIS								
9. TB can be tested by your local clinic. <sup>g</sup>	105 (46.7)	120 (53.3)	69 (44.2)	87 (55.8)	36 (52.2)	33 (47.8)	1.4 (0.8-2.4)	0.7 (0.4-1.4)
10. How is TB diagnosed? <sup>h</sup>	47 (20.9)	178 (79.1)	29 (18.6)	127 (81.4)	18 (26.1)	51 (73.9)	1.5 (0.8-3.0)	0.9 (0.4-1.8)
TB TREATMENT								

TB knowledge evaluation content	Category of TB exposure						POR (95%CI) <sup>c</sup>	
	All (n=225)		High (n=156)		Medium/Low (n=69)			
	Correct <sup>a</sup>	Incorrect <sup>b</sup>	Correct <sup>a</sup>	Incorrect <sup>b</sup>	Correct <sup>a</sup>	Incorrect <sup>b</sup>	Crude	Adjusted
11. How is TB treated? <sup>g</sup>	96 (42.7)	129 (57.3)	67 (43.0)	89 (57.1)	29 (42.0)	40 (58.0)	1.0 (0.5-1.7)	0.4 (0.2-0.7)
12. TB can be cured. <sup>f</sup>	106 (47.1)	119 (52.9)	69 (44.2)	87 (55.8)	37 (53.6)	32 (46.4)	1.5 (0.8-2.6)	1.0 (0.5-1.8)
13. Untreated TB can be fatal. <sup>d</sup>	142 (63.1)	83 (36.9)	86 (55.1)	70 (44.9)	56 (81.2)	13 (18.8)	3.5 (1.8-6.9)	1.9 (0.8-4.2)
14. TB can be cured drinking tea and making natural home remedies. <sup>d</sup>	103 (45.8)	122 (54.2)	59 (37.8)	97 (62.2)	44 (63.8)	25 (26.2)	2.9 (1.6-5.2)	1.2 (0.5-2.5)
BOVINE TB								
15. Cattle can also experience a TB infection called bovine tuberculosis. <sup>i</sup>	145 (64.4)	80 (35.6)	96 (61.5)	60 (38.5)	49 (71.0)	20 (29.0)	1.5 (0.8-2.8)	0.7 (0.3-1.5)
16. Transmission of bovine TB can happen between cattle and humans. <sup>g</sup>	139 (61.8)	86 (38.2)	95 (60.9)	61 (39.1)	44 (63.8)	25 (36.2)	1.1 (0.6-2.0)	0.6 (0.3-1.1)
17. How can bovine TB be transmitted? <sup>d</sup>	76 (33.8)	149 (66.2)	49 (31.4)	107 (68.6)	27 (39.1)	42 (60.9)	1.4 (0.8-2.5)	0.9 (0.5-1.8)
Mean (SD) TB Knowledge Score (max. 17)	7.1 (4.9)		6.7 (5.1)		8.0 (4.6)			

<sup>a</sup>Frequency and percentage of correct answers chosen;<sup>b</sup>Incorrect is the sum of wrong and 'don't know' answers;<sup>c</sup>Prevalence odds ratio (POR) and corresponding 95% confidence interval;<sup>d</sup>Adjusted for age primary language, education, and years of dairy work experience;<sup>e</sup>Adjusted for education;<sup>f</sup>Adjusted for primary language;<sup>g</sup>Adjusted for primary language and education;<sup>h</sup>Adjusted for age and education;<sup>i</sup>Adjusted for primary language, education, and years of dairy work experience.