

Archived Editions (COVID-19 Genomics and Precision Public Health Weekly Update)

Published on 03/04/2021

COVID-19 Genomics and Precision Public Health Weekly Update Content

- Pathogen and Human Genomics Studies
- Non-Genomics Precision Health Studies
- News, Reviews and Commentaries

Pathogen and Human Genomics Studies

- Report Outlines Strategy for Surveillance of Emerging Coronavirus Variants (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=813)
Stephenson J. JAMA Network, Feb 23, 2021.

- COVID-19 disruptions to endoscopic surveillance in Lynch syndrome. (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=814)
McKenna Danielle B et al. Cancer prevention research (Philadelphia, Pa.) 2021 2

This study demonstrates that the COVID-19 pandemic shutdown led to delayed endoscopic surveillance in Lynch syndrome, with potentially impactful delays among young patients. This data also emphasizes the importance of timely surveillance in Lynch syndrome during this current, as well as potential future, global pandemics.

- Microbial signatures in the lower airways of mechanically ventilated COVID19 patients associated with poor clinical outcome (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=817)
I Sulaiman et al, MEDRXIV, February 26, 2021

We found that isolation of a hospital-acquired respiratory pathogen was not associated with fatal outcome. However, poor clinical outcome was associated with enrichment of the lower airway microbiota with an oral commensal (*Mycoplasma salivarium*), while high SARS-CoV-2 viral burden, poor anti-SARS-CoV-2 antibody response, together with a unique host transcriptome profile of the lower airways were most predictive of mortality.

- A Novel SARS-CoV-2 Variant of Concern, B.1.526, Identified in New York (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=818)
MK Annavajhala et al, MEDRXIV, February 25, 2021

To survey for these novel variants in our patient population in New York City, PCR assays were designed to identify viruses with two signature mutations, E484K and N501Y. We observed a steady increase in the detection rate from late December to mid-February, with an alarming rise to 12.3% in the past two weeks.

- A Neanderthal OAS1 isoform protects individuals of European ancestry against COVID-19 susceptibility and severity (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=819)
S Zhou et al, Nature Medicine, February 25, 2021

To identify circulating proteins influencing Coronavirus Disease 2019 (COVID-19) susceptibility and severity, we undertook a two-sample Mendelian randomization (MR) study, rapidly scanning hundreds of circulating proteins while reducing bias due to reverse causation and confounding. In up to 14,134 cases and 1.2 million controls, we found that an s.d. increase in OAS1 levels was associated with reduced COVID-19 death or ventilation, hospitalization and susceptibility.

- Common genetic variation in humans impacts in vitro susceptibility to SARS-CoV-2 infection. (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=831)
Dobrindt Kristina et al. Stem cell reports 2021 2

We hypothesize that, in addition to viral load and host antibody repertoire, host genetic variants influence vulnerability to infection. Here we apply human induced pluripotent stem cell (hiPSC)-based models and CRISPR engineering to explore the host genetics of SARS-CoV-2. We demonstrate that a single-nucleotide polymorphism (rs4702), common in the population and located in the 3' UTR of the protease *FURIN*, influences alveolar and neuron infection by SARS-CoV-2 in vitro.

- Escape of SARS-CoV-2 501Y.V2 variants from neutralization by convalescent plasma (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=832)

S Cele et al, MEDRXIV, February 27, 2021

Here, using a live virus neutralization assay (LVNA), we compared neutralization of a first wave virus (B.1.1.117) versus the 501Y.V2 variant using plasma collected from adults hospitalized with COVID-19 from two South African infection waves, with the second wave dominated by 501Y.V2 infections.

- Increased hazard of mortality in cases compatible with SARS-CoV-2 variant of concern 202012/1 - a matched cohort study (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=833)

R Challen et al, MEDRXIV, February 27, 2021

In this matched cohort study of 54,773 pairs of participants matched by age, gender, ethnicity, index of multiple deprivation, lower tier local authority region, and sample date of positive specimen, and differing only by detectability of the spike protein gene. Death within 28 days after first positive test was higher among patients with VOC-202012/01. (hazard ratio 1.7, 95% CI 1.3-2.2).

- A statewide analysis of SARS-CoV-2 transmission in New York (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=836)

EL Nesselquist et al, MEDRXIV, February 28, 2021

The Wadsworth Center sought to characterize the transmission dynamics of SARS-CoV-2 across the entire state of New York from March to September with the addition of over 600 genomes from under-sampled and previously unsampled New York counties and to more fully understand the breadth of the initial outbreak in Westchester County. Additional sequencing confirmed the dominance of B.1 and descendant lineages (collectively referred to as B.1.X) in New York State.

- Medical vulnerability of individuals with Down syndrome to severe COVID-19-data from the Trisomy 21 Research Society and the UK ISARIC4C survey (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=838)

A Huls et al, Ebiomedicine, February 2021

Health conditions, immune dysfunction, and premature aging associated with trisomy 21 (Down syndrome, DS) may impact the clinical course of COVID-19. The T21RS COVID-19 Initiative launched an international survey for clinicians or caregivers on patients with COVID-19 and DS. Data collected between April and October 2020 (N=1046) were analyzed and compared with the UK ISARIC4C survey of hospitalized COVID-19 patients with and without DS.

- Genetics of symptom remission in outpatients with COVID-19 (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=846)

MP Dube et al, MEDRXIV, March 1, 2021

We conducted a genome-wide association study of time to remission of COVID-19 symptoms in 1723 outpatients with at least one risk factor for disease severity from the COLCORONA clinical trial. We found a significant association at 5p13.3 (rs1173773; $P = 4.94 \times 10^{-8}$) near the natriuretic peptide receptor 3 gene (NPR3). By day 15 of the study, 44%, 54% and 59% of participants with 0, 1, or 2 copies of the effect allele respectively, had symptom remission.

- Transmission of SARS-CoV-2 in domestic cats imposes a narrow bottleneck. (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=849)

Braun Katarina M et al. PLoS pathogens 2021 2 (2) e1009373

"In RNA virus evolution, stochastic processes like narrow transmission bottlenecks and genetic drift typically act to constrain the overall pace of adaptive evolution. Our data suggest that here, positive selection in index cats followed by a narrow transmission bottleneck may have instead accelerated the fixation of S H655Y, a potentially beneficial SARS-CoV-2 variant. Overall, our study suggests species- and context-specific adaptations are likely to continue to emerge. This underscores the importance of continued genomic surveillance for new SARS-CoV-2 variants as well as heightened scrutiny for signatures of SARS-CoV-2 positive selection in humans and mammalian model systems."

- Sequence Analysis of 20,453 SARS-CoV-2 Genomes from the Houston Metropolitan Area Identifies the Emergence and

Widespread Distribution of Multiple Isolates of All Major Variants of Concern (/PHGKB/phgHome.action?

action=forward&dbsource=covUpdate&id=850)

Long SW, et al. medRxiv, Mar 2, 2021.

"Houston is the first city in the United States to have all variants documented by genome sequencing. As vaccine deployment accelerates worldwide, increased genomic surveillance of SARS-CoV-2 is essential to understanding the presence and frequency of consequential variants and their patterns and trajectory of dissemination."

- High prevalence of SARS-CoV-2 antibodies in care homes affected by COVID-19: Prospective cohort study, England.

(/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=851)

Ladhani Shamez N et al. EclinicalMedicine 2020 11 100597

"Seropositivity across these seven homes [with no reported COVID cases] varied between 10.7-56.5%, with four exceeding community seroprevalence in London (14.8%). Seropositivity rates for staff and residents correlated significantly (r_s 0.84, [95% CI 0.51-0.95] $p < 0.001$) across the 13 homes. WGS identified multiple introductions into some homes and silent transmission of a single lineage between staff and residents in one home.

- Estimated transmissibility and impact of SARS-CoV-2 lineage B.1.1.7 in England (/PHGKB/phgHome.action?

action=forward&dbsource=covUpdate&id=858)

NG Davies et al, Science, March 3, 2021

Using statistical and dynamic modelling approaches, we estimate that this variant has a 43–90%, higher reproduction number than preexisting variants. A fitted two-strain dynamic transmission model shows that VOC 202012/01 will lead to large resurgences of COVID-19 cases. Without stringent control measures, including limited closure of educational institutions and an accelerated vaccine roll-out, COVID-19 hospitalizations and deaths across England in 2021 will exceed those in 2020.

- Delayed Large Local Reactions to mRNA-1273 Vaccine against SARS-CoV-2. (/PHGKB/phgHome.action?

action=forward&dbsource=covUpdate&id=861)

Blumenthal Kimberly G et al. The New England journal of medicine 2021 3

We have observed delayed large local reactions to the mRNA-1273 vaccine, with a median onset on day 8 (range, 4 to 11) after the first dose. These reactions had a variable appearance. Here, we report on a series of 12 patients with these reactions, all of which appeared near the injection site after complete resolution of the initial local and systemic symptoms associated with vaccination.

Non-Genomics Precision Health Studies

- Report Outlines Strategy for Surveillance of Emerging Coronavirus Variants (/PHGKB/phgHome.action?

action=forward&dbsource=covUpdate&id=813)

Stephenson J. JAMA Network, Feb 23, 2021.

- COVID-19 disruptions to endoscopic surveillance in Lynch syndrome. (/PHGKB/phgHome.action?

action=forward&dbsource=covUpdate&id=814)

McKenna Danielle B et al. Cancer prevention research (Philadelphia, Pa.) 2021 2

This study demonstrates that the COVID-19 pandemic shutdown led to delayed endoscopic surveillance in Lynch syndrome, with potentially impactful delays among young patients. This data also emphasizes the importance of timely surveillance in Lynch syndrome during this current, as well as potential future, global pandemics.

- Microbial signatures in the lower airways of mechanically ventilated COVID19 patients associated with poor clinical outcome

(/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=817)

I Sulaiman et al, MEDRXIV, February 26, 2021

We found that isolation of a hospital-acquired respiratory pathogen was not associated with fatal outcome. However, poor clinical outcome was associated with enrichment of the lower airway microbiota with an oral commensal (*Mycoplasma salivarium*), while high SARS-CoV-2 viral burden, poor anti-SARS-CoV-2 antibody response, together with a unique host transcriptome profile of the lower airways were most predictive of mortality.

- A Novel SARS-CoV-2 Variant of Concern, B.1.526, Identified in New York (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=818)
MK Annavajhala et al, MEDRXIV, February 25, 2021

To survey for these novel variants in our patient population in New York City, PCR assays were designed to identify viruses with two signature mutations, E484K and N501Y. We observed a steady increase in the detection rate from late December to mid-February, with an alarming rise to 12.3% in the past two weeks.

- A Neanderthal OAS1 isoform protects individuals of European ancestry against COVID-19 susceptibility and severity (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=819)
S Zhou et al, Nature Medicine, February 25, 2021

To identify circulating proteins influencing Coronavirus Disease 2019 (COVID-19) susceptibility and severity, we undertook a two-sample Mendelian randomization (MR) study, rapidly scanning hundreds of circulating proteins while reducing bias due to reverse causation and confounding. In up to 14,134 cases and 1.2 million controls, we found that an s.d. increase in OAS1 levels was associated with reduced COVID-19 death or ventilation, hospitalization and susceptibility.

- Common genetic variation in humans impacts in vitro susceptibility to SARS-CoV-2 infection. (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=831)
Dobrindt Kristina et al. Stem cell reports 2021 2

We hypothesize that, in addition to viral load and host antibody repertoire, host genetic variants influence vulnerability to infection. Here we apply human induced pluripotent stem cell (hiPSC)-based models and CRISPR engineering to explore the host genetics of SARS-CoV-2. We demonstrate that a single-nucleotide polymorphism (rs4702), common in the population and located in the 3' UTR of the protease *FURIN*, influences alveolar and neuron infection by SARS-CoV-2 in vitro.

- Escape of SARS-CoV-2 501Y.V2 variants from neutralization by convalescent plasma (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=832)
S Cele et al, MEDRXIV, February 27, 2021

Here, using a live virus neutralization assay (LVNA), we compared neutralization of a first wave virus (B.1.1.117) versus the 501Y.V2 variant using plasma collected from adults hospitalized with COVID-19 from two South African infection waves, with the second wave dominated by 501Y.V2 infections.

- Increased hazard of mortality in cases compatible with SARS-CoV-2 variant of concern 202012/1 - a matched cohort study (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=833)
R Challen et al, MEDRXIV, February 27, 2021

In this matched cohort study of 54,773 pairs of participants matched by age, gender, ethnicity, index of multiple deprivation, lower tier local authority region, and sample date of positive specimen, and differing only by detectability of the spike protein gene. Death within 28 days after first positive test was higher among patients with VOC-202012/01. (hazard ratio 1.7, 95% CI 1.3-2.2).

- A statewide analysis of SARS-CoV-2 transmission in New York (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=836)
EL Nesselquist et al, MEDRXIV, February 28, 2021

The Wadsworth Center sought to characterize the transmission dynamics of SARS-CoV-2 across the entire state of New York from March to September with the addition of over 600 genomes from under-sampled and previously unsampled New York counties and to more fully understand the breadth of the initial outbreak in Westchester County. Additional sequencing confirmed the dominance of B.1 and descendant lineages (collectively referred to as B.1.X) in New York State.

- Medical vulnerability of individuals with Down syndrome to severe COVID-19-data from the Trisomy 21 Research Society and the UK ISARIC4C survey (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=838)
A Huls et al, Ebiomedicine, February 2021

Health conditions, immune dysfunction, and premature aging associated with trisomy 21 (Down syndrome, DS) may impact the clinical course of COVID-19. The T21RS COVID-19 Initiative launched an international survey for clinicians or caregivers on patients with COVID-19 and DS. Data collected between April and October 2020 (N=1046) were analyzed and compared with the UK ISARIC4C survey of hospitalized COVID-19 patients with and without DS.

- Genetics of symptom remission in outpatients with COVID-19 (/PHGKB/phgHome.action?)

action=forward&dbsource=covUpdate&id=846)

MP Dube et al, MEDRXIV, March 1, 2021

We conducted a genome-wide association study of time to remission of COVID-19 symptoms in 1723 outpatients with at least one risk factor for disease severity from the COLCORONA clinical trial. We found a significant association at 5p13.3 (rs1173773; $P = 4.94 \times 10^{-8}$) near the natriuretic peptide receptor 3 gene (NPR3). By day 15 of the study, 44%, 54% and 59% of participants with 0, 1, or 2 copies of the effect allele respectively, had symptom remission.

- Transmission of SARS-CoV-2 in domestic cats imposes a narrow bottleneck. (/PHGKB/phgHome.action?

action=forward&dbsource=covUpdate&id=849)

Braun Katarina M et al. PLoS pathogens 2021 2 (2) e1009373

"In RNA virus evolution, stochastic processes like narrow transmission bottlenecks and genetic drift typically act to constrain the overall pace of adaptive evolution. Our data suggest that here, positive selection in index cats followed by a narrow transmission bottleneck may have instead accelerated the fixation of S H655Y, a potentially beneficial SARS-CoV-2 variant. Overall, our study suggests species- and context-specific adaptations are likely to continue to emerge. This underscores the importance of continued genomic surveillance for new SARS-CoV-2 variants as well as heightened scrutiny for signatures of SARS-CoV-2 positive selection in humans and mammalian model systems."

- Sequence Analysis of 20,453 SARS-CoV-2 Genomes from the Houston Metropolitan Area Identifies the Emergence and Widespread Distribution of Multiple Isolates of All Major Variants of Concern (/PHGKB/phgHome.action?

action=forward&dbsource=covUpdate&id=850)

Long SW, et al. medRxiv, Mar 2, 2021.

"Houston is the first city in the United States to have all variants documented by genome sequencing. As vaccine deployment accelerates worldwide, increased genomic surveillance of SARS-CoV-2 is essential to understanding the presence and frequency of consequential variants and their patterns and trajectory of dissemination."

- High prevalence of SARS-CoV-2 antibodies in care homes affected by COVID-19: Prospective cohort study, England.

(/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=851)

Ladhani Shamez N et al. EClinicalMedicine 2020 11 100597

"Seropositivity across these seven homes [with no reported COVID cases] varied between 10.7-56.5%, with four exceeding community seroprevalence in London (14.8%). Seropositivity rates for staff and residents correlated significantly ($r_s 0.84$, [95% CI 0.51-0.95] $p < 0.001$) across the 13 homes. WGS identified multiple introductions into some homes and silent transmission of a single lineage between staff and residents in one home.

- Estimated transmissibility and impact of SARS-CoV-2 lineage B.1.1.7 in England (/PHGKB/phgHome.action?

action=forward&dbsource=covUpdate&id=858)

NG Davies et al, Science, March 3, 2021

Using statistical and dynamic modelling approaches, we estimate that this variant has a 43–90%, higher reproduction number than preexisting variants. A fitted two-strain dynamic transmission model shows that VOC 202012/01 will lead to large resurgences of COVID-19 cases. Without stringent control measures, including limited closure of educational institutions and an accelerated vaccine roll-out, COVID-19 hospitalizations and deaths across England in 2021 will exceed those in 2020.

- Delayed Large Local Reactions to mRNA-1273 Vaccine against SARS-CoV-2. (/PHGKB/phgHome.action?

action=forward&dbsource=covUpdate&id=861)

Blumenthal Kimberly G et al. The New England journal of medicine 2021 3

We have observed delayed large local reactions to the mRNA-1273 vaccine, with a median onset on day 8 (range, 4 to 11) after the first dose. These reactions had a variable appearance. Here, we report on a series of 12 patients with these reactions, all of which appeared near the injection site after complete resolution of the initial local and systemic symptoms associated with vaccination.

News, Reviews and Commentaries

- Report Outlines Strategy for Surveillance of Emerging Coronavirus Variants (/PHGKB/phgHome.action?

action=forward&dbsource=covUpdate&id=813)

Stephenson J. JAMA Network, Feb 23, 2021.

- COVID-19 disruptions to endoscopic surveillance in Lynch syndrome. (/PHGKB/phgHome.action?

action=forward&dbsource=covUpdate&id=814)

McKenna Danielle B et al. Cancer prevention research (Philadelphia, Pa.) 2021 2

This study demonstrates that the COVID-19 pandemic shutdown led to delayed endoscopic surveillance in Lynch syndrome, with potentially impactful delays among young patients. This data also emphasizes the importance of timely surveillance in Lynch syndrome during this current, as well as potential future, global pandemics.

- Microbial signatures in the lower airways of mechanically ventilated COVID19 patients associated with poor clinical outcome (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=817)

I Sulaiman et al, MEDRXIV, February 26, 2021

We found that isolation of a hospital-acquired respiratory pathogen was not associated with fatal outcome. However, poor clinical outcome was associated with enrichment of the lower airway microbiota with an oral commensal (*Mycoplasma salivarium*), while high SARS-CoV-2 viral burden, poor anti-SARS-CoV-2 antibody response, together with a unique host transcriptome profile of the lower airways were most predictive of mortality.

- A Novel SARS-CoV-2 Variant of Concern, B.1.526, Identified in New York (/PHGKB/phgHome.action?

action=forward&dbsource=covUpdate&id=818)

MK Annavajhala et al, MEDRXIV, February 25, 2021

To survey for these novel variants in our patient population in New York City, PCR assays were designed to identify viruses with two signature mutations, E484K and N501Y. We observed a steady increase in the detection rate from late December to mid-February, with an alarming rise to 12.3% in the past two weeks.

- A Neanderthal OAS1 isoform protects individuals of European ancestry against COVID-19 susceptibility and severity (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=819)

S Zhou et al, Nature Medicine, February 25, 2021

To identify circulating proteins influencing Coronavirus Disease 2019 (COVID-19) susceptibility and severity, we undertook a two-sample Mendelian randomization (MR) study, rapidly scanning hundreds of circulating proteins while reducing bias due to reverse causation and confounding. In up to 14,134 cases and 1.2 million controls, we found that an s.d. increase in OAS1 levels was associated with reduced COVID-19 death or ventilation, hospitalization and susceptibility.

- Common genetic variation in humans impacts in vitro susceptibility to SARS-CoV-2 infection. (/PHGKB/phgHome.action?

action=forward&dbsource=covUpdate&id=831)

Dobrindt Kristina et al. Stem cell reports 2021 2

We hypothesize that, in addition to viral load and host antibody repertoire, host genetic variants influence vulnerability to infection. Here we apply human induced pluripotent stem cell (hiPSC)-based models and CRISPR engineering to explore the host genetics of SARS-CoV-2. We demonstrate that a single-nucleotide polymorphism (rs4702), common in the population and located in the 3' UTR of the protease *FURIN*, influences alveolar and neuron infection by SARS-CoV-2 in vitro.

- Escape of SARS-CoV-2 501Y.V2 variants from neutralization by convalescent plasma (/PHGKB/phgHome.action?

action=forward&dbsource=covUpdate&id=832)

S Cele et al, MEDRXIV, February 27, 2021

Here, using a live virus neutralization assay (LVNA), we compared neutralization of a first wave virus (B.1.1.117) versus the 501Y.V2 variant using plasma collected from adults hospitalized with COVID-19 from two South African infection waves, with the second wave dominated by 501Y.V2 infections.

- Increased hazard of mortality in cases compatible with SARS-CoV-2 variant of concern 202012/1 - a matched cohort study (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=833)

R Challen et al, MEDRXIV< February 27, 2021

In this matched cohort study of 54,773 pairs of participants matched by age, gender, ethnicity, index of multiple deprivation, lower tier local authority region, and sample date of positive specimen, and differing only by detectability of the spike protein gene. Death within 28 days after first positive test was higher among patients with VOC-202012/01. (hazard ratio 1.7, 95%

CI 1.3-2.2).

- A statewide analysis of SARS-CoV-2 transmission in New York (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=836)
EL Nesselquist et al, MEDRXIV, February 28, 2021

The Wadsworth Center sought to characterize the transmission dynamics of SARS-CoV-2 across the entire state of New York from March to September with the addition of over 600 genomes from under-sampled and previously unsampled New York counties and to more fully understand the breadth of the initial outbreak in Westchester County. Additional sequencing confirmed the dominance of B.1 and descendant lineages (collectively referred to as B.1.X) in New York State.

- Medical vulnerability of individuals with Down syndrome to severe COVID-19-data from the Trisomy 21 Research Society and the UK ISARIC4C survey (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=838)
A Huls et al, Ebiomedicine, February 2021

Health conditions, immune dysfunction, and premature aging associated with trisomy 21 (Down syndrome, DS) may impact the clinical course of COVID-19. The T21RS COVID-19 Initiative launched an international survey for clinicians or caregivers on patients with COVID-19 and DS. Data collected between April and October 2020 (N=1046) were analyzed and compared with the UK ISARIC4C survey of hospitalized COVID-19 patients with and without DS.

- Genetics of symptom remission in outpatients with COVID-19 (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=846)
MP Dube et al, MEDRXIV, March 1, 2021

We conducted a genome-wide association study of time to remission of COVID-19 symptoms in 1723 outpatients with at least one risk factor for disease severity from the COLCORONA clinical trial. We found a significant association at 5p13.3 (rs1173773; $P = 4.94 \times 10^{-8}$) near the natriuretic peptide receptor 3 gene (NPR3). By day 15 of the study, 44%, 54% and 59% of participants with 0, 1, or 2 copies of the effect allele respectively, had symptom remission.

- Transmission of SARS-CoV-2 in domestic cats imposes a narrow bottleneck. (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=849)
Braun Katarina M et al. PLoS pathogens 2021 2 (2) e1009373

"In RNA virus evolution, stochastic processes like narrow transmission bottlenecks and genetic drift typically act to constrain the overall pace of adaptive evolution. Our data suggest that here, positive selection in index cats followed by a narrow transmission bottleneck may have instead accelerated the fixation of S H655Y, a potentially beneficial SARS-CoV-2 variant. Overall, our study suggests species- and context-specific adaptations are likely to continue to emerge. This underscores the importance of continued genomic surveillance for new SARS-CoV-2 variants as well as heightened scrutiny for signatures of SARS-CoV-2 positive selection in humans and mammalian model systems."

- Sequence Analysis of 20,453 SARS-CoV-2 Genomes from the Houston Metropolitan Area Identifies the Emergence and Widespread Distribution of Multiple Isolates of All Major Variants of Concern (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=850)
Long SW, et al. medRxiv, Mar 2, 2021.

"Houston is the first city in the United States to have all variants documented by genome sequencing. As vaccine deployment accelerates worldwide, increased genomic surveillance of SARS-CoV-2 is essential to understanding the presence and frequency of consequential variants and their patterns and trajectory of dissemination."

- High prevalence of SARS-CoV-2 antibodies in care homes affected by COVID-19: Prospective cohort study, England. (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=851)
Ladhani Shamez N et al. EClinicalMedicine 2020 11 100597

"Seropositivity across these seven homes [with no reported COVID cases] varied between 10.7-56.5%, with four exceeding community seroprevalence in London (14.8%). Seropositivity rates for staff and residents correlated significantly (rs 0.84, [95% CI 0.51-0.95] $p < 0.001$) across the 13 homes. WGS identified multiple introductions into some homes and silent transmission of a single lineage between staff and residents in one home.

- Estimated transmissibility and impact of SARS-CoV-2 lineage B.1.1.7 in England (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=858)
NG Davies et al, Science, March 3, 2021

Using statistical and dynamic modelling approaches, we estimate that this variant has a 43–90%, higher reproduction number than preexisting variants. A fitted two-strain dynamic transmission model shows that VOC 202012/01 will lead to large resurgences of COVID-19 cases. Without stringent control measures, including limited closure of educational institutions and an accelerated vaccine roll-out, COVID-19 hospitalizations and deaths across England in 2021 will exceed those in 2020.

- Delayed Large Local Reactions to mRNA-1273 Vaccine against SARS-CoV-2. (/PHGKB/phgHome.action?action=forward&dbsource=covUpdate&id=861)

Blumenthal Kimberly G et al. The New England journal of medicine 2021 3

We have observed delayed large local reactions to the mRNA-1273 vaccine, with a median onset on day 8 (range, 4 to 11) after the first dose. These reactions had a variable appearance. Here, we report on a series of 12 patients with these reactions, all of which appeared near the injection site after complete resolution of the initial local and systemic symptoms associated with vaccination.

Disclaimer: Articles listed in COVID-19 Genomics and Precision Public Health Weekly Update are selected by the CDC Office of Public Health Genomics to provide current awareness of the scientific literature and news. Inclusion in the update does not necessarily represent the views of the Centers for Disease Control and Prevention nor does it imply endorsement of the article's methods or findings. CDC and DHHS assume no responsibility for the factual accuracy of the items presented. The selection, omission, or content of items does not imply any endorsement or other position taken by CDC or DHHS. Opinion, findings and conclusions expressed by the original authors of items included in the Clips, or persons quoted therein, are strictly their own and are in no way meant to represent the opinion or views of CDC or DHHS. References to publications, news sources, and non-CDC Websites are provided solely for informational purposes and do not imply endorsement by CDC or DHHS.

Page last reviewed: Oct 1, 2020

Page last updated: Mar 28, 2021

Content source: Office of Genomics and Precision Public Health (<http://www.cdc.gov/genomics/>), CDC Office of Science (<https://www.cdc.gov/od/science/index.htm>)