


FIGURE. Estimated proportion of adults aged ≥19 years who received selected vaccines, by age group and risk status – National Health Interview Survey, United States, 2010–2018

	Unweighted sample size (2010–2018)	2010	2011	2012	2013	2014	2015	2016	2017	2018	Average annual change (95% CI)	Linear trend test p-value
Influenza - age ≥19 yrs	252,236	37.2	38.5	39.6	42.8	43.2	44.8	43.5	45.4	46.1	1.1 (0.8, 1.5)	<0.01
Influenza - age ≥19 yrs, high risk	68,818	54.7	56.6	56.6	59.9	58.0	58.8	58.8	59.7	61.0	0.6 (0.3, 1.0)	<0.01
Pneumococcal - age 19–64 yrs, increased risk	72,651	18.5	20.1	20.0	21.2	20.3	23.0	24.0	24.5	23.3	0.7 (0.5, 0.9)	<0.01
Pneumococcal - age ≥65 yrs	65,428	59.7	62.3	59.9	59.7	61.3	63.6	66.9	69.0	69.0	1.3 (1.1, 1.5)	<0.01
Herpes zoster - age ≥60 yrs	89,209	14.4	15.8	20.1	24.2	27.9	30.6	33.4	34.9	34.5	2.8 (2.6, 3.0)	<0.01
Tetanus-toxoid (Td or Tdap) - age ≥19 yrs	266,638	62.0	62.6	62.4	62.0	62.2	61.6	62.2	63.4	62.9	0.1 (-0.0, 0.2)	0.15
Tdap* - age ≥19 yrs	179,114	6.9	10.5	14.2	17.2	20.1	23.1	26.6	31.7	31.2	3.2 (3.0, 3.4)	<0.01



Abbreviations: Td = tetanus and diphtheria toxoids; Tdap = tetanus toxoid, reduced diphtheria toxoid, and acellular pertussis vaccine.

* Tdap vaccination coverage data among adults aged ≥65 years are available beginning in the NHIS 2012 survey.