



Centers for Disease Control and Prevention
CDC 24/7: Saving Lives, Protecting People™



COVID-19

SARS-CoV-2 Variant Classifications and Definitions

Updated June 4, 2021

[Print](#)

Key Points:

- Genetic variants of SARS-CoV-2 have been emerging and circulating around the world throughout the COVID-19 pandemic.
- Viral mutations and variants in the United States are routinely monitored through sequence-based surveillance, laboratory studies, and epidemiological investigations.
- A US government interagency group developed a Variant Classification scheme that defines three classes of SARS-CoV-2 variants:
 - [Variant of Interest](#)
 - [Variant of Concern](#)
 - [Variant of High Consequence](#)
- The B.1.1.7, B.1.351, P.1, B.1.427, and B.1.429 variants circulating in the United States are classified as variants of concern.
- To date, no variants of high consequence have been identified in the United States.
- Laboratory studies suggest specific monoclonal antibody treatments may be less effective for treating cases of COVID-19 caused by variants with the L452R or E484K substitution in the spike protein, the combination of K417N, E484K, and N501Y, or the combination of K417T, E484K, and N501Y substitutions in the spike protein.
 - L452R is present in B.1.526.1, B.1.427, B.1.429, and B.1.617 lineages and sub-lineages.
 - E484K is present in B.1.525, P.2, P.1, and B.1.351, but only some strains of B.1.526 and B.1.1.7.
 - The combination of K417N, E484K, and N501Y substitutions is present in B.1.351.
 - The combination of K417T, E484K, and N501Y substitutions is present in P.1.

Viruses constantly change through mutation. A variant has one or more mutations that differentiate it from other variants in circulation. As expected, multiple variants of SARS-CoV-2 have been documented in the [United States](#) and [globally](#) throughout this pandemic. To inform local outbreak investigations and understand national trends, scientists compare genetic differences between viruses to identify variants and how they are related to each other.

Variant classifications

The US Department of Health and Human Services (HHS) established a SARS-CoV-2 Interagency Group (SIG) to improve coordination among the Centers for Disease Control and Prevention (CDC), National Institutes of Health (NIH), Food and Drug Administration (FDA), Biomedical Advanced Research and Development Authority (BARDA), and Department of Defense (DoD). This interagency group is focused on the rapid characterization of emerging variants and actively monitors their potential impact on critical SARS-CoV-2 countermeasures, including vaccines, therapeutics, and diagnostics.

- [Variants of Interest \(VOI\)](#)
- [Variants of Concern \(VOC\)](#)
- [Variants of High Consequence \(VOHC\)](#)

Notes: Each classification of variant includes the possible attributes of lower classes (i.e., VOC includes the possible attributes of VOI); variant status might escalate or deescalate based on scientific evidence. This page will be updated as needed to show the

variants that belong to each class. The [World Health Organization](#) (WHO) also classifies variant viruses as Variants of Concern and Variants of Interest; US classifications may differ from those of WHO since the importance of variants may differ by location.

See [Variant Proportions in the U.S.](#)

Variant of Interest

A variant with specific genetic markers that have been associated with changes to receptor binding, reduced neutralization by antibodies generated against previous infection or vaccination, reduced efficacy of treatments, potential diagnostic impact, or predicted increase in transmissibility or disease severity.

Possible attributes of a variant of interest:

- Specific genetic markers that are predicted to affect transmission, diagnostics, therapeutics, or immune escape
- Evidence that it is the cause of an increased proportion of cases or unique outbreak clusters
- Limited prevalence or expansion in the US or in other countries

A variant of interest might require one or more appropriate public health actions, including enhanced sequence surveillance, enhanced laboratory characterization, or epidemiological investigations to assess how easily the virus spreads to others, the severity of disease, the efficacy of therapeutics and whether currently authorized vaccines offer protection.

Current variants of interest in the United States that are being monitored and characterized are listed in the table below. The table will be updated when a new variant of interest is identified.

Selected Characteristics of SARS-CoV-2 Variants of Interest⁺

Name (Pango lineage ) ^a	Spike Protein Substitutions	Name (Nextstrain ) ^b	WHO Label ^d	First Detected	Relevance
B.1.525	Spike: A67V, 69del, 70del, 144del, E484K, D614G, Q677H, F888L	20A/S:484K	Eta	United Kingdom/Nigeria – December 2020	•
B.1.526	Spike: (L5F*), T95I, D253G, (S477N*), (E484K*), D614G, (A701V*)	20C/S:484K	Iota	United States (New York) – November 2020	•

B.1.526.1	Spike: D80G, 144del, F157S, L452R, D614G, (T791I*), (T859N*), D950H	20C		United States (New York) – October 2020	
B.1.617	Spike: L452R, E484Q, D614G	20A		India – February 2021	
B.1.617.1	Spike: (T95I), G142D, E154K, L452R, E484Q, D614G, P681R, Q1071H	20A/S:154K	Kappa	India – December 2020	

B.1.617.2	Spike: T19R, (G142D), 156del, 157del, R158G, L452R, T478K, D614G, P681R, D950N	20A/S:478K	Delta	India – December 2020	
B.1.617.3	Spike: T19R, G142D, L452R, E484Q, D614G, P681R, D950N	20A		India – October 2020	
P.2	Spike: E484K, (F565L*), D614G, V1176F	20J	Zeta	Brazil – April 2020	

--	--	--	--	--	--

(*) = detected in some sequences but not all

** These variants share one specific mutation called D614G. This mutation was one of the first documented in the US in the initial stages of the pandemic, after having initially circulated in Europe^[13]. There is evidence that variants with this mutation spread more quickly than viruses without this mutation^[12] [\[12\]](#).*

a – Phylogenetic Assignment of Named Global Outbreak (PANGO) Lineages is software tool developed by members of the Rambaut Lab. The associated web application was developed by the Centre for Genomic Pathogen Surveillance in South Cambridgeshire and is intended to implement the dynamic nomenclature of SARS-CoV-2 lineages, known as the PANGO nomenclature.

b – Nextstrain, a collaboration between researchers in Seattle, USA and Basel, Switzerland, provides open-source tools for visualizing the genetics of outbreaks. The goal is to support public health surveillance by facilitating understanding of the spread and evolution of pathogens.

c – The Biodefense and Emerging Infections Research Resources (BEI Resources) is a NIAID-funded repository to provide reagents, tools, and information to the research community. The reference viruses proposed here facilitate the harmonization of information among all stakeholders in the COVID-19 pandemic research community. Please note that the reference viruses provided in the tables below are based on what is currently available through the BEI Resources.

d – To assist with public discussions of variants, WHO proposed using labels consisting of the Greek Alphabet, i.e., Alpha, Beta, Gamma, as a practical way to discuss variants by non-scientific audiences. [Tracking SARS-CoV-2 variants \(who.int\)](#) [\[14\]](#)

Variant of Concern

A variant for which there is evidence of an increase in transmissibility, more severe disease (e.g., increased hospitalizations or deaths), significant reduction in neutralization by antibodies generated during previous infection or

vaccination, reduced effectiveness of treatments or vaccines, or diagnostic detection failures.

Possible attributes of a variant of concern:

In addition to the possible attributes of a variant of interest

- Evidence of impact on diagnostics, treatments, or vaccines
 - Widespread interference with diagnostic test targets
 - Evidence of substantially decreased susceptibility to one or more class of therapies
 - Evidence of significant decreased neutralization by antibodies generated during previous infection or vaccination
 - Evidence of reduced vaccine-induced protection from severe disease
- Evidence of increased transmissibility
- Evidence of increased disease severity

Variants of concern might require one or more appropriate public health actions, such as notification to WHO under the International Health Regulations, reporting to CDC, local or regional efforts to control spread, increased testing, or research to determine the effectiveness of vaccines and treatments against the variant. Based on the characteristics of the variant, additional considerations may include the development of new diagnostics or the modification of vaccines or treatments.

Current variants of concern in the United States that are being closely monitored and characterized by federal agencies are included in the table below. The table will be updated when a new variant of concern is identified.

Selected Characteristics of SARS-CoV-2 Variants of Concern⁺

Name (Pango lineage) ^a	Spike Protein Substitutions	Name (Nextstrain) ^b	WHO Label ^c	First Detected	BEI  Reference Isolate
B.1.1.7	69del, 70del, 144del, (E484K*), (S494P*), N501Y, A570D, D614G, P681H, T716I, S982A,	20I/501Y.V1	Alpha	United Kingdom	NR-540C 

	D1118H (K1191N*)					•
						•
B.1.351	D80A, D215G, 241del, 242del, 243del, K417N, E484K, N501Y, D614G, A701V	20H/501.V2	Beta	South Africa	NR-5528 	• •
						•
B.1.427	L452R, D614G	20C/S:452R	Epsilon	United States- (California)		• •

●

* These variants share one specific mutation called D614G. This mutation was one of the first documented in the US in the initial stages of the pandemic, after having initially circulated in Europe^[13]. There is evidence that variants with this mutation spread more quickly than viruses without this mutation^[12].

b – Nextstrain, a collaboration between researchers in Seattle, USA and Basel, Switzerland, provides open-source tools for visualizing the genetics of outbreaks. The goal is to support public health surveillance by facilitating understanding of the spread and evolution of pathogens.

d – The Biodefense and Emerging Infections Research Resources (BEI Resources) is a NIAID-funded repository to provide reagents, tools, and information to the research community. The reference viruses proposed here facilitate the harmonization of information among all stakeholders in the COVID-19 pandemic research community. Please note that the reference viruses provided in the tables below are based on what is currently available through the BEI resources.

e – Attributes listed are based on data available from pseudoviruses or recombinant viruses containing combinations of substitutions characteristic of specific lineages or from reference virus isolates.

Variant of High Consequence

A variant of high consequence has clear evidence that prevention measures or medical countermeasures (MCMs) have significantly reduced effectiveness relative to previously circulating variants.

Possible attributes of a variant of high consequence:

In addition to the possible attributes of a variant of concern

- Impact on Medical Countermeasures (MCM)
 - Demonstrated failure of diagnostics
 - Evidence to suggest a significantly reduction in vaccine effectiveness, a disproportionately high number of vaccine breakthrough cases, or very low vaccine-induced protection against severe disease
 - Significantly reduced susceptibility to multiple Emergency Use Authorization (EUA) or approved therapeutics
 - More severe clinical disease and increased hospitalizations

A variant of high consequence would require notification to WHO under the International Health Regulations, reporting to CDC, an announcement of strategies to prevent or contain transmission, and recommendations to update treatments and vaccines.

Currently there are no SARS-CoV-2 variants that rise to the level of high consequence.

Treatment considerations for healthcare providers

Substitutions of Concern for SARS-CoV-2 Monoclonal Antibody Therapies

In the United States, there are two anti-SARS-CoV-2 monoclonal antibody treatments with FDA Emergency Use Authorization (EUA) for the treatment of COVID-19: [bamlanivimab plus etesevimab](#) [↗](#) and [casirivimab plus imdevimab](#). [↗](#)

CDC's national genomic surveillance program identifies new and emerging SARS-CoV-2 variants to determine implications for COVID-19 diagnostics, treatments, or vaccines authorized for use in the United States. Sequences with similar genetic changes are grouped into lineages, and multiple lineages can have the same substitutions. For example, the E484K substitution is found in lineages B.1.351, P.1, B.1.526, and many others. Genomic surveillance efforts provide the capability to detect viruses that have reduced susceptibility to treatments more quickly.

In laboratory studies, SARS-CoV-2 variants that contain certain substitutions in the spike protein cause a marked reduction in susceptibility to bamlanivimab and may have reduced sensitivity to etesevimab and casirivimab. The L452R substitution found in the B.1.427 and B.1.429 variants has been shown to cause a significant reduction in susceptibility to bamlanivimab. The L452R substitution also causes a modest decrease in susceptibility to the combination of bamlanivimab and etesevimab, although the clinical implications of this decrease are not known.⁷ In addition, the E484K substitution, which is found in the B.1.351, P.1, and B.1.526 variants, also results in a marked reduction in susceptibility to bamlanivimab, and laboratory studies suggest that the B.1.351, P.1, and B.1.526 variants have markedly reduced susceptibility to the combination of bamlanivimab and etesevimab.⁷ Laboratory studies also suggest that the K417N and K417T substitutions, which are present in the B.1.351 and P.1 variants, respectively, along with the E484K mutation, reduces virus susceptibility to casirivimab, although the combination of casirivimab and imdevimab appears to retain activity.¹⁴

The table below shows the national and regional unweighted proportions of SARS-CoV-2 that contain the L452R or E484K substitution, individually, as well as the unweighted proportions of SARS-CoV-2 that contain the combination of K417N, E484K, and N501Y substitutions or the combination of K417T, E484K, and N501Y substitutions. As new data become available, additional substitutions may be added to the table below. The national and regional proportions provided in the table below will be updated weekly.

Resources

[Monoclonal Antibody COVID-19 Infusion](#) [↗](#)

[Statement on Anti-SARS-CoV-2 Monoclonal Antibodies EUA | COVID-19 Treatment Guidelines \(nih.gov\)](#) [↗](#)

Unweighted Proportions of SARS-CoV-2 Substitutions of Therapeutic Concern

Spike Protein Substitution	National Proportion ^a	Regional Proportions ^b		Common Pango Lineages with Spike Protein Substitutions ^c
L452R	6.9%	Region 1	6.5%	B.1.526.1 B.1.429 B.1.427 B.1.617.2 B.1 B.1.617.1 C.36 A.2.5
		Region 2	7.2%	
		Region 3	7.1%	
		Region 4	4.3%	
		Region 5	5.9%	
		Region 6	4.5%	
		Region 7	3.8%	
		Region 8	9.0%	
		Region 9	13.3%	
		Region 10	13.8%	
E484K	16.3%	Region 1	25.5%	P.1 B.1.526 B.1.1.318 B.1.351 B.1.525 B.1.621 R.1 B.1
		Region 2	22.5%	
		Region 3	14.0%	
		Region 4	16.3%	
		Region 5	13.0%	
		Region 6	11.4%	
		Region 7	12.9%	
		Region 8	9.2%	
		Region		

		9	17.9%	
		Region 10	14.7%	
K417N, E484K, N501Y	0.6%	Region 1	0.1%	B.1.351 B.1
		Region 2	0.4%	
		Region 3	0.7%	
		Region 4	0.5%	
		Region 5	0.7%	
		Region 6	0.2%	
		Region 7	1.3%	
		Region 8	0.7%	
		Region 9	1.4%	
		Region 10	0.9%	
K417T, E484K, N501Y	7.9%	Region 1	12.4%	P.1
		Region 2	5.3%	
		Region 3	2.2%	
		Region 4	8.0%	
		Region 5	8.3%	
		Region 6	5.4%	
		Region 7	8.3%	
		Region 8	3.9%	
		Region 9	10.2%	
		Region 10	9.4%	

a – The unweighted proportion of SARS-CoV-2 circulating in the United States that contain the designated substitution, based on >34,000 sequences collected through CDC's national genomic surveillance during the two-week period ending May 8, 2021.

b – The unweighted regional proportion of SARS-CoV-2 circulating in each HHS region that contain the designated substitution, based on >34,000 sequences collected through CDC's national genomic surveillance during the two-week period ending May 8, 2021.

c – The lineages listed are the most common lineages within CDC's national genomic surveillance with these substitutions, but this list is not intended to be a complete list of the lineages that contain the spike protein substitutions.

References

1. Zhou, B., Thi Nhu Thao, T., Hoffmann, D. et al. SARS-CoV-2 spike D614G change enhances replication and transmission. *Nature* (2021). <https://doi.org/10.1038/s41586-021-03361-1> 
2. Volz E, Hill V, McCrone J, et al. Evaluating the Effects of SARS-CoV-2 Spike Mutation D614G on Transmissibility and Pathogenicity. *Cell* 2021; 184(64-75). doi: <https://doi.org/10.1016/j.cell.2020.11.020> 
3. Korber B, Fischer WM, Gnanakaran S, et al. Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G Increases Infectivity of the COVID-19 Virus. *Cell* 2021; 182(812-7) doi: <https://doi.org/10.1016/j.cell.2020.06.043> 
4. Yurkovetskiy L, Wang X, Pascal KE, et al. Structural and Functional Analysis of the D614G SARS-CoV-2 Spike Protein Variant. *Cell* 2020; 183(3): 739-751. doi: <https://doi.org/10.1016/j.cell.2020.09.032> 
5. *Davies NG, Abbott S, Barnard RC, et al. Estimated transmissibility and impact of SARS-CoV-2 lineage B.1.1.7 in England. *MedRxiv* 2021. doi: <https://doi.org/10.1101/2020.12.24.20248822> 
6. Horby P, Huntley C, Davies N et al. NERVTAG note on B.1.1.7 severity. New & Emerging Threats Advisory Group, Jan. 21, 2021. Retrieved from [NERVTAG note on variant severity](#) 
7. [Fact Sheet For Health Care Providers Emergency Use Authorization \(Eua\) Of Bamlanivimab And Etesevimab 02092021 \(fda.gov\)](#) 
8. *Wang P, Nair MS, Liu L, et al. Antibody Resistance of SARS-CoV-2 Variants B.1.351 and B.1.1.7. *BioRxiv* 2021. doi: <https://doi.org/10.1101/2021.01.25.428137> 
9. *Shen X, Tang H, McDanal C, et al. SARS-CoV-2 variant B.1.1.7 is susceptible to neutralizing antibodies elicited by ancestral Spike vaccines. *BioRxiv* 2021. doi: <https://doi.org/10.1101/2021.01.27.428516> 
10. *Edara VV, Floyd K, Lai L, et al. Infection and mRNA-1273 vaccine antibodies neutralize SARS-CoV-2 UK variant. *MedRxiv* 2021. doi: <https://doi.org/10.1101/2021.02.02.21250799> 
11. *Collier DA, DeMarco A, Ferreira I, et al. SARS-CoV-2 B.1.1.7 sensitivity to mRNA

- vaccine-elicited, convalescent and monoclonal antibodies. MedRxiv 2021.
doi: <https://doi.org/10.1101/2021.01.19.21249840> 
12. *Wu K, Werner AP, Moliva JI, et al. mRNA-1273 vaccine induces neutralizing antibodies against spike mutants from global SARS-CoV-2 variants. BioRxiv 2021.
doi: <https://doi.org/10.1101/2021.01.25.427948> 
 13. Emary KRW, Golubchik T, Aley PK, et al. Efficacy of ChAdOx1 nCoV-19 (AZD1222) Vaccine Against SARS-CoV-2 VOC 202012/01 (B.1.1.7). 2021. The Lancet.
doi: <http://dx.doi.org/10.2139/ssrn.3779160> 
 14. [FACT SHEET FOR HEALTH CARE PROVIDERS EMERGENCY USE AUTHORIZATION \(EUA\) OF REGEN-COV \(fda.gov\)](#) 
 15. *Wang P, Wang M, Yu J, et al. Increased Resistance of SARS-CoV-2 Variant P.1 to Antibody Neutralization. BioRxiv 2021.
doi: <https://doi.org/10.1101/2021.03.01.433466> 
 16. Pearson CAB, Russell TW, Davies NG, et al. Estimates of severity and transmissibility of novel South Africa SARS-CoV-2 variant 501Y.V2. Retrieved from: [pdf \(cmmid.github.io\)](#) 
 17. Liu Y, Liu J, Xia H, et al. Neutralizing Activity of BNT162b2-Elicited Serum. 2021. NEJM. DOI: 10.1056/NEJMc2102017
 18. *Madhi SA, Ballie V, Cutland CL, et al. Safety and efficacy of the ChAdOx1 nCoV-19 (AZD1222) Covid-19 vaccine against the B.1.351 variant in South Africa. MedRxiv 2021. doi: <https://doi.org/10.1101/2021.02.10.21251247> 
 19. [Novavax COVID-19 Vaccine Demonstrates 89.3% Efficacy in UK Phase 3 Trial | Novavax Inc. – IR Site](#) 
 20. [Johnson & Johnson COVID-19 Vaccine Authorized by U.S. FDA For Emergency Use | Johnson & Johnson \(jnj.com\)](#) 
 21. *Deng X, Garcia-Knight MA, Khalid MM, et al. Transmission, infectivity, and antibody neutralization of an emerging SARS-CoV-2 variant in California carrying a L452R spike protein mutation. MedRxiv 2021. doi: <https://doi.org/10.1101/2021.03.07.21252647> 
 22. Xie X, Liu Y, Liu J, et al. SARS-CoV-2 spike E484K mutation reduces antibody neutralisation. The Lancet 2021. doi: [https://doi.org/10.1016/S2666-5247\(21\)00068-9](https://doi.org/10.1016/S2666-5247(21)00068-9) 
 23. Garcia-Beltran W, Lam EC, St. Denis K, et al. Multiple SARS-CoV-2 variants escape neutralization by vaccine-induced humoral immunity. Cell 2021. doi: <https://doi.org/10.1016/j.cell.2021.03.013> 
 24. *Annavajhala MK, Mohri H, Zucker JE, et al. A Novel SARS-CoV-2 Variant of Concern, B.1.526, Identified in New York. MedRxiv 2021. DOI: [10.1101/2021.02.23.21252259](https://doi.org/10.1101/2021.02.23.21252259) 
 25. *Yadav PD, Sapkal GN, Abraham P, et al. Neutralization of variant under investigation B.1.617 with sera of BBV152 vaccinees. BioRxiv 2021. DOI: <https://doi.org/10.1101/2021.04.23.441101> 
 26. Greaney AJ, Loes AN, Crawford KHD, et al. Comprehensive mapping of mutations in the SARS-CoV-2 receptor-binding domain that affect recognition by polyclonal human

plasma antibodies. Cell 2021. DOI: <https://doi.org/10.1016/j.chom.2021.02.003> 

27. *Edara VV, Lai L, Sahoo MK, et al. Infection and vaccine-induced neutralizing antibody responses to the SARS-CoV-2 B.1.617.1 variant. BioRxiv 2021. DOI: <https://doi.org/10.1101/2021.05.09.443299> 

*Non-peer-reviewed

Related Resources

[Emerging SARS-CoV-2 Variants](#)

[New Variants of the Virus that Causes COVID-19](#)

[Cases, Data, and Surveillance](#)

[COVID-19 Genomic Epidemiology Toolkit](#)

Last Updated June 4, 2021

Content source: [National Center for Immunization and Respiratory Diseases \(NCIRD\), Division of Viral Diseases](#)