



COVID-19

Preparing for Your COVID-19 Vaccination

Updated Aug. 19, 2021

[Print](#)

NOTICE: HHS [announced a plan](#) to begin offering COVID-19 vaccine booster shots this fall. CDC's independent advisory committee, the Advisory Committee on Immunization Practices, will continue to meet and discuss data on the evolution of the pandemic and the use of COVID-19 vaccines. ACIP will make further recommendations on the use of boosters for the public after a thorough review of the evidence.

[COVID-19 vaccines](#) are effective at protecting you from getting sick even after you have had COVID-19. Vaccination is an important tool to help us get back to normal. This information will help you prepare for your COVID-19 vaccination.

Learn more about the different types of COVID-19 vaccines and [how they work](#).

- [Pfizer-BioNTech COVID-19 Vaccine](#)
- [Moderna COVID-19 Vaccine](#)
- [Johnson & Johnson's Janssen COVID-19 Vaccine](#)

Learn more about the [benefits of getting a COVID-19 vaccination](#).



Find a COVID-19 Vaccine: Search [vaccines.gov](#), text your ZIP code to 438829, or call 1-800-232-0233 to find locations near you.

Plan and Prepare for Your COVID-19 Vaccination

- Find out [how to get a COVID-19 vaccine](#)
- Get vaccinated regardless of whether you already had COVID-19.
- Get a COVID-19 vaccine or other vaccines at the [same visit or without waiting 14 days](#) between vaccines.
- Learn more about [routine medical procedures and screenings](#) and COVID-19 vaccination.
- If you are getting a COVID-19 vaccine that requires two doses, be sure to [schedule an appointment for your second shot](#).
- People with moderately to severely compromised immune systems should [receive an additional dose](#) of mRNA COVID-19 vaccine after the initial 2 doses.


 **Watch Video:** [What to Expect at Your COVID-19 Vaccination Appointment \[00:00:48\]](#)

Get Vaccinated Even If You Have Had COVID-19

Already had **COVID-19**?

Studies show that getting a **COVID-19 vaccine** after you recover from COVID-19 provides added protection to your immune system.

Vaccines are a safe way to keep you from getting and spreading COVID-19. COVID-19 vaccination also helps **protect you** from serious illness if you get sick again.



 [JPG - 2 MB]

Get vaccinated to protect against serious illness.

You should get a COVID-19 vaccine, even if you have already had COVID-19 because:

- Research has not yet shown how long you are protected from getting COVID-19 again after you recover from COVID-19.
- Vaccination helps protect you even if you've already had COVID-19.

Evidence is emerging that people **get better protection by being fully vaccinated** compared with having had COVID-19. [One study](#) showed that unvaccinated people who already had COVID-19 are more than 2 times as likely than fully vaccinated people to get COVID-19 again. Learn more about [why getting vaccinated is a safer way to build protection](#) than getting infected.

If you were treated for COVID-19 with monoclonal antibodies or convalescent plasma, you should wait 90 days before getting a COVID-19 vaccine. Talk to your healthcare professional if you are unsure what treatments you received or if you have more questions about getting a COVID-19 vaccine.

If you or your child have a history of multisystem inflammatory syndrome in adults or children ([MIS-A](#) or [MIS-C](#)), consider delaying vaccination until you have recovered from being sick and for 90 days after the date of diagnosis of MIS-A or MIS-C. Learn more about the [clinical considerations](#) people with a history of MIS-A or MIS-C.

Experts are still learning more about how long vaccines protect against COVID-19 in real-world conditions. CDC will keep the public informed as new evidence becomes available.

Considerations for Taking Medication before Getting Vaccinated

For most people, it is not recommended to avoid, discontinue, or delay medications for underlying medical conditions around the time of COVID-19 vaccination. However, your healthcare provider should talk to you about what is currently known and not known about the effectiveness of getting a COVID-19 vaccine when taking medications that suppress the immune system.

It is not recommended you take over-the-counter medicine – such as ibuprofen, aspirin, or acetaminophen – before vaccination for the purpose of trying to prevent vaccine-related side effects. It is not known how these medications might affect how well the vaccine works. However, if you take these medications regularly for other reasons, you should keep taking them before you get vaccinated. It is also not recommended to take antihistamines before getting a COVID-19 vaccine to try to prevent allergic reactions.

If you have questions about medications that you are taking, talk to your healthcare professional or your vaccination provider.

Learn more about medications to relieve [post-vaccination side effects](#).

Last Updated Aug. 19, 2021