

**SUPPLEMENTARY TABLE 1. Clinical severity of COVID-19 sequelae at 1 month after positive test, 2 months after positive test, and date of interview by disease severity at diagnosis (N = 365)\* — Long Beach, California, April 1-December 10, 2020**

	Severity of symptoms, no. of persons, relative to date of positive test									Adjusted % of symptomatic persons, relative to date of positive test		
	1 month (n = 364)			2 months (n = 363)			Interview date (n = 365)			1 month (n = 174)	2 months (n = 127)	Interview date (n = 114)
Severity at Diagnosis	Asym	Mild	Mod	Asym	Mild	Mod	Asym	Mild	Mod	N/A	N/A	N/A
<b>Asym (n = 27)</b>	27	0	0	26	1	0	25	2	0	0.0	2.6	5.4
<b>Mild (n = 222<sup>†</sup>)</b>	131	85	5	156	61	4	167	50	5	22.3	20.8	18.0
<b>Moderate (n = 97)</b>	26	19	52	46	13	38	51	17	29	40.1	37.2	34.5
<b>Severe/critical (n = 19<sup>§</sup>)</b>	6	5	8	8	5	5	8	7	4	37.5	39.3	42.1

Abbreviations: Asym = asymptomatic; mod = moderate; N/A = not applicable.

\* Excludes 1 person for whom disease severity at diagnosis was undetermined.

<sup>†</sup> At months 1 and 2, data were insufficient for 1 person (n = 221).