

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Trends in Agricultural Triazole Fungicide Use in the United States, 1992–2016 and Possible Implications for Antifungal-Resistant Fungi in Human Disease

Mitsuru Toda, Karlyn D. Beer, Kathryn M. Kuivila, Tom M. Chiller, and Brendan R. Jackson

Table of Contents

Table S1. Low, high, and average estimates of agricultural triazole use by year in metric tons, United States, 1992–2016.

Table S2. Average agricultural triazole fungicide use by year and compound type in metric tons, United States, 1992–2016.

Table S3. Average agricultural triazole use by year and crop type in metric tons, United States, 1992–2016.

Table S4. Triazole use by state over five time periods in metric tons, United States, 1992–2016.

Table S5. Average agricultural triazole fungicide use by state over five time periods in metric tons, United States, 1992–2016.

Table S6. Triazole fungicide use in California, 2017.

Figure S1. Agricultural triazole fungicide usage by state and crop type in metric tons, United States, 1992–2016. A) Wheat, B) Corn, C) Soybeans, D) Other crops (cropland for pasture, fallow and idle cropland, pastureland, other hay, alfalfa, sorghum, non-wheat grains, tobacco, peanuts, sugarcane, sugar beets, and other miscellaneous crops) E) Orchards and grapes (stone fruit trees, citrus, nut trees, apples, pears, and grapevines), F) Rice, G) Vegetables and fruit (all vegetables and non-orchard fruit, including beans, peas, greens, berries, and melons), H) Cotton.