

SUPPLEMENTARY TABLE 1. Annual average estimated number† and percentage of adults aged ≥18 years who had suicidal thoughts during the previous year, by state and sex — National Survey on Drug Use and Health, United States, 2015-2019

	Male			Female		
	No.	%	(95% CI)	No.	%	(95% CI)
US total	4,860	4.1	4.0–4.3	5,727	4.5	4.4–4.6
State						
Alabama	56	3.3	2.5–4.2	89	4.6	3.7–5.6
Alaska	17	6.3	5.3–7.6	15	5.9	5.0–7.0
Arizona	111	4.3	3.3–5.7	127	4.7	3.8–5.8
Arkansas	46	4.3	3.3–5.6	60	5.2	4.1–6.5
California	669	4.6	4.1–5.2	590	3.9	3.5–4.4
Colorado	114	5.4	4.2–7.0	146	6.8	5.3–8.6
Connecticut	49	3.7	2.7–5.2	51	3.5	2.9–4.3
Delaware	18	5.2	4.0–6.7	21	5.4	4.0–7.4
District of Columbia	9	3.4	2.5–4.7	12	4.0	3.1–5.2
Florida	260	3.3	2.8–3.9	328	3.9	3.3–4.5
Georgia	161	4.5	3.5–5.7	135	3.3	2.7–4.1
Hawaii	24	4.7	3.5–6.3	22	4.0	3.0–5.5
Idaho	27	4.4	3.5–5.6	43	6.7	5.7–8.0
Illinois	153	3.3	2.8–3.9	198	4.0	3.4–4.6
Indiana	116	4.8	3.8–6.1	159	6.2	5.2–7.3
Iowa	50	4.3	3.3–5.5	61	5.1	4.0–6.5
Kansas	51	4.9	3.7–6.3	62	5.6	4.6–6.9
Kentucky	96	6.0	4.8–7.5	74	4.3	3.5–5.2
Louisiana	60	3.7	2.9–4.8	92	5.1	4.1–6.3
Maine	22	4.2	3.2–5.5	26	4.8	3.8–6.1
Maryland	81	3.7	2.9–4.8	106	4.4	3.5–5.4
Massachusetts	110	4.3	3.2–5.7	131	4.7	3.6–6.1
Michigan	133	3.6	3.0–4.3	179	4.5	3.9–5.3
Minnesota	90	4.4	3.4–5.7	97	4.6	3.7–5.7

Mississippi	40	3.9	2.7–5.5	58	5.0	3.9–6.3
Missouri	82	3.7	2.9–4.8	107	4.5	3.4–5.9
Montana	20	5.0	4.0–6.1	23	5.7	4.7–7.0
Nebraska	25	3.7	2.9–4.7	37	5.1	4.1–6.5
Nevada	41	3.7	2.6–5.0	76	6.6	5.4–8.0
New Hampshire	24	4.6	3.6–5.9	31	5.8	4.8–7.0
New Jersey	80	2.5	1.9–3.2	141	4.0	3.1–5.1
New Mexico	34	4.5	3.2–6.2	39	4.9	3.8–6.1
New York	249	3.5	2.9–4.1	316	4.0	3.4–4.6
North Carolina	155	4.3	3.4–5.3	173	4.2	3.5–5.1
North Dakota	11	3.9	2.9–5.2	14	5.2	4.3–6.4
Ohio	222	5.2	4.5–6.1	257	5.6	4.9–6.3
Oklahoma	51	3.6	2.7–4.8	68	4.5	3.6–5.7
Oregon	91	5.8	4.6–7.3	91	5.5	4.5–6.7
Pennsylvania	196	4.1	3.4–4.9	252	4.9	4.2–5.8
Rhode Island	22	5.6	4.5–7.1	18	4.2	3.2–5.4
South Carolina	68	3.8	2.7–5.3	96	4.8	3.7–6.1
South Dakota	12	3.8	2.9–4.9	12	3.9	2.9–5.2
Tennessee	105	4.3	3.4–5.5	131	4.9	4.1–5.9
Texas	323	3.3	2.8–3.8	403	3.9	3.4–4.4
Utah	73	6.8	5.3–8.6	77	7.1	5.6–8.9
Vermont	13	5.2	4.0–6.7	18	6.9	5.7–8.4
Virginia	127	4.2	3.5–5.0	131	3.9	3.3–4.7
Washington	143	5.2	4.2–6.4	157	5.5	4.3–7.0
West Virginia	31	4.5	3.5–5.8	40	5.4	4.5–6.6
Wisconsin	92	4.2	3.3–5.4	120	5.3	4.3–6.6
Wyoming	8	3.8	2.7–5.2	15	7.1	5.7–8.8
Region						
Northeast	765	3.7	3.4–4.0	985	4.3	4.0–4.7
Midwest	1,037	4.2	3.9–4.5	1,302	4.9	4.6–5.2

South	1,686	3.8	3.6–4.1		2,018	4.2	4.0–4.4
West	1,371	4.8	4.5–5.2		1,421	4.8	4.5–5.1

Abbreviations: No. = number, CI = confidence interval.

[†]In thousands.

NOTE: Respondents were asked, "At any time in the past 12 months, did you seriously think about trying to kill yourself?" If they answered "Yes," they were categorized as having serious thoughts of suicide in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2015-2019.

SUPPLEMENTARY TABLE 2. Annual average estimated number[†] and percentage of adults aged ≥18 years who had suicidal thoughts during the previous year, by state and age group — National Survey on Drug Use and Health, United States, 2015-2019

Age group	18-39 years			40-55 years			≥ 56 years		
	No.	%	(95% CI)	No.	%	(95% CI)	No.	%	(95% CI)
US total	6,515	6.9	6.8–7.1	2,218	3.4	3.2–3.6	1,854	2.2	2.0–2.3
State									
Alabama	94	7.1	5.9–8.5	39	4.0	2.8–5.9	13	0.9	0.4–2.3
Alaska	22	10.2	9.0–11.7	5	3.6	2.3–5.6	5	2.8	1.6–4.8
Arizona	151	7.7	6.5–9.0	59	4.4	2.9–6.5	29	1.5	0.7–3.0
Arkansas	59	7.1	5.8–8.6	26	4.7	3.1–6.9	20	2.5	1.4–4.3
California	799	6.6	6.0–7.2	219	2.7	2.2–3.4	242	2.5	2.0–3.2
Colorado	146	8.5	7.2–10.1	57	5.0	3.4–7.2	56	4.1	2.6–6.4
Connecticut	67	6.7	5.8–7.8	17	2.1	1.4–3.3	17	1.7	0.8–3.7
Delaware	21	7.8	6.4–9.5	8	4.4	3.0–6.3	10	3.6	2.4–5.3
District of	15	5.1	4.1–6.4	4	3.1	1.9–5.1	2	1.5	0.8–2.9
Florida	350	6.3	5.7–6.9	105	2.5	1.9–3.3	133	2.0	1.5–2.8
Georgia	169	5.6	4.8–6.4	75	3.4	2.4–4.8	52	2.2	1.3–3.5
Hawaii	29	7.9	6.5–9.5	8	3.2	2.0–5.1	8	1.9	1.0–3.8
Idaho	41	8.5	7.5–9.5	16	5.1	3.2–7.9	13	2.9	1.7–4.7
Illinois	232	6.2	5.5–7.1	69	2.7	2.0–3.6	49	1.5	0.9–2.4
Indiana	183	9.6	8.2–11.2	60	4.6	3.3–6.3	33	1.8	1.2–2.9
Iowa	67	7.5	6.3–8.9	23	4.0	2.6–6.1	21	2.3	1.4–3.9
Kansas	63	7.4	6.3–8.7	26	4.9	3.6–6.5	23	3.1	1.7–5.6
Kentucky	97	7.8	6.6–9.3	40	4.5	3.1–6.6	34	2.8	1.7–4.5
Louisiana	82	6.0	4.9–7.4	39	4.3	2.9–6.4	32	2.7	1.8–4.1
Maine	27	7.8	6.6–9.3	12	4.7	3.1–7.0	9	2.0	1.3–3.1
Maryland	120	6.9	5.6–8.4	29	2.3	1.3–4.0	37	2.3	1.5–3.7
Massachusetts	151	7.3	5.8–9.0	50	3.8	2.4–5.8	40	2.0	1.1–3.8
Michigan	186	6.7	6.0–7.5	84	4.2	3.3–5.4	42	1.5	0.9–2.3
Minnesota	129	7.9	6.9–9.0	34	3.0	1.7–5.4	25	1.7	0.8–3.7
Mississippi	57	6.9	5.7–8.3	21	3.5	2.2–5.6	20	2.6	1.5–4.5
Missouri	132	7.7	6.3–9.4	30	2.3	1.4–3.9	27	1.7	0.9–3.1
Montana	25	8.5	7.2–9.9	10	5.1	3.7–6.9	9	2.7	1.8–4.1
Nebraska	37	6.6	5.4–7.9	14	3.9	2.6–5.8	12	2.4	1.4–3.9
Nevada	66	7.5	6.3–8.9	32	5.6	3.9–8.1	18	2.2	1.3–3.7
New Hampshire	31	8.7	7.5–10.1	11	3.8	2.5–5.6	14	3.2	2.0–5.3
New Jersey	143	5.8	5.0–6.7	45	2.2	1.5–3.3	34	1.4	0.7–3.1
New Mexico	42	7.4	6.3–8.7	13	3.3	2.2–5.0	17	2.9	1.6–5.4
New York	363	6.2	5.6–6.8	92	2.3	1.7–3.2	110	2.1	1.4–3.0
North Carolina	183	6.4	5.5–7.4	76	3.5	2.5–5.0	70	2.6	1.7–3.9
North Dakota	18	7.7	6.5–9.1	3	1.9	1.2–3.2	5	2.4	1.4–4.1

Ohio	269	8.3	7.6–9.1		97	4.3	3.5–5.3		114	3.4	2.5–4.5
Oklahoma	70	6.3	5.3–7.5		34	4.4	2.9–6.5		14	1.4	0.7–2.6
Oregon	110	9.3	8.1–10.6		42	4.8	3.5–6.5		31	2.7	1.6–4.3
Pennsylvania	248	7.2	6.6–7.8		110	4.2	3.1–5.6		90	2.4	1.6–3.5
Rhode Island	27	8.5	7.3–9.8		6	3.1	2.1–4.5		7	2.4	1.3–4.4
South Carolina	83	6.1	5.0–7.5		39	3.8	2.5–5.7		43	3.0	1.6–5.5
South Dakota	17	6.8	5.5–8.3		5	3.0	1.8–4.9		3	1.3	0.7–2.5
Tennessee	141	7.5	6.3–9.0		64	4.7	3.3–6.6		31	1.7	0.9–3.2
Texas	528	6.2	5.5–6.9		110	2.0	1.5–2.6		88	1.4	1.0–2.0
Utah	105	10.3	8.6–12.2		36	6.6	4.5–9.6		8	1.4	0.7–2.9
Vermont	16	8.9	7.4–10.7		8	6.3	4.6–8.6		7	3.5	2.2–5.4
Virginia	165	6.9	6.0–7.9		55	3.1	2.2–4.3		38	1.7	0.9–3.1
Washington	166	7.5	6.5–8.6		76	5.1	3.7–6.9		58	3.1	1.8–5.2
West Virginia	37	7.9	6.9–9.1		19	5.1	3.7–7.1		14	2.4	1.3–4.5
Wisconsin	123	7.6	6.5–8.9		64	5.5	3.8–7.8		25	1.5	0.8–2.8
Wyoming	14	8.4	7.0–10.0		4	4.2	2.7–6.4		5	3.2	1.7–6.1
Region											
Northeast	1,073	6.7	6.3–7.0		350	3.0	2.6–3.5		328	2.1	1.7–2.5
Midwest	1,453	7.5	7.2–7.8		509	3.8	3.4–4.2		378	2.0	1.7–2.4
South	2,272	6.5	6.2–6.8		782	3.1	2.8–3.5		650	2.0	1.8–2.3
West	1,717	7.4	7.0–7.8		577	3.8	3.4–4.2		499	2.6	2.2–3.0

Abbreviations: No. = number, CI = confidence interval.

[†]In thousands.

NOTE: Respondents were asked, "At any time in the past 12 months, did you seriously think about trying to kill yourself?" If they answered "Yes," they were categorized as having serious thoughts of suicide in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2015-2019.

SUPPLEMENTARY TABLE 3. Annual average estimated number[†] and percentage of adults aged ≥18 years who had suicidal thoughts during the previous year, by state and race/ethnicity — National Survey on Drug Use and Health, United States, 2015-2019

Race/ethnicity	NH white			NH black			NH Asian			NH AI/AN			Hispanic		
	No.	%	(95% CI)	No.	%	(95% CI)	No.	%	(95% CI)	No.	%	(95% CI)	No.	%	(95% CI)
US total	7,169	4.6	4.4–4.7	1,081	3.7	3.5–4.0	400	2.9	2.5–3.4	68	5.0	4.2–6.0	1,514	3.9	3.6–4.1
State															
Alabama	106	4.2	3.3–5.4	27	2.9	2.1–4.1	*	*	*	*	*	*	7	5.5	2.2–13.3
Alaska	18	5.3	4.5–6.3	*	*	*	2	5.0	3.0–8.3	5	6.7	4.5–9.7	3	10.2	6.9–15.0
Arizona	136	4.3	3.5–5.3	5	2.2	1.0–4.9	5	2.9	1.2–6.8	3	1.6	0.7–4.0	79	5.5	4.2–7.2
Arkansas	82	4.9	3.9–6.0	15	4.6	3.0–6.9	*	*	*	*	*	*	6	4.3	2.1–8.7
California	605	5.0	4.5–5.6	72	4.4	3.2–6.1	147	3.2	2.4–4.3	*	*	*	384	3.7	3.2–4.3
Colorado	195	6.4	5.2–7.8	14	9.0	4.1–18.4	*	*	*	*	*	*	34	4.3	2.8–6.5
Connecticut	63	3.2	2.5–4.0	9	3.5	2.2–5.6	2	1.6	0.5–4.6	*	*	*	22	5.7	3.7–8.8
Delaware	30	6.1	5.0–7.5	6	3.6	2.5–5.4	1	1.9	0.5–6.8	*	*	*	2	4.0	2.3–7.1
District of	8	3.5	2.4–5.0	9	3.9	2.8–5.2	1	5.0	2.3–10.6	*	*	*	2	3.2	1.7–6.2
Florida	359	3.8	3.3–4.5	83	3.6	2.7–4.7	9	2.3	1.1–4.6	*	*	*	120	3.0	2.3–4.0
Georgia	191	4.5	3.6–5.5	78	3.3	2.7–4.2	8	3.0	1.3–6.7	*	*	*	12	1.9	1.1–3.5
Hawaii	10	4.0	2.9–5.5	*	*	*	10	2.5	1.7–3.8	*	*	*	6	7.1	4.9–10.2
Idaho	57	5.3	4.5–6.3	*	*	*	*	*	*	*	*	*	8	6.1	3.8–9.7
Illinois	238	3.8	3.3–4.5	49	3.8	2.5–5.7	17	3.2	1.7–5.7	*	*	*	42	2.9	2.1–4.0
Indiana	224	5.5	4.6–6.5	22	5.1	3.0–8.4	*	*	*	*	*	*	18	6.5	3.5–11.7
Iowa	98	4.7	3.8–5.8	3	4.4	2.0–9.5	*	*	*	*	*	*	3	3.1	1.7–5.7
Kansas	88	5.2	4.2–6.4	5	4.6	1.7–11.7	1	2.0	0.7–5.6	*	*	*	11	5.4	3.3–8.8
Kentucky	156	5.4	4.5–6.4	11	4.3	2.0–9.0	*	*	*	*	*	*	2	1.6	0.6–4.6
Louisiana	104	4.9	3.9–6.1	35	3.4	2.4–4.8	*	*	*	*	*	*	5	2.9	1.1–7.2
Maine	45	4.5	3.8–5.2	*	*	*	*	*	*	*	*	*	*	*	*
Maryland	109	4.4	3.5–5.6	55	4.1	2.8–6.1	9	2.9	1.4–5.9	*	*	*	10	2.5	1.5–4.1
Massachusetts	191	4.7	3.7–6.1	15	4.1	2.0–8.0	6	1.6	0.6–3.8	*	*	*	22	4.1	2.6–6.7
Michigan	246	4.1	3.6–4.7	30	3.1	2.3–4.2	5	2.2	1.2–4.0	*	*	*	14	4.3	2.6–7.0
Minnesota	164	4.7	3.8–5.7	6	2.9	1.6–5.1	8	4.4	1.7–11.1	*	*	*	3	1.6	0.7–3.8
Mississippi	63	4.8	3.8–6.2	30	3.8	2.8–5.0	*	*	*	*	*	*	*	*	*
Missouri	152	4.1	3.3–5.0	21	4.3	2.7–6.6	*	*	*	*	*	*	3	2.0	0.8–5.1
Montana	35	5.0	4.3–5.7	*	*	*	*	*	*	4	10.6	6.7–16.4	2	7.9	3.5–17.2
Nebraska	50	4.3	3.5–5.2	6	10.2	5.7–17.4	*	*	*	*	*	*	3	2.7	1.6–4.7
Nevada	65	5.4	4.2–7.0	9	4.7	2.8–7.8	10	5.8	3.2–10.4	*	*	*	27	4.7	3.5–6.3
New	49	5.0	4.3–5.9	*	*	*	*	*	*	*	*	*	3	7.8	4.2–13.9
New Jersey	136	3.4	2.8–4.3	41	4.8	2.5–9.0	9	1.4	0.8–2.3	*	*	*	30	2.3	1.6–3.4
New Mexico	37	5.8	4.2–8.0	*	*	*	*	*	*	3	2.7	1.5–4.9	28	4.0	3.1–5.2
New York	338	3.9	3.3–4.4	68	3.2	2.5–4.1	19	1.4	0.9–2.2	*	*	*	130	4.8	3.7–6.4
North Carolina	238	4.6	3.9–5.5	50	3.1	2.2–4.4	5	2.3	0.9–5.3	1	1.4	0.7–2.8	26	4.5	3.0–6.7
North Dakota	21	4.2	3.5–5.1	*	*	*	*	*	*	1	5.1	3.1–8.4	*	*	*
Ohio	385	5.3	4.8–5.9	53	5.2	3.8–7.1	8	4.1	2.4–6.7	*	*	*	17	6.2	4.1–9.3

Oklahoma	82	4.1	3.2–5.2	8	3.9	2.0–7.3	*	*	*	11	4.6	2.7–7.9	7	3.0	1.6–5.6
Oregon	145	5.7	4.8–6.8	*	*	*	4	2.9	1.5–5.4	*	*	*	20	5.8	3.8–8.6
Pennsylvania	350	4.5	3.9–5.1	32	3.4	2.5–4.5	6	1.9	1.0–3.6	*	*	*	44	7.4	4.6–11.6
Rhode Island	28	4.4	3.5–5.5	*	*	*	1	3.1	1.1–8.9	*	*	*	8	7.6	5.4–10.7
South Carolina	119	4.7	3.6–6.1	32	3.3	2.4–4.4	*	*	*	*	*	*	4	2.6	1.3–5.2
South Dakota	20	3.6	2.9–4.6	*	*	*	*	*	*	3	6.5	4.2–9.9	0	1.5	0.8–2.7
Tennessee	183	4.7	3.9–5.6	28	3.5	2.6–4.7	*	*	*	*	*	*	20	9.2	4.8–17.0
Texas	363	3.9	3.4–4.6	69	2.9	2.3–3.7	32	3.2	1.8–5.5	*	*	*	234	3.2	2.8–3.7
Utah	113	6.5	5.4–7.8	*	*	*	*	*	*	*	*	*	19	7.2	4.7–10.9
Vermont	29	6.3	5.3–7.4	*	*	*	*	*	*	*	*	*	*	*	*
Virginia	172	4.2	3.5–5.0	47	4.0	2.7–5.7	12	3.1	1.8–5.4	*	*	*	14	2.6	1.5–4.4
Washington	232	5.7	4.7–6.9	7	3.4	1.3–8.5	17	3.5	1.3–8.8	*	*	*	24	4.2	2.5–6.9
West Virginia	64	4.8	4.1–5.7	1	2.7	1.3–5.5	*	*	*	*	*	*	*	*	*
Wisconsin	162	4.3	3.6–5.2	16	6.7	3.4–12.7	5	4.4	2.0–9.5	*	*	*	22	8.9	4.7–16.2
Wyoming	19	5.1	4.2–6.2	*	*	*	*	*	*	*	*	*	3	7.9	4.0–15.0
Region															
Northeast	1,228	4.1	3.8–4.5	169	3.6	2.9–4.5	44	1.6	1.2–2.0	2	3.8	1.9–7.4	261	4.6	3.9–5.5
Midwest	1,846	4.6	4.3–4.8	212	4.3	3.7–5.0	62	3.8	2.9–4.9	18	7.7	5.7–10.3	140	4.1	3.4–4.9
South	2,428	4.4	4.1–4.6	583	3.5	3.2–3.8	93	3.0	2.3–3.9	21	4.5	3.0–6.7	477	3.2	2.9–3.6
West	1,667	5.3	5.0–5.7	118	4.7	3.7–5.9	201	3.3	2.6–4.1	27	4.5	3.4–5.9	637	4.1	3.7–4.6

Abbreviations: No. = number, CI = confidence interval, NH = Non-Hispanic, AI/AN = American Indian or Alaska Native.

†In thousands.

NOTE: Estimated numbers appearing as 0 in this table mean that the estimate is greater than 0 but less than 500 (because estimated numbers are shown in thousands).

NOTE: Respondents were asked, "At any time in the past 12 months, did you seriously think about trying to kill yourself?" If they answered "Yes," they were categorized as having serious thoughts of suicide in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2015-2019.

*Low precision; no estimate reported.

SUPPLEMENTARY Table 4. Annual average estimated number[†] and percentage of adults aged ≥18 years who made a suicide plan during the previous year, by state and sex — National Survey on Drug Use and Health, United States, 2015-2019

	Male			Female		
	No.	%	(95% CI)	No.	%	(95% CI)
US total	1,356	1.1	1.1–1.2	1,728	1.4	1.3–1.4
Alabama	16	0.9	0.5–1.6	25	1.3	0.9–1.8
Alaska	7	2.6	1.9–3.4	6	2.2	1.6–3.0
Arizona	29	1.1	0.7–1.7	52	1.9	1.4–2.6
Arkansas	13	1.2	0.7–2.2	18	1.5	1.1–2.2
California	191	1.3	1.1–1.6	167	1.1	0.9–1.3
Colorado	26	1.3	0.8–2.1	45	2.1	1.5–3.0
Connecticut	10	0.7	0.4–1.2	12	0.8	0.4–1.5
Delaware	5	1.6	1.0–2.4	5	1.2	0.6–2.3
District of Columbia	2	0.7	0.3–1.4	4	1.3	0.8–2.1
Florida	65	0.8	0.6–1.1	88	1.0	0.8–1.3
Georgia	45	1.2	0.8–1.9	48	1.2	0.8–1.7
Hawaii	6	1.2	0.7–1.9	7	1.3	0.8–2.2
Idaho	8	1.3	0.9–2.0	17	2.6	2.1–3.3
Illinois	46	1.0	0.7–1.3	69	1.4	1.0–1.9
Indiana	41	1.7	1.2–2.4	58	2.2	1.6–3.1
Iowa	18	1.6	1.1–2.2	24	2.0	1.4–2.9
Kansas	15	1.4	0.9–2.3	20	1.9	1.3–2.7
Kentucky	31	1.9	1.2–3.0	25	1.4	1.0–2.0
Louisiana	20	1.2	0.7–2.2	34	1.9	1.3–2.7
Maine	6	1.2	0.7–2.0	7	1.3	0.8–1.9
Maryland	18	0.8	0.5–1.3	29	1.2	0.8–1.8
Massachusetts	26	1.0	0.7–1.5	29	1.0	0.6–1.6
Michigan	42	1.1	0.9–1.5	45	1.1	0.9–1.5
Minnesota	24	1.1	0.6–2.2	32	1.5	1.0–2.2
Mississippi	16	1.6	0.9–2.7	18	1.5	1.1–2.1
Missouri	25	1.1	0.7–1.7	32	1.3	0.8–2.2
Montana	7	1.7	1.1–2.6	6	1.5	1.0–2.1
Nebraska	10	1.5	0.9–2.4	12	1.7	1.3–2.4
Nevada	15	1.3	0.8–2.3	28	2.4	1.6–3.6
New Hampshire	7	1.3	0.8–2.1	10	1.9	1.2–3.0
New Jersey	15	0.4	0.3–0.7	39	1.1	0.7–1.6
New Mexico	6	0.8	0.5–1.3	11	1.3	0.9–1.9
New York	58	0.8	0.6–1.1	92	1.2	0.9–1.6
North Carolina	43	1.2	0.8–1.8	58	1.4	1.0–1.9
North Dakota	4	1.3	0.8–2.3	4	1.6	1.1–2.2

Ohio	64	1.5	1.1–2.0		74	1.6	1.3–2.1
Oklahoma	13	1.0	0.6–1.6		25	1.7	1.1–2.4
Oregon	20	1.3	0.9–1.9		30	1.8	1.3–2.6
Pennsylvania	42	0.9	0.6–1.3		70	1.4	1.0–1.8
Rhode Island	8	2.0	1.3–3.2		4	0.9	0.6–1.4
South Carolina	22	1.2	0.6–2.4		34	1.7	1.0–2.7
South Dakota	5	1.5	1.0–2.4		3	0.9	0.6–1.6
Tennessee	33	1.4	0.9–2.2		35	1.3	0.9–1.9
Texas	89	0.9	0.7–1.2		107	1.0	0.8–1.3
Utah	24	2.3	1.5–3.4		20	1.8	1.2–2.8
Vermont	4	1.5	1.0–2.4		5	1.8	1.1–2.7
Virginia	31	1.0	0.7–1.5		41	1.2	0.9–1.7
Washington	33	1.2	0.7–2.1		52	1.8	1.3–2.6
West Virginia	13	1.9	1.2–3.0		14	1.9	1.3–2.7
Wisconsin	39	1.8	1.2–2.6		36	1.6	1.1–2.3
Wyoming	2	0.9	0.6–1.5		4	1.8	1.3–2.6
Region							
Northeast	175	0.8	0.7–1.0		267	1.2	1.0–1.4
Midwest	331	1.3	1.2–1.5		410	1.5	1.4–1.7
South	477	1.1	1.0–1.2		607	1.3	1.2–1.4
West	374	1.3	1.1–1.5		444	1.5	1.3–1.7

Abbreviations: No. = number, CI = confidence interval.

†In thousands.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2015-2019.

NOTE: Only respondents who reported suicide ideation were asked about suicide plans.

SUPPLEMENTARY Table 5. Annual average estimated number[†] and percentage of adults aged ≥18 years who made a suicide plan during the previous year, by state and age group — National Survey on Drug Use and Health, United States, 2015-2019

Age group	18-39 years			40-55 years			≥ 56 years		
	No.	%	(95% CI)	No.	%	(95% CI)	No.	%	(95% CI)
US total	2,004	2.1	2.1–2.2	655	1.0	0.9–1.1	425	0.5	0.4–0.6
State									
Alabama	28	2.1	1.5–2.9	11	1.2	0.6–2.5	2	0.2	0.0–1.1
Alaska	9	4.3	3.5–5.3	1	1.1	0.5–2.0	2	0.9	0.4–2.4
Arizona	53	2.7	2.1–3.4	19	1.4	0.8–2.7	8	0.4	0.1–1.3
Arkansas	16	1.9	1.4–2.6	10	1.7	0.9–3.2	5	0.7	0.2–1.8
California	240	2.0	1.7–2.3	60	0.8	0.5–1.2	58	0.6	0.4–1.0
Colorado	42	2.4	1.8–3.4	23	2.0	1.1–3.5	7	0.5	0.2–1.7
Connecticut	12	1.2	0.8–1.7	6	0.7	0.3–1.6	4	0.4	0.1–1.4
Delaware	5	1.9	1.2–3.0	4	2.2	1.2–4.2	1	0.3	0.1–1.1
District of	4	1.3	0.9–2.0	1	1.1	0.4–2.8	1	0.4	0.2–1.1
Florida	96	1.7	1.4–2.0	35	0.8	0.5–1.4	22	0.3	0.2–0.7
Georgia	55	1.8	1.4–2.3	20	0.9	0.5–1.8	18	0.7	0.3–1.7
Hawaii	10	2.7	2.0–3.8	1	0.5	0.2–1.2	2	0.4	0.1–1.4
Idaho	16	3.4	2.6–4.4	6	1.9	1.1–3.5	2	0.5	0.2–1.3
Illinois	73	2.0	1.6–2.4	22	0.8	0.5–1.4	21	0.6	0.2–1.6
Indiana	73	3.8	2.9–5.0	18	1.4	0.7–2.6	8	0.4	0.1–1.9
Iowa	28	3.1	2.4–4.0	9	1.5	0.9–2.7	5	0.6	0.2–1.9
Kansas	20	2.3	1.8–3.1	8	1.5	0.9–2.5	7	1.0	0.4–2.2
Kentucky	39	3.2	2.3–4.3	9	1.0	0.5–2.3	7	0.6	0.2–1.5
Louisiana	27	2.0	1.4–2.8	21	2.4	1.4–3.8	6	0.5	0.2–1.7
Maine	9	2.7	2.0–3.7	3	1.3	0.5–3.2	1	0.1	0.0–0.9
Maryland	31	1.8	1.3–2.5	11	0.9	0.4–2.1	4	0.3	0.1–1.1
Massachusetts	45	2.2	1.5–3.1	7	0.5	0.2–1.2	3	0.2	0.0–1.1
Michigan	55	2.0	1.6–2.4	27	1.4	0.9–2.1	4	0.1	0.0–0.4
Minnesota	32	2.0	1.4–2.8	16	1.4	0.6–3.2	7	0.5	0.2–1.5
Mississippi	20	2.5	1.8–3.4	6	1.0	0.5–2.3	8	1.0	0.4–2.6
Missouri	47	2.7	2.0–3.8	9	0.7	0.3–1.8	2	0.1	0.0–0.8
Montana	6	2.1	1.4–2.9	3	1.5	0.8–2.8	4	1.2	0.6–2.2
Nebraska	12	2.2	1.7–2.8	5	1.3	0.6–2.6	6	1.1	0.5–2.8
Nevada	25	2.8	2.1–3.8	9	1.5	0.7–3.3	9	1.1	0.4–3.0
New	8	2.3	1.8–2.9	2	0.7	0.3–1.6	7	1.6	0.8–3.2
New Jersey	41	1.7	1.3–2.2	7	0.4	0.2–0.9	5	0.2	0.0–1.4
New Mexico	12	2.1	1.6–2.6	3	0.8	0.4–1.6	2	0.3	0.1–1.2
New York	95	1.6	1.3–1.9	26	0.7	0.4–1.1	29	0.5	0.2–1.2
North Carolina	55	1.9	1.4–2.6	19	0.9	0.5–1.5	28	1.0	0.5–2.0
North Dakota	5	2.2	1.6–3.0	1	0.6	0.3–1.2	2	1.1	0.5–2.6
Ohio	85	2.6	2.2–3.1	27	1.2	0.7–1.9	26	0.8	0.4–1.4

Oklahoma	26	2.4	1.8–3.1	10	1.3	0.7–2.3	2	0.2	0.0–1.2
Oregon	34	2.8	2.2–3.7	8	0.9	0.5–1.7	9	0.8	0.3–2.0
Pennsylvania	72	2.1	1.8–2.4	24	0.9	0.4–1.9	15	0.4	0.2–0.9
Rhode Island	7	2.3	1.7–3.1	1	0.7	0.3–1.5	3	1.1	0.4–3.0
South Carolina	29	2.2	1.4–3.4	15	1.5	0.8–2.7	11	0.8	0.3–2.3
South Dakota	5	2.1	1.5–2.9	2	1.1	0.5–2.6	1	0.4	0.1–1.6
Tennessee	47	2.5	1.7–3.5	20	1.5	0.7–3.2	2	0.1	0.0–0.7
Texas	154	1.8	1.5–2.2	28	0.5	0.3–0.8	14	0.2	0.1–0.4
Utah	33	3.2	2.5–4.0	11	1.9	0.9–4.0	1	0.2	0.0–1.1
Vermont	5	2.7	1.9–3.8	3	2.1	1.1–3.9	1	0.5	0.2–1.5
Virginia	50	2.1	1.6–2.7	17	1.0	0.5–1.7	5	0.2	0.1–0.7
Washington	51	2.3	1.8–3.0	20	1.3	0.6–2.9	14	0.7	0.3–1.9
West Virginia	13	2.8	2.2–3.5	6	1.7	1.1–2.8	7	1.3	0.5–3.1
Wisconsin	43	2.6	2.0–3.4	23	2.0	1.1–3.6	9	0.5	0.2–1.5
Wyoming	5	2.9	2.2–3.8	1	0.6	0.2–1.8	1	0.3	0.1–1.3
Region									
Northeast	295	1.8	1.7–2.0	79	0.7	0.5–0.9	68	0.4	0.3–0.7
Midwest	478	2.5	2.3–2.7	166	1.2	1.0–1.5	97	0.5	0.4–0.7
South	695	2.0	1.8–2.1	246	1.0	0.8–1.2	143	0.4	0.3–0.6
West	536	2.3	2.1–2.5	165	1.1	0.9–1.3	117	0.6	0.4–0.8

Abbreviations: No. = number, CI = confidence interval.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2015-2019.

*Low precision; no estimate reported.

NOTE: Only respondents who reported suicide ideation were asked about suicide plans.

SUPPLEMENTARY Table 6. Annual average estimated number[†] and percentage of adults aged ≥18 years who made a suicide plan during the previous year, by state and race/ethnicity — National Survey on Drug Use and Health, United States, 2015–2019

Race/Ethnicity	NH white			NH black			NH Asian			NH AI/AN			Hispanic		
	No.	%	(95% CI)	No.	%	(95% CI)	No.	%	(95% CI)	No.	%	(95% CI)	No.	%	(95% CI)
US total	2,014	1.3	1.2–1.3	349	1.2	1.1–1.4	121	.9	0.7–1.1	26	1.9	1.4–2.7	446	1.1	1.0–1.3
State															
Alabama	33	1.3	0.9–1.9	8	0.8	0.5–1.5	*	*	*	*	*	*	*	*	*
Alaska	5	1.5	1.1–2.1	0	2.7	1.0–7.1	1	2.6	1.5–4.8	3	4.1	2.3–7.1	2	5.4	3.0–9.6
Arizona	42	1.3	0.9–2.0	2	0.9	0.3–2.7	*	*	*	2	0.8	0.2–2.7	33	2.3	1.5–3.4
Arkansas	24	1.4	1.0–2.1	4	1.2	0.6–2.6	*	*	*	*	*	*	1	1.0	0.4–2.6
California	147	1.2	1.0–1.5	24	1.5	0.8–2.9	48	1.1	0.6–1.9	*	*	*	123	1.2	0.9–1.5
Colorado	60	2.0	1.4–2.6	3	1.9	0.6–6.4	*	*	*	*	*	*	5	0.6	0.3–1.2
Connecticut	14	0.7	0.4–1.2	3	1.1	0.5–2.3	*	*	*	*	*	*	3	0.8	0.3–2.3
Delaware	7	1.4	0.9–2.2	2	1.0	0.5–1.9	*	*	*	*	*	*	1	2.0	0.8–4.8
District of Columbia	1	0.5	0.2–1.0	4	1.5	0.9–2.4	*	*	*	*	*	*	1	1.6	0.5–5.1
Florida	89	0.9	0.7–1.3	33	1.4	0.9–2.2	2	0.6	0.1–2.9	*	*	*	23	0.6	0.4–0.8
Georgia	54	1.3	0.8–1.9	28	1.2	0.8–1.7	*	*	*	*	*	*	7	1.2	0.6–2.3
Hawaii	2	1.0	0.5–2.0	*	*	*	3	0.7	0.4–1.3	*	*	*	1	1.5	0.7–3.4
Idaho	20	1.9	1.5–2.4	*	*	*	*	*	*	*	*	*	3	1.9	1.0–3.7
Illinois	66	1.1	0.8–1.3	24	1.9	0.9–3.9	7	1.3	0.5–3.4	*	*	*	17	1.2	0.7–2.0
Indiana	76	1.9	1.4–2.5	7	1.7	0.7–3.9	*	*	*	*	*	*	8	2.7	1.1–6.6
Iowa	35	1.7	1.2–2.2	2	3.0	1.1–7.9	*	*	*	*	*	*	2	1.8	0.9–3.4
Kansas	26	1.5	1.1–2.2	1	1.1	0.2–5.0	*	*	*	*	*	*	6	2.7	1.3–5.7
Kentucky	52	1.8	1.3–2.4	3	1.0	0.3–3.5	*	*	*	*	*	*	0	0.3	0.0–1.9
Louisiana	40	1.9	1.3–2.7	11	1.0	0.6–1.8	*	*	*	*	*	*	0	0.2	0.1–0.7
Maine	13	1.3	0.9–1.8	*	*	*	*	*	*	*	*	*	*	*	*
Maryland	21	0.8	0.5–1.4	22	1.7	0.9–2.9	0	0.1	0.0–1.0	*	*	*	3	0.8	0.3–1.8
Massachusetts	34	0.8	0.5–1.3	9	2.6	1.0–6.4	3	0.9	0.3–3.2	*	*	*	4	0.7	0.4–1.4
Michigan	69	1.2	0.9–1.5	8	0.8	0.5–1.3	2	0.8	0.3–2.1	*	*	*	2	0.7	0.3–1.7
Minnesota	45	1.3	0.9–2.0	*	*	*	*	*	*	*	*	*	*	*	*
Mississippi	21	1.6	1.1–2.4	12	1.5	0.9–2.4	*	*	*	*	*	*	*	*	*
Missouri	47	1.2	0.9–1.7	5	1.0	0.4–2.3	*	*	*	*	*	*	2	1.2	0.3–4.6
Montana	11	1.5	1.1–2.0	*	*	*	*	*	*	1	2.4	1.1–4.9	1	2.5	0.9–6.8
Nebraska	17	1.5	1.1–2.0	3	4.2	1.9–9.2	*	*	*	*	*	*	1	1.1	0.4–2.8

Nevada	21	1.8	1.1–3.0		5	2.6	1.3–5.1		3	1.7	0.6–4.7		*	*	*		11	1.9	1.2–2.8
New Hampshire	15	1.5	1.0–2.2		*	*	*		*	*	*		*	*	*		1	2.9	1.0–8.5
New Jersey	32	0.8	0.5–1.2		7	0.8	0.4–1.8		5	0.8	0.3–1.8		*	*	*		7	0.6	0.3–1.1
New Mexico	8	1.2	0.7–1.9		*	*	*		*	*	*		1	0.8	0.3–2.2		8	1.1	0.8–1.5
New York	81	0.9	0.7–1.2		15	0.7	0.4–1.1		9	0.7	0.4–1.2		*	*	*		41	1.5	0.9–2.7
North Carolina	76	1.5	1.1–1.9		14	0.9	0.4–1.7		*	*	*		0	1.0	0.2–3.7		7	1.3	0.6–2.7
North Dakota	7	1.5	1.0–2.1		*	*	*		*	*	*		0	1.1	0.3–3.8		0	0.6	0.1–2.9
Ohio	112	1.6	1.3–1.9		14	1.4	0.7–2.6		*	*	*		*	*	*		6	2.2	1.0–4.8
Oklahoma	26	1.3	0.9–1.8		2	0.8	0.3–2.3		*	*	*		4	1.6	0.6–3.9		3	1.0	0.4–2.6
Oregon	39	1.5	1.1–2.2		*	*	*		1	0.7	0.2–2.7		*	*	*		7	2.0	1.1–3.6
Pennsylvania	79	1.0	0.8–1.3		10	1.0	0.6–1.6		2	0.7	0.2–2.3		*	*	*		16	2.6	1.0–6.7
Rhode Island	9	1.5	1.0–2.2		1	2.3	0.8–6.6		*	*	*		*	*	*		2	1.4	0.7–2.8
South Carolina	42	1.7	1.1–2.6		7	0.7	0.4–1.2		*	*	*		*	*	*		1	0.3	0.1–1.4
South Dakota	6	1.1	0.8–1.5		*	*	*		*	*	*		1	3.3	1.6–6.8		0	1.0	0.7–1.5
Tennessee	53	1.4	0.9–2.0		12	1.5	0.9–2.4		*	*	*		*	*	*		2	0.9	0.2–3.2
Texas	101	1.1	0.9–1.3		23	1.0	0.6–1.6		7	0.7	0.3–2.0		*	*	*		53	0.7	0.5–1.0
Utah	32	1.9	1.4–2.5		*	*	*		*	*	*		*	*	*		5	2.1	1.0–4.3
Vermont	8	1.6	1.2–2.3		*	*	*		*	*	*		*	*	*		*	*	*
Virginia	46	1.1	0.8–1.5		15	1.2	0.7–2.2		2	0.4	0.1–1.3		*	*	*		5	1.0	0.4–2.5
Washington	63	1.5	1.1–2.1		2	1.2	0.4–3.5		2	0.5	0.2–1.4		*	*	*		11	1.8	0.7–4.9
West Virginia	26	2.0	1.4–2.7		*	*	*		*	*	*		*	*	*		*	*	*
Wisconsin	55	1.5	1.1–2.0		3	1.4	0.4–5.3		2	1.6	0.4–6.2		*	*	*		*	*	*
Wyoming	5	1.4	1.1–1.9		*	*	*		*	*	*		*	*	*		0	1.2	0.5–2.5
Region																			
Northeast	284	1.0	0.8–1.1		45	1.0	0.7–1.3		21	0.7	0.5–1.1		1	1.4	0.4–4.3		74	1.3	0.9–2.0
Midwest	562	1.4	1.3–1.5		68	1.4	1.0–1.9		25	1.5	1.0–2.4		6	2.6	0.9–7.0		55	1.6	1.2–2.3
South	713	1.3	1.2–1.4		196	1.2	1.0–1.3		16	0.5	0.3–1.0		10	2.1	1.2–3.6		108	0.7	0.6–0.9
West	455	1.5	1.3–1.6		40	1.6	1.0–2.5		59	1.0	0.6–1.5		10	1.6	1.1–2.4		208	1.3	1.2–1.6

Abbreviations: No. = number, CI = confidence interval, NH = Non-Hispanic, AI/AN = American Indian or Alaska Native.

†In thousands.

NOTE: Estimated numbers appearing as 0 in this table mean that the estimate is greater than 0 but less than 500 (because estimated numbers are shown in thousands).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2015-2019.

*Low precision; no estimate reported.

NOTE: Only respondents who reported suicide ideation were asked about suicide plans.

SUPPLEMENTARY Table 7. Annual average estimated number[†] and percentage of adults aged ≥18 years who made a suicide attempt during the previous year, by state and sex— National Survey on Drug Use and Health, United States, 2015-2019

	Male			Female		
	No.	%	(95% CI)	No.	%	(95% CI)
US total	565	0.5	0.4–0.5	820	0.6	0.6–0.7
State						
Alabama	10	0.6	0.3–1.0	9	0.5	0.3–0.7
Alaska	2	0.9	0.5–1.6	2	0.7	0.4–1.2
Arizona	13	0.5	0.3–1.0	27	1.0	0.7–1.4
Arkansas	6	0.6	0.3–1.3	8	0.7	0.5–1.1
California	73	0.5	0.4–0.7	75	0.5	0.4–0.7
Colorado	6	0.3	0.1–0.7	18	0.9	0.5–1.5
Connecticut	4	0.3	0.2–0.6	3	0.2	0.1–0.4
Delaware	2	0.6	0.3–1.3	2	0.6	0.3–1.3
District of Columbia	1	0.5	0.2–1.1	2	0.7	0.4–1.4
Florida	22	0.3	0.2–0.4	52	0.6	0.4–0.9
Georgia	22	0.6	0.3–1.1	28	0.7	0.4–1.2
Hawaii	3	0.5	0.3–1.0	4	0.7	0.4–1.2
Idaho	2	0.4	0.2–0.8	4	0.7	0.4–1.2
Illinois	27	0.6	0.4–0.9	24	0.5	0.3–0.8
Indiana	18	0.7	0.5–1.2	25	1.0	0.6–1.6
Iowa	9	0.8	0.5–1.3	11	0.9	0.5–1.6
Kansas	5	0.5	0.2–1.1	10	0.9	0.5–1.4
Kentucky	15	0.9	0.5–1.6	11	0.6	0.4–1.1
Louisiana	12	0.7	0.4–1.4	19	1.1	0.7–1.7
Maine	1	0.3	0.2–0.5	3	0.5	0.3–0.9
Maryland	11	0.5	0.3–1.0	8	0.3	0.2–0.6
Massachusetts	8	0.3	0.2–0.6	19	0.7	0.4–1.1
Michigan	17	0.5	0.3–0.7	25	0.6	0.4–0.9
Minnesota	7	0.3	0.2–0.7	11	0.5	0.3–0.9
Mississippi	6	0.6	0.3–1.1	9	0.8	0.5–1.3
Missouri	10	0.5	0.3–0.8	17	0.7	0.4–1.4

Montana	2	0.5	0.2–0.9		2	0.6	0.4–1.0
Nebraska	3	0.4	0.1–1.1		5	0.7	0.4–1.1
Nevada	6	0.5	0.3–0.9		12	1.0	0.5–2.0
New Hampshire	2	0.3	0.2–0.5		2	0.4	0.3–0.7
New Jersey	5	0.2	0.1–0.4		23	0.6	0.4–1.1
New Mexico	2	0.3	0.2–0.6		5	0.6	0.4–0.9
New York	26	0.4	0.2–0.6		57	0.7	0.5–1.0
North Carolina	21	0.6	0.3–1.1		29	0.7	0.5–1.1
North Dakota	1	0.5	0.2–0.9		2	0.7	0.4–1.2
Ohio	25	0.6	0.4–0.9		27	0.6	0.4–0.8
Oklahoma	7	0.5	0.3–0.9		10	0.7	0.4–1.4
Oregon	7	0.5	0.2–0.9		11	0.7	0.4–1.0
Pennsylvania	17	0.4	0.2–0.5		42	0.8	0.5–1.3
Rhode Island	3	0.7	0.3–1.5		2	0.5	0.3–0.9
South Carolina	6	0.3	0.2–0.7		14	0.7	0.3–1.7
South Dakota	2	0.7	0.3–1.2		2	0.6	0.3–1.0
Tennessee	16	0.6	0.4–1.1		19	0.7	0.5–1.1
Texas	41	0.4	0.3–0.6		54	0.5	0.4–0.7
Utah	7	0.7	0.4–1.2		10	1.0	0.6–1.5
Vermont	1	0.3	0.2–0.6		2	0.7	0.4–1.3
Virginia	18	0.6	0.4–0.9		17	0.5	0.3–0.8
Washington	13	0.5	0.2–1.3		18	0.6	0.3–1.3
West Virginia	6	0.9	0.4–2.3		7	0.9	0.6–1.4
Wisconsin	13	0.6	0.4–1.0		22	1.0	0.6–1.7
Wyoming	1	0.6	0.3–0.9		1	0.6	0.4–1.0
Region							
Northeast	67	0.3	0.3–0.4		153	0.7	0.5–0.8
Midwest	138	0.6	0.5–0.7		179	0.7	0.6–0.8
South	222	0.5	0.4–0.6		299	0.6	0.5–0.7
West	138	0.5	0.4–0.6		188	0.6	0.5–0.8

Abbreviations: No. = number, CI = confidence interval.

†In thousands.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2015-2019

NOTE: Only respondents who reported suicide ideation were asked about suicide attempt(s).

SUPPLEMENTARY Table 8. Annual average estimated number[†] and percentage of adults aged ≥18 years who made a suicide attempt during the previous year, by state and age group — National Survey on Drug Use and Health, United States, 2015-2019

Age groups	18-39 years			40-55 years			≥ 56 years		
	No.	%	(95% CI)	No.	%	(95% CI)	No.	%	(95% CI)
US total	968	1.0	1.0–1.1	252	0.4	0.3–0.5	166	0.2	0.1–0.3
State									
Alabama	15	1.1	0.8–1.7	4	0.4	0.1–1.3	*	*	*
Alaska	3	1.6	1.0–2.4	1	0.4	0.1–1.2	0	0.1	0.0–1.0
Arizona	28	1.4	1.0–2.1	9	0.6	0.3–1.4	3	0.1	0.0–1.0
Arkansas	9	1.0	0.7–1.5	5	0.8	0.3–2.1	1	0.2	0.0–1.2
California	111	0.9	0.7–1.2	26	0.3	0.2–0.6	11	0.1	0.0–0.4
Colorado	17	1.0	0.7–1.5	1	0.1	0.0–0.8	6	0.4	0.1–1.7
Connecticut	6	0.6	0.4–1.0	1	0.1	0.0–0.6	*	*	*
Delaware	3	1.3	0.7–2.4	1	0.3	0.1–1.2	0	0.2	0.0–1.1
District of	2	0.7	0.5–1.2	1	0.8	0.3–2.6	0	0.2	0.0–0.7
Florida	53	1.0	0.8–1.2	17	0.4	0.2–0.9	4	0.1	0.0–0.4
Georgia	28	0.9	0.6–1.3	12	0.5	0.3–1.1	11	0.5	0.1–1.5
Hawaii	5	1.2	0.8–1.8	1	0.2	0.0–0.9	1	0.3	0.1–1.1
Idaho	5	1.0	0.7–1.7	2	0.5	0.2–1.6	*	*	*
Illinois	32	0.8	0.6–1.1	11	0.4	0.2–0.9	9	0.3	0.1–1.1
Indiana	32	1.7	1.1–2.5	3	0.2	0.1–0.7	8	0.4	0.1–1.9
Iowa	13	1.5	1.0–2.1	3	0.5	0.2–1.4	4	0.4	0.1–1.8
Kansas	10	1.2	0.8–1.9	2	0.3	0.1–0.9	3	0.4	0.1–1.6
Kentucky	19	1.5	1.0–2.3	3	0.3	0.1–0.9	4	0.3	0.1–1.4
Louisiana	17	1.2	0.9–1.7	12	1.3	0.7–2.5	2	0.2	0.0–1.3
Maine	4	1.1	0.7–1.5	0	0.0	0.0–0.3	1	0.1	0.0–0.9
Maryland	13	0.7	0.5–1.2	2	0.2	0.0–1.2	4	0.3	0.1–1.1
Massachusetts	25	1.2	0.8–1.8	2	0.1	0.0–0.7	*	*	*
Michigan	25	0.9	0.7–1.2	15	0.8	0.4–1.4	1	0.0	0.0–0.3
Minnesota	14	0.9	0.5–1.4	3	0.3	0.1–1.2	*	*	*
Mississippi	13	1.6	1.0–2.4	1	0.2	0.1–0.5	1	0.2	0.0–1.1
Missouri	21	1.2	0.8–1.9	7	0.5	0.2–1.8	*	*	*
Montana	3	1.1	0.7–1.8	1	0.6	0.2–1.6	*	*	*
Nebraska	5	0.9	0.6–1.4	2	0.6	0.2–1.9	*	*	*
Nevada	12	1.3	0.9–2.0	2	0.4	0.1–1.3	*	*	*
New Hampshire	4	1.1	0.8–1.5	0	0.0	0.0–0.2	*	*	*
New Jersey	20	0.8	0.5–1.2	3	0.2	0.0–0.7	5	0.2	0.0–1.4
New Mexico	6	1.1	0.7–1.6	0	0.1	0.0–0.3	0	0.1	0.0–0.5
New York	57	1.0	0.7–1.3	10	0.2	0.1–0.6	17	0.3	0.1–0.8
North Carolina	25	0.9	0.6–1.3	8	0.4	0.1–1.4	16	0.6	0.2–1.5

North Dakota	3	1.1	0.8–1.6	0	0.2	0.1–0.8	0	0.1	0.0–1.0
Ohio	38	1.2	0.9–1.4	8	0.4	0.2–0.8	6	0.2	0.1–0.6
Oklahoma	10	0.9	0.5–1.6	6	0.7	0.3–1.7	2	0.2	0.0–1.2
Oregon	16	1.4	0.9–2.0	2	0.2	0.1–0.7	*	*	*
Pennsylvania	44	1.3	1.0–1.6	11	0.4	0.1–1.6	4	0.1	0.0–0.4
Rhode Island	4	1.3	0.9–2.0	0	0.0	0.0–0.2	1	0.3	0.0–2.1
South Carolina	18	1.3	0.6–2.6	2	0.2	0.1–0.9	*	*	*
South Dakota	3	1.2	0.9–1.8	1	0.5	0.2–1.5	*	*	*
Tennessee	24	1.3	0.9–1.9	10	0.7	0.3–1.6	*	*	*
Texas	73	0.9	0.7–1.1	14	0.2	0.1–0.6	9	0.1	0.0–0.6
Utah	16	1.5	1.1–2.2	1	0.1	0.0–0.9	1	0.2	0.0–1.1
Vermont	2	1.0	0.6–1.7	1	0.6	0.2–1.7	*	*	*
Virginia	22	0.9	0.6–1.3	12	0.7	0.3–1.3	2	0.1	0.0–0.6
Washington	12	0.5	0.3–0.9	5	0.3	0.1–1.8	14	0.7	0.2–2.5
West Virginia	7	1.5	1.0–2.2	2	0.6	0.3–1.3	4	0.6	0.1–3.0
Wisconsin	20	1.2	0.8–1.8	7	0.6	0.2–1.7	8	0.5	0.2–1.5
Wyoming	2	1.4	1.0–1.9	0	0.1	0.0–0.6	0	0.1	0.0–0.5
Region									
Northeast	166	1.0	0.9–1.2	27	0.2	0.1–0.5	27	0.2	0.1–0.3
Midwest	216	1.1	1.0–1.2	63	0.5	0.3–0.6	39	0.2	0.1–0.4
South	350	1.0	0.9–1.1	111	0.4	0.3–0.6	60	0.2	0.1–0.3
West	236	1.0	0.9–1.2	51	0.3	0.2–0.5	40	0.2	0.1–0.4

Abbreviations: No. = number, CI = confidence interval.

†In thousands.

NOTE: Estimated numbers appearing as 0 in this table mean that the estimate is greater than 0 but less than 500 (because estimated numbers are shown in thousands).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2015-2019.

*Low precision; no estimate reported.

NOTE: Only respondents who reported suicide ideation were asked about suicide attempt(s).

SUPPLEMENTARY Table 9. Annual average estimated number[†] and percentage of adults aged ≥18 years who made a suicide attempt during the previous year, by state and race/ethnicity — National Survey on Drug Use and Health, United States, 2015-2019

Race/Ethnicity	NH white			NH black			NH Asian			NH AI/AN			Hispanic		
	No.	%	(95% CI)	No.	%	(95% CI)	No.	%	(95% CI)	No.	%	(95% CI)	No.	%	(95% CI)
US total	805	0.5	0.5–0.6	204	0.7	0.6–0.8	64	0.5	0.3–0.6	11	0.8	0.5–1.1	243	0.6	0.5–0.7
State															
Alabama	14	0.5	0.3–0.9	5	0.5	0.3–1.0	*	*	*	*	*	*	*	*	*
Alaska	1	0.4	0.2–0.7	0	1.3	0.3–5.0	0	0.6	0.1–3.0	1	1.4	0.5–4.0	1	1.8	0.6–5.2
Arizona	18	0.6	0.3–1.0	1	0.2	0.1–0.9	*	*	*	1	0.5	0.1–1.6	19	1.3	0.9–1.9
Arkansas	9	0.5	0.3–0.9	4	1.4	0.5–3.5	*	*	*	*	*	*	1	1.0	0.3–3.3
California	45	0.4	0.3–0.6	13	0.8	0.4–1.5	15	0.3	0.1–0.8	*	*	*	68	0.6	0.5–0.9
Colorado	19	0.6	0.4–1.1	*	*	*	*	*	*	*	*	*	2	0.2	0.1–0.7
Connecticut	4	0.2	0.1–0.4	1	0.4	0.1–1.4	*	*	*	*	*	*	1	0.3	0.1–0.8
Delaware	3	0.7	0.4–1.2	0	0.3	0.1–0.7	*	*	*	*	*	*	0	0.9	0.2–4.3
District of	0	0.1	0.0–0.4	2	1.0	0.5–1.9	0	1.0	0.2–4.0	*	*	*	1	1.1	0.3–4.5
Florida	43	0.5	0.3–0.7	14	0.6	0.4–1.1	2	0.5	0.2–1.8	*	*	*	13	0.3	0.2–0.5
Georgia	25	0.6	0.3–1.1	20	0.9	0.6–1.3	*	*	*	*	*	*	4	0.6	0.2–1.5
Hawaii	1	0.3	0.1–0.8	*	*	*	1	0.4	0.2–0.8	*	*	*	1	1.2	0.5–3.1
Idaho	6	0.6	0.4–0.9	*	*	*	*	*	*	*	*	*	0	0.2	0.0–0.5
Illinois	26	0.4	0.3–0.6	14	1.1	0.4–2.6	1	0.2	0.0–1.7	*	*	*	10	0.7	0.3–1.4
Indiana	36	0.9	0.6–1.3	2	0.5	0.2–1.6	*	*	*	*	*	*	*	*	*
Iowa	16	0.8	0.5–1.2	1	1.9	0.6–5.8	*	*	*	*	*	*	1	1.2	0.4–3.8
Kansas	9	0.5	0.3–0.9	*	*	*	*	*	*	*	*	*	3	1.4	0.5–3.9
Kentucky	23	0.8	0.5–1.2	1	0.3	0.0–2.2	*	*	*	*	*	*	0	0.3	0.0–1.9
Louisiana	20	0.9	0.6–1.5	9	0.8	0.4–1.6	*	*	*	*	*	*	0	0.2	0.0–0.7
Maine	4	0.4	0.3–0.6	*	*	*	*	*	*	*	*	*	*	*	*
Maryland	11	0.5	0.2–0.9	7	0.5	0.2–1.2	*	*	*	*	*	*	1	0.2	0.0–0.5
Massachusetts	16	0.4	0.2–0.7	3	0.9	0.2–3.5	3	0.9	0.3–3.2	*	*	*	4	0.7	0.3–1.4
Michigan	31	0.5	0.4–0.7	7	0.7	0.4–1.2	1	0.3	0.1–1.6	*	*	*	1	0.2	0.0–0.7
Minnesota	14	0.4	0.2–0.6	*	*	*	*	*	*	*	*	*	*	*	*
Mississippi	7	0.5	0.3–0.9	6	0.7	0.4–1.3	*	*	*	*	*	*	*	*	*
Missouri	22	0.6	0.3–1.0	3	0.5	0.2–1.1	*	*	*	*	*	*	1	0.9	0.2–4.0
Montana	3	0.4	0.2–0.7	*	*	*	*	*	*	1	2.7	1.3–5.3	0	0.4	0.1–1.8
Nebraska	6	0.5	0.3–0.9	0	0.3	0.0–1.8	*	*	*	*	*	*	1	1.0	0.4–2.8
Nevada	7	0.6	0.2–1.5	3	1.8	0.7–4.3	1	0.7	0.2–2.7	*	*	*	4	0.8	0.4–1.5
New	3	0.3	0.2–0.4	*	*	*	*	*	*	*	*	*	1	2.1	0.7–6.2
New Jersey	15	0.4	0.2–0.8	5	0.5	0.2–1.5	5	0.7	0.3–1.4	*	*	*	4	0.3	0.1–0.8
New Mexico	2	0.3	0.1–0.5	*	*	*	*	*	*	1	0.8	0.2–3.0	4	0.6	0.4–0.9
New York	41	0.5	0.3–0.7	10	0.5	0.2–0.9	4	0.3	0.1–0.8	*	*	*	25	0.9	0.5–1.8
North Carolina	38	0.7	0.5–1.1	8	0.5	0.2–1.4	*	*	*	0	0.2	0.0–1.4	4	0.6	0.2–1.7
North Dakota	2	0.5	0.3–0.7	*	*	*	*	*	*	0	0.8	0.2–3.5	0	0.2	0.0–1.3
Ohio	40	0.5	0.4–0.7	5	0.5	0.2–1.1	*	*	*	*	*	*	3	1.1	0.4–3.1

Oklahoma	10	0.5	0.2–1.0	1	0.6	0.2–1.7	*	*	*	2	1.0	0.4–2.9	0	0.1	0.0–0.2
Oregon	12	0.5	0.3–0.8	*	*	*	0	0.1	0.0–0.7	*	*	*	4	1.1	0.5–2.7
Pennsylvania	33	0.4	0.3–0.6	9	1.0	0.6–1.6	1	0.3	0.0–2.2	*	*	*	12	2.0	0.6–6.9
Rhode Island	4	0.6	0.3–1.0	0	0.2	0.0–1.4	0	0.2	0.0–1.4	*	*	*	1	1.2	0.8–1.8
South Carolina	8	0.3	0.2–0.6	6	0.6	0.3–1.1	*	*	*	*	*	*	*	*	*
South Dakota	3	0.5	0.3–0.8	*	*	*	*	*	*	1	2.0	0.9–4.5	*	*	*
Tennessee	21	0.5	0.3–0.9	10	1.3	0.7–2.2	*	*	*	*	*	*	2	1.1	0.3–3.3
Texas	41	0.4	0.3–0.7	13	0.6	0.3–1.0	5	0.5	0.1–2.0	*	*	*	32	0.4	0.3–0.7
Utah	11	0.7	0.4–1.0	*	*	*	*	*	*	*	*	*	4	1.4	0.7–2.8
Vermont	3	0.5	0.3–0.9	*	*	*	*	*	*	*	*	*	*	*	*
Virginia	17	0.4	0.3–0.7	10	0.9	0.4–1.8	3	0.8	0.3–2.1	*	*	*	4	0.8	0.3–2.1
Washington	21	0.5	0.3–1.0	*	*	*	*	*	*	*	*	*	1	0.1	0.0–0.5
West Virginia	13	1.0	0.5–1.7	0	0.2	0.0–1.3	*	*	*	*	*	*	*	*	*
Wisconsin	29	0.8	0.5–1.3	3	1.1	0.4–3.6	*	*	*	*	*	*	2	0.7	0.2–2.6
Wyoming	2	0.6	0.4–0.8	*	*	*	*	*	*	*	*	*	0	1.0	0.4–2.7
Region															
Northeast	122	0.4	0.3–0.5	28	0.6	0.4–0.9	13	0.5	0.3–0.8	0	0.6	0.1–2.7	47	0.8	0.5–1.4
Midwest	233	0.6	0.5–0.7	37	0.8	0.5–1.1	9	0.5	0.3–1.1	2	0.9	0.5–1.8	25	0.7	0.5–1.1
South	302	0.5	0.5–0.6	118	0.7	0.6–0.8	14	0.5	0.2–0.9	3	0.7	0.3–1.5	63	0.4	0.3–0.6
West	148	0.5	0.4–0.6	21	0.8	0.5–1.3	27	0.4	0.2–0.9	5	0.8	0.5–1.4	107	0.7	0.6–0.9

Abbreviations: No. = number, CI = confidence interval, NH = Non-Hispanic, AI/AN = American Indian or Alaska Native.

†In thousands.

NOTE: Estimated numbers appearing as 0 in this table mean that the estimate is greater than 0 but less than 500 (because estimated numbers are shown in thousands).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2015-2019.

*Low precision; no estimate reported.

NOTE: Only respondents who reported suicide ideation were asked about suicide attempt(s).