



## COVID-19

# COVID-19 Vaccines That Require 2 Shots

Updated Nov. 18, 2021

**NOTICE:** CDC now recommends that children between the ages of 5 and 11 years receive the Pfizer-BioNTech pediatric COVID-19 Vaccine. Learn more about [vaccines for children and teens](#).

Learn more about who is [eligible for a COVID-19 booster shot](#).

## What You Need to Know

- If you receive a Pfizer-BioNTech or Moderna COVID-19 vaccine, you will need 2 initial shots to get the most protection.
- **CDC does not recommend mixing products for a two dose primary vaccine series or an additional primary dose.** If you received a Pfizer-BioNTech or Moderna COVID-19 vaccine, you should get the same product when you need a second shot or additional primary dose. However, mixing and matching COVID-19 vaccines is allowed for booster shots.
- **You should get your second shot even if you have side effects after the first shot,** unless a vaccination provider or your doctor tells you [not to get it](#).
- If you have a [weakened immune system](#) due to other diseases or medications, you can receive a COVID-19 vaccine if you have not had a severe allergic reaction to any of the [ingredients](#) in the vaccine.
- People with [moderately to severely compromised immune systems](#) should [receive an additional primary dose](#) of mRNA COVID-19 vaccine (Pfizer-BioNTech or Moderna) 28 days after their initial 2 doses.
- Certain groups of people who received both doses of the Pfizer-BioNTech COVID-19 vaccine or Moderna COVID-19 vaccine are eligible to get a [booster shot](#).

## Timing of Your Second Shot

The timing between your first and second shots depends on which vaccine you received. If you received the:

### [Pfizer-BioNTech COVID-19 vaccine](#)

Get your second shot **3 weeks (or 21 days)** after your first

### [Moderna COVID-19 vaccine](#)

Get your second shot **4 weeks (or 28 days)** after your first

You should **get your second shot as close to the recommended 3-week or 4-week interval as possible.** However, your second dose may be given up to [6 weeks \(42 days\)](#) after the first dose, if necessary. You should not get the second dose early. There is currently limited information on the effectiveness of receiving your second shot earlier than recommended or later than 6 weeks after the first shot.

However, if you do receive your second shot of COVID-19 vaccine up to 4 days before or at any time after the recommended date, you do not have to restart the vaccine series, and you can be considered fully vaccinated. This guidance might be updated as more information becomes available.

# Scheduling Your Second Shot

- Planning for your second shot is important.
- If you need help scheduling your vaccination appointment for your second shot, contact the location that set up your first appointment.
- If you are having trouble or have questions about using a vaccination management or scheduling system, reach out to the organization that enrolled you in the system. This may be your state or local health department, employer, or vaccination provider.
- Scheduling an appointment for your second shot at the time you get your first shot is recommended, but not required.
- If you need to get your second shot in a location that is different from where you received your first shot (for example, if you moved to a different state or attend school in a different state), there are several ways you can [find a vaccine provider](#) for your second dose.



## Your CDC COVID-19 Vaccination Record Card and Your Second Shot

At your first vaccination appointment, you should have received a CDC COVID-19 Vaccination Record card that tells you what COVID-19 vaccine you received, the date you received it, and where you received it.

- **Bring your vaccination card with you to your second shot appointment** so your provider can fill in the information about your second dose.
- If you did not receive a CDC COVID-19 Vaccination Record card at your first appointment, contact the vaccination provider site where you got your first shot or your [state health department](#) to find out how you can get a card.

Learn more about what to do if you need a copy of your [CDC COVID-19 Vaccination Record card](#).

## After Getting Your Second Shot

You may experience [side effects](#) after getting a COVID-19 vaccine. These are normal signs that your body is building protection. Get helpful tips on [how to reduce any pain or discomfort](#).

Cases of myocarditis and pericarditis in adolescents and young adults have been reported more often after getting the second dose than after the first dose of one of the two mRNA COVID-19 vaccines, Pfizer-BioNTech or Moderna. **These reports are rare and the known and potential benefits of COVID-19 vaccination outweigh the known and potential risks, including the [possible risk of myocarditis or pericarditis](#).**

Use [v-safe](#) on your smartphone to tell CDC about any side effects after getting the COVID-19 vaccine. If you [enter your second shot](#) in your [v-safe](#) account, the system will send you daily health check-ins. Please note that [v-safe](#) is not automatically notified when you receive a second shot of vaccine, so you must enter the information yourself.

## When You Are Fully Vaccinated

People are considered fully vaccinated:

- 2 weeks after their second shot in a 2-dose series, like the Pfizer-BioNTech or Moderna vaccines, or
- 2 weeks after a single-shot vaccine, like Johnson & Johnson's Janssen COVID-19 Vaccine

Administration of an [additional primary dose](#) or a [booster dose](#) is not required to be considered fully vaccinated for public health purposes at this time.

You are **not** fully vaccinated if:

- It has been less than 2 weeks since your single-dose shot
- It has been less than 2 weeks since your second shot of a two-dose vaccine
- You haven't received your second dose of a two-dose vaccine

## Related Pages

- › [When Getting Your Vaccine](#)
- › [Types of Vaccines Available](#)
- › [Possible Side Effects](#)

Last Updated Nov. 18, 2021