

Weekly U.S. Influenza Surveillance Report

Updated January 28, 2022

FLUVIEW



A Weekly Influenza Surveillance Report Prepared by the Influenza Division

Note: CDC is tracking the COVID-19 pandemic in a weekly publication called [COVID Data Tracker Weekly Review](https://www.cdc.gov/coronavirus/2019-ncov/covid-data/covidview/).
(<https://www.cdc.gov/coronavirus/2019-ncov/covid-data/covidview/>)

Key Updates for Week 3, ending January 22, 2022

The percent of specimens testing positive for influenza remains stable, indicating that influenza virus circulation has remained at similar levels during the past two weeks, even while overall levels of respiratory illness have declined.

Viruses

Clinical Lab

1.9%
positive for influenza
this week

Public Health Lab

The majority of viruses
detected are influenza A(H3N2).

Virus Characterization

Genetic characterization and
antiviral susceptibility are
summarized in this report.

</flu/weekly/#VirusCharacterization>

Illness

Outpatient Respiratory Illness

2.8%
of visits to a health care provider are for respiratory illness this week
(*above baseline*)

Outpatient Respiratory Illness: Activity Map

This week, 6 jurisdictions experienced moderate activity and 12 jurisdictions experienced high or very high activity.

Long-term Care Facilities

0.8%

of facilities reported
≥ 1 influenza-positive test
among residents this week

Severe Disease

FluSurv-NET 4.3 per 100,000 cumulative hospitalization rate	HHS Protect Hospitalizations 971 patients admitted to hospitals with influenza this week.
NCHS Mortality 28.4% of deaths attributed to pneumonia, influenza, or COVID-19 this week (<i>above threshold</i>)	Pediatric Deaths 0 influenza-associated deaths reported this week for a total of 5 so far this season

All data are preliminary and may change as more reports are received.

A description of the CDC influenza surveillance system, including methodology and detailed descriptions of each data component is available on the [surveillance methods](http://www.cdc.gov/flu/weekly/overview.htm) (<http://www.cdc.gov/flu/weekly/overview.htm>) page.

Additional information on the current and previous influenza seasons for each surveillance component are available on [FluView Interactive](https://www.cdc.gov/flu/weekly/fluviewinteractive.htm) (<https://www.cdc.gov/flu/weekly/fluviewinteractive.htm>).

Key Points

- The percent of specimens testing positive for influenza remains stable, indicating that influenza virus circulation has remained at similar levels during the past two weeks, even while overall levels of respiratory illness have declined. While influenza activity is difficult to predict, it is expected to continue for several more weeks.
- The majority of influenza viruses detected are A(H3N2). Most of the H3N2 viruses identified so far this season are genetically closely related to the vaccine virus, but there are some antigenic differences that have developed as H3N2 viruses have continued to evolve.
- The percentage of outpatient visits due to respiratory illness decreased nationally again this week but remains above baseline. Influenza is contributing to levels of respiratory illness, but other respiratory viruses are also circulating. The relative contribution of influenza varies by location.
- The number of hospital admissions reported to HHS Protect declined slightly again this week.
- The cumulative hospitalization rate in the FluSurv-NET system is higher than the rate for the entire 2020-2021 season, but lower than the rate seen at this time during the four seasons preceding the COVID-19 pandemic.
- CDC estimates that so far this season there have been at least 2 million flu illnesses, 20,000 hospitalizations, and 1,200 deaths from flu.
- There's still time to get vaccinated. An annual flu vaccine is the best way to protect against flu and its potentially serious complications. CDC recommends everyone 6 months and older get a flu vaccine.
- Flu vaccination coverage so far is lower this season compared to last.
- Flu vaccines are available at many different locations, including pharmacies and health departments. Visit www.vaccines.gov to find a flu vaccine near you.
- There are also flu antiviral drugs that can be used to treat flu illness.

U.S. Virologic Surveillance

(https://www.cdc.gov/flu/weekly/overview.htm#anchor_1633697372803)

Influenza A(H3N2) viruses have been the most frequently detected influenza viruses this season. For regional and state level data and age group distribution, please visit [FluView Interactive](https://gis.cdc.gov/grasp/fluview/fluportaldashboard.html) (<https://gis.cdc.gov/grasp/fluview/fluportaldashboard.html>). Viruses known to be associated with recent live attenuated influenza vaccine (LAIV) receipt or found upon further testing to be a vaccine virus are not included as they are not circulating influenza viruses.

Clinical Laboratories

The results of tests performed by clinical laboratories nationwide are summarized below. Data from clinical laboratories (the percentage of specimens tested that are positive for influenza) are used to monitor whether influenza activity is increasing or decreasing.

	Week 3	Data Cumulative since October 3, 2021 (Week 40)
No. of specimens tested	79,667	1,360,940
No. of positive specimens (%)	1,543 (1.9%)	35,913 (2.6%)
<i>Positive specimens by type</i>		
Influenza A	1,503 (97.4%)	35,104 (97.7%)
Influenza B	40 (2.6%)	809 (2.3%)

(<http://gis.cdc.gov/grasp/fluview/fluportaldashboard.html>)

[View Chart Data \(/flu/weekly/weeklyarchives2021-2022/data/whoAllregt_cl03.html\)](/flu/weekly/weeklyarchives2021-2022/data/whoAllregt_cl03.html) | [View Full Screen \(/flu/weekly/weeklyarchives2021-2022/WhoNPHL03.html\)](/flu/weekly/weeklyarchives2021-2022/WhoNPHL03.html)

Public Health Laboratories

The results of tests performed by public health laboratories nationwide are summarized below. Data from public health laboratories are used to monitor the proportion of circulating viruses that belong to each influenza subtype/lineage.

	Week 3	Data Cumulative since October 3, 2021 (Week 40)
No. of specimens tested	38,977	490,190
No. of positive specimens	503	10,012
<i>Positive specimens by type/subtype</i>		
Influenza A	502 (99.8%)	9,928 (99.2%)
(H1N1)pdm09	0	5 (0.1%)
H3N2	177 (100%)	6,919 (99.9%)
H3N2v	0	1 (<0.1%)

	Week 3	Data Cumulative since October 3, 2021 (Week 40)
Subtyping not performed	325	3,003
Influenza B	1 (0.2%)	84 (0.8%)
Yamagata lineage	0	1 (3.0%)
Victoria lineage	0	32 (97.0%)
Lineage not performed	1	51

(<http://gis.cdc.gov/grasp/fluview/fluportaldashboard.html>)

[View Chart Data \(/flu/weekly/weeklyarchives2021-2022/data/whoAllregt_phl03.html\)](/flu/weekly/weeklyarchives2021-2022/data/whoAllregt_phl03.html) | [View Full Screen \(/flu/weekly/weeklyarchives2021-2022/WhoPHL03.html\)](/flu/weekly/weeklyarchives2021-2022/WhoPHL03.html)

Additional virologic surveillance information for current and past seasons:
[Surveillance Methods \(https://wcms-wp.cdc.gov/flu/weekly/overview.htm#anchor_1633697372803\)](https://wcms-wp.cdc.gov/flu/weekly/overview.htm#anchor_1633697372803) | [FluView Interactive: National, Regional, and State Data \(http://gis.cdc.gov/grasp/fluview/fluportaldashboard.html\)](#) or [Age Data \(https://gis.cdc.gov/grasp/fluview/flu_by_age_virus.html\)](https://gis.cdc.gov/grasp/fluview/flu_by_age_virus.html)

Influenza Virus Characterization

(/flu/weekly/overview.htm#anchor_1633697390939)

CDC performs [genetic \(https://www.cdc.gov/flu/professionals/laboratory/genetic-characterization.htm\)](https://www.cdc.gov/flu/professionals/laboratory/genetic-characterization.htm) and [antigenic \(https://www.cdc.gov/flu/professionals/laboratory/antigenic.htm\)](https://www.cdc.gov/flu/professionals/laboratory/antigenic.htm) characterization of U.S. viruses submitted from state and local public health laboratories using the Right Size Roadmap submission guidance. These data are used to compare how similar the currently circulating influenza viruses are to the reference viruses representing viruses contained in the current influenza vaccines. The data are also used to monitor evolutionary changes that continually occur in influenza viruses circulating in humans. CDC also tests susceptibility of circulating influenza viruses to antiviral medications including the neuraminidase inhibitors (oseltamivir, zanamivir, and peramivir) and the PA endonuclease inhibitor baloxavir.

CDC has genetically characterized 460 influenza viruses collected since October 3, 2021. Most of the H3N2 viruses so far are genetically closely related to the vaccine virus, but there are some antigenic differences that have developed as H3N2 viruses have continued to evolve. Virus antigenic data will be reported later this season when a sufficient number of specimens have been tested.

Virus Subtype or Lineage	Genetic Characterization				
	Total No. of Subtype/Lineage Tested	HA Clade	Number (% of subtype/lineage tested)	HA Subclade	Number (% of subtype/lineage tested)
A/H1	3				
		6B.1A	3 (100%)	5a.1	2 (66.7%)
				5a.2	1 (33.3%)
A/H3	437				
		3C.2a1b	437(100%)	1a	1 (0.2%)
				1b	1 (0.2%)
				2a	0
				2a.1	0
				2a.2	435 (99.5%)
		3C.3a	0	3a	0
B/Victoria	20				
		V1A	20 (100%)	V1A	0
				V1A.1	0
				V1A.3	9 (45.0%)
				V1A.3a	0
				V1A.3a.1	0
				V1A.3a.2	11 (55.0%)
B/Yamagata	0				
		Y3	0		

CDC assesses susceptibility of influenza viruses to the antiviral medications including the neuraminidase inhibitors (oseltamivir, zanamivir, and peramivir) and the PA endonuclease inhibitor baloxavir using next generation sequence analysis supplemented by laboratory assays. Information about antiviral susceptibility test methods can be found at [U.S. Influenza Surveillance: Purpose and Methods | CDC \(https://www.cdc.gov/flu/weekly/overview.htm#anchor_1633697390939\)](https://www.cdc.gov/flu/weekly/overview.htm#anchor_1633697390939).

Viruses collected in the United States since October 3, 2021, were tested for antiviral susceptibility as follows:

Antiviral Medication			Total Viruses	A/H1	A/H3	B/Victoria	B/Yamagata
Neuraminidase Inhibitors	Oseltamivir	Viruses Tested	484	3	461	20	0
		Reduced Inhibition	(0.0%)	(0.0%)	(0.0%)	(0.0%)	(0.0%)
		Highly Reduced Inhibition	(0.0%)	(0.0%)	(0.0%)	(0.0%)	(0.0%)
	Peramivir	Viruses Tested	484	3	461	20	0
		Reduced Inhibition	(0.0%)	(0.0%)	(0.0%)	(0.0%)	(0.0%)
		Highly Reduced Inhibition	(0.0%)	(0.0%)	(0.0%)	(0.0%)	(0.0%)
	Zanamivir	Viruses Tested	484	3	461	20	0
		Reduced Inhibition	(0.0%)	(0.0%)	(0.0%)	(0.0%)	(0.0%)
		Highly Reduced Inhibition	(0.0%)	(0.0%)	(0.0%)	(0.0%)	(0.0%)
PA Cap-Dependent Endonuclease Inhibitor	Baloxavir	Viruses Tested	479	3	456	20	0
		Reduced Susceptibility	(0.0%)	(0.0%)	(0.0%)	(0.0%)	(0.0%)

High levels of resistance to the adamantanes (amantadine and rimantadine) persist among influenza A(H1N1)pdm09 and influenza A(H3N2) viruses (the adamantanes are not effective against influenza B viruses). Therefore, use of these antivirals for treatment and prevention of influenza A virus infection is not recommended and data from adamantane resistance testing are not presented.

Outpatient Respiratory Illness Surveillance
(https://www.cdc.gov/flu/weekly/overview.htm#anchor_1539281266932)

The U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet) monitors outpatient visits for influenza-like illness [ILI (fever plus cough or sore throat)], not laboratory-confirmed influenza, and will therefore capture respiratory illness visits due to infection with any pathogen that can present with similar symptoms, including influenza, SARS-CoV-2, and RSV. Due to the COVID-19 pandemic, health care-seeking behaviors have changed, and people may be accessing the health care system in alternative settings not captured as a part of ILINet or at a different point in their illness than they might have before the pandemic. Therefore, it is important to evaluate syndromic surveillance data, including that from ILINet, in the context of other sources of surveillance data to obtain a complete and accurate picture of influenza, SARS-CoV-2, and other respiratory virus activity. CDC is tracking the COVID-19 pandemic in a weekly publication called [COVID Data Tracker Weekly Review](https://www.cdc.gov/coronavirus/2019-ncov/covid-data/covidview/index.html) (<https://www.cdc.gov/coronavirus/2019-ncov/covid-data/covidview/index.html>). Information about other respiratory virus activity can be found on [CDC's National Respiratory and Enteric Virus Surveillance System \(NREVSS\) website](https://www.cdc.gov/surveillance/nrevss/index.html) (<https://www.cdc.gov/surveillance/nrevss/index.html>).

Outpatient Respiratory Illness Visits

Nationwide, during week 3, 2.8% of patient visits reported through ILINet were due to respiratory illness that included fever plus a cough or sore throat, also referred to as ILI. This percentage is above the national baseline. Five of the 10 HHS regions are above their region-specific baselines (Regions 3,4,6,7, and 10), and the remaining regions are below their baselines. Multiple respiratory viruses are co-circulating, and the relative contribution of influenza virus infection to ILI varies by location.

<http://gis.cdc.gov/grasp/fluview/fluportaldashboard.html>

* Effective October 3, 2021 (week 40), the ILI definition (fever plus cough or sore throat) no longer includes “without a known cause other than influenza.”

[View Chart Data \(current season only\) \(/flu/weekly/weeklyarchives2021-2022/data/senAllregt03.html\)](/flu/weekly/weeklyarchives2021-2022/data/senAllregt03.html) | [View Full Screen](#)
[\(/flu/weekly/weeklyarchives2021-2022/ILI03.html\)](/flu/weekly/weeklyarchives2021-2022/ILI03.html)

Outpatient Respiratory Illness Visits by Age Group

More than 70% of ILINet participants provide both the number of patient visits for respiratory illness and the total number of patient visits for the week broken out by age group. Data from this subset of providers are used to calculate the percentages of patient visits for respiratory illness by age group.

The percentage of visits for respiratory illness reported in ILINet are trending downward for all age groups (0–4 years, 5–24 years, 25-49 years, 50–64 years, and 65+ years).



<http://gis.cdc.gov/grasp/fluview/fluportaldashboard.html>

* Effective October 3, 2021 (week 40), the ILI definition (fever plus cough or sore throat) no longer includes “without a known cause other than influenza.”

[View Chart Data \(/flu/weekly/weeklyarchives2021-2022/data/iliage03.html\)](/flu/weekly/weeklyarchives2021-2022/data/iliage03.html) | [View Full Screen \(/flu/weekly/weeklyarchives2021-2022/ILIAge03.html\)](/flu/weekly/weeklyarchives2021-2022/ILIAge03.html)

Outpatient Respiratory Illness Activity Map

Data collected in ILINet are used to produce a measure of [ILI activity](#)* [by state/jurisdiction and Core Based Statistical Areas \(CBSA\).](https://www.cdc.gov/flu/weekly/overview.htm#anchor_1633697504110)

Activity Level	Number of Jurisdictions		Number of CBSAs	
	Week 3 (Week ending Jan. 22, 2022)	Week 2 (Week ending Jan. 15, 2022)	Week 3 (Week ending Jan. 22, 2022)	Week 2 (Week ending Jan. 15, 2022)
Very High	1	2	10	14
High	11	14	44	87

Activity Level	Number of Jurisdictions		Number of CBSAs	
	Week 3 (Week ending Jan. 22, 2022)	Week 2 (Week ending Jan. 15, 2022)	Week 3 (Week ending Jan. 22, 2022)	Week 2 (Week ending Jan. 15, 2022)
Moderate	6	12	71	103
Low	10	14	151	159
Minimal	25	13	379	305
Insufficient Data	2	0	274	261

*Data collected in ILINet may disproportionately represent certain populations within a jurisdiction or CBSA, and therefore, may not accurately depict the full picture of influenza activity for the entire jurisdiction or CBSA. Differences in the data presented here by CDC and independently by some health departments likely represent differing levels of data completeness with data presented by the health department likely being the more complete.

Additional information about medically attended visits for ILI for current and past seasons:
Surveillance Methods (https://wcms-wp.cdc.gov/flu/weekly/overview.htm#anchor_1539281266932) | FluView Interactive: National, Regional, and State Data (<http://gis.cdc.gov/grasp/fluview/fluportaldashboard.html>) or ILI Activity Map (<https://gis.cdc.gov/grasp/fluview/main.html>)

Long-term Care Facility (LTCF) Surveillance

(https://www.cdc.gov/flu/weekly/overview.htm#anchor_1633698386507)

LTCFs (e.g., nursing homes/skilled nursing, long-term care for the developmentally disabled, and assisted living facilities) from all 50 states and U.S. territories report data on influenza virus infections among residents through the [National Healthcare Safety Network \(NHSN\) Long-term Care Facility Component](https://www.cdc.gov/nhsn/ltc/index.html) (<https://www.cdc.gov/nhsn/ltc/index.html>). During week 3, 114 (0.8%) of 14,249 reporting LTCFs reported at least one influenza positive test among their residents.

(</flu/weekly/weeklyarchives2021-2022/LTCF03.html>)[View Chart Data](#)  (</flu/weekly/weeklyarchives2021-2022/data/LTCFData03.csv>) | [View Full Screen](#) (</flu/weekly/weeklyarchives2021-2022/LTCF03.html>)

Additional information about long-term care facility surveillance:
Surveillance Methods (https://www.cdc.gov/flu/weekly/overview.htm#anchor_1633698386507) | [Additional Data](#)  (<https://data.cms.gov/covid-19/covid-19-nursing-home-data>)

Hospitalization Surveillance

(http://www.cdc.gov/flu/weekly/overview.htm#anchor_1634240269291)

FluSurv–NET

The Influenza Hospitalization Surveillance Network (FluSurv-NET) conducts population-based surveillance for laboratory-confirmed influenza-related hospitalizations in select counties in 14 states and represents approximately 9% of the U.S. population. FluSurv-NET hospitalization data are preliminary. As data are received each week, prior case counts and rates are updated accordingly.

A total of 1,250 laboratory-confirmed influenza-associated hospitalizations were reported by FluSurv-NET sites between October 1, 2021, and January 22, 2022. The overall cumulative hospitalization rate is 4.3 per 100,000 population. This cumulative hospitalization rate is higher than the cumulative in-season hospitalization rate observed for week 3 during the 2020-2021 season (0.5 per 100,000), but lower than the in-season rates observed for week 3 during the 4 seasons preceding the COVID-19 pandemic (these ranged from 14.8 to 41.9 per 100,000 during the 2016-17 through 2019-20 seasons).

When examining rates by age, the highest rate of hospitalization per 100,000 population was among adults aged ≥ 65 years (11.3); within this group, rates were highest among adults aged ≥ 85 years (24.0). Among persons aged < 65 years, hospitalization rates per 100,000 population were highest among children aged 0-4 years (6.2) followed by adults aged 50-64 years (4.0). When examining rates by race and ethnicity, the highest rate of hospitalization per 100,000 population was among non-Hispanic American Indian or Alaska Native persons (5.4) followed by non-Hispanic Black persons (5.0).

Among 1,250 hospitalizations, 1,178 (94.2%) were associated with influenza A virus, 62 (5.0%) with influenza B virus, 5 (0.4%) with influenza A virus and influenza B virus co-infection, and 5 (0.4%) with influenza virus for which the type was not determined. Among 295 hospitalizations with influenza A subtype information, 292 (99.0%) were A(H3N2) and 3 (1.0%) were A(H1N1)pdm09. Based on preliminary data, of the 1,250 laboratory-confirmed influenza-associated hospitalizations, 1.9% also tested positive for SARS-CoV-2.

Among 444 hospitalized adults with information on underlying medical conditions, 89.6% had at least one reported underlying medical condition. Among 52 hospitalized children with information on underlying medical conditions, 55.8% had at least one reported underlying medical condition; the most commonly reported was asthma.

Additional FluSurv-NET hospitalization surveillance information for current and past seasons and additional age groups:
Surveillance Methods (https://www.cdc.gov/flu/weekly/overview.htm#anchor_1633698456778) | FluView Interactive: Rates by Age, Sex, and Race/Ethnicity (<http://gis.cdc.gov/GRASP/Fluview/FluHospRates.html>) or Data on Patient Characteristics (<http://gis.cdc.gov/grasp/fluview/FluHospChars.html>)

FluSurv-Net data are used to generate national estimates of the total numbers of influenza cases, medical visits, hospitalizations, and deaths. This season, CDC is reporting preliminary cumulative in-season estimates, which are available at <https://www.cdc.gov/flu/about/burden/preliminary-in-season-estimates.htm> (<https://www.cdc.gov/flu/about/burden/preliminary-in-season-estimates.htm>)

HHS–Protect Hospitalization Surveillance

Hospitals report to HHS-Protect the number of patients admitted with laboratory-confirmed influenza. During week 3, 971 patients with laboratory-confirmed influenza were admitted to the hospital.

(</flu/weekly/weeklyarchives2021-2022/Protect03.html>)[View Chart Data](#)  (</flu/weekly/weeklyarchives2021-2022/data/ProtectData03.csv>) | [View Full Screen](#) (</flu/weekly/weeklyarchives2021-2022/Protect03.html>)

Additional HHS Protect hospitalization surveillance information:
Surveillance Methods (https://www.cdc.gov/flu/weekly/overview.htm#anchor_1633698474047) | [Additional Data](#) 
(<https://healthdata.gov/Hospital/COVID-19-Reported-Patient-Impact-and-Hospital-Capa/anag-cw7u>)

Mortality Surveillance

(https://www.cdc.gov/flu/weekly/overview.htm#anchor_1634311686144)

National Center for Health Statistics (NCHS) Mortality Surveillance

Based on NCHS mortality surveillance data available on January 27, 2022, 28.4% of the deaths that occurred during the week ending January 22, 2022 (week 3), were due to pneumonia, influenza, and/or COVID-19 (PIC). This percentage is above the epidemic threshold of 7.1% for this week. Among the 5,940 PIC deaths reported for this week, 5,160 had COVID-19 listed as an underlying or contributing cause of death on the death certificate, and 27 listed influenza, indicating that current PIC mortality is due primarily to COVID-19 and not influenza. The data presented are preliminary and may change as more data are received and processed.

[View Chart Data](https://gis.cdc.gov/grasp/fluview/mortality.html)  [\(/flu/weekly/weeklyarchives2021-2022/data/NCHSData03.csv\)](/flu/weekly/weeklyarchives2021-2022/data/NCHSData03.csv) | [View Full Screen](#) [\(/flu/weekly/weeklyarchives2021-2022/NCHS03.html\)](/flu/weekly/weeklyarchives2021-2022/NCHS03.html)

Additional pneumonia, influenza and COVID-19 mortality surveillance information for current and past seasons:
[Surveillance Methods](https://www.cdc.gov/flu/weekly/overview.htm#anchor_1633698570680) [\(/https://www.cdc.gov/flu/weekly/overview.htm#anchor_1633698570680\)](https://www.cdc.gov/flu/weekly/overview.htm#anchor_1633698570680) | [FluView Interactive](#)
[\(/https://gis.cdc.gov/grasp/fluview/mortality.html\)](https://gis.cdc.gov/grasp/fluview/mortality.html)

Influenza–Associated Pediatric Mortality

No influenza-associated pediatric deaths were reported to CDC during week 3.

A total of five influenza-associated pediatric deaths occurring during the 2021-2022 season have been reported to CDC.

[\(<http://gis.cdc.gov/GRASP/Fluview/PedFluDeath.html>\)](http://gis.cdc.gov/GRASP/Fluview/PedFluDeath.html)
[View Full Screen \(/flu/weekly/weeklyarchives2021-2022/PedFlu03.html\)](/flu/weekly/weeklyarchives2021-2022/PedFlu03.html)

Additional pediatric mortality surveillance information for current and past seasons:
[Surveillance Methods \(\[https://www.cdc.gov/flu/weekly/overview.htm#anchor_1633698596803\]\(https://www.cdc.gov/flu/weekly/overview.htm#anchor_1633698596803\)\)](https://www.cdc.gov/flu/weekly/overview.htm#anchor_1633698596803) | [FluView Interactive](#)
(<https://gis.cdc.gov/GRASP/Fluview/PedFluDeath.html>)

Additional National and International Influenza Surveillance Information


FluView Interactive: FluView includes enhanced web-based interactive applications that can provide dynamic visuals of the influenza data collected and analyzed by CDC. These [FluView Interactive applications](http://www.cdc.gov/flu/weekly/fluviewinteractive.htm) (<http://www.cdc.gov/flu/weekly/fluviewinteractive.htm>) allow people to create customized, visual interpretations of influenza data, as well as make comparisons across flu seasons, regions, age groups and a variety of other demographics.

National Institute for Occupational Safety and Health: Monthly surveillance data on the prevalence of health-related workplace absenteeism among full-time workers in the United States are [available from NIOSH](https://www.cdc.gov/niosh/topics/absences/default.html) (<https://www.cdc.gov/niosh/topics/absences/default.html>).

U.S. State and local influenza surveillance: Select a jurisdiction below to access the latest local influenza information.

Alabama (http://adph.org/influenza/)	Alaska (http://dhss.alaska.gov/dph/Epi/id/Pages/influenza/flui)
Colorado (https://www.colorado.gov/pacific/cdphe/influenza)	Connecticut (https://portal.ct.gov/DPH/Epidemiology-and-En)

Georgia (https://dph.georgia.gov/epidemiology/influenza/flu-activity-georgia)	Hawaii (http://health.hawaii.gov/docd/resources/reports/influ)
Iowa (http://idph.iowa.gov/influenza/surveillance)	Kansas (http://www.kdheks.gov/flu/surveillance.htm)
Maryland (https://phpa.health.maryland.gov/influenza/fluwatch/)	Massachusetts (https://www.mass.gov/influenza)
Missouri (http://health.mo.gov/living/healthcondiseases/communicable/influenza/reports.php)	Montana (https://dphhs.mt.gov/publichealth/cdepi/diseases/)
New Jersey (http://www.nj.gov/health/cd/topics/flu.shtml)	New Mexico (https://nmhealth.org/about/erd/ideb/isp/)
Ohio (http://www.flu.ohio.gov)	Oklahoma (https://www.ok.gov/health/Prevention_and_Preparedness/Acu)
South Carolina (http://www.scdhec.gov/Health/DiseasesandConditions/InfectiousDiseases/Flu/FluData/)	South Dakota (https://doh.sd.gov/diseases/infectious/flu/sui)
Vermont (http://www.healthvermont.gov/immunizations-infectious-disease/influenza/flu-activity-and-surveillance)	Virginia (http://www.vdh.virginia.gov/epidemiology/influenza-)
Wyoming (https://health.wyo.gov/publichealth/infectious-disease-epidemiology-unit/disease/influenza/)	New York City (http://www1.nyc.gov/site/doh/providers/hea)



World Health Organization:

Additional influenza surveillance information from participating WHO member nations is available through [FluNet](https://www.who.int/tools/flunet)  (<https://www.who.int/tools/flunet>) and the [Global Epidemiology Reports](https://www.who.int/teams/global-influenza-programme/surveillance-and-monitoring/influenza-surveillance-outputs).  (<https://www.who.int/teams/global-influenza-programme/surveillance-and-monitoring/influenza-surveillance-outputs>)

WHO Collaborating Centers for Influenza:

[Australia](http://www.influenzacentre.org/Surveillance_Samples_Received.html)  (http://www.influenzacentre.org/Surveillance_Samples_Received.html), [China](http://www.chinaivdc.cn/cnic/)  (<http://www.chinaivdc.cn/cnic/>), [Japan](http://idsc.nih.gov/jp/index.html)  (<http://idsc.nih.gov/jp/index.html>), the [United Kingdom](https://www.crick.ac.uk/research/worldwide-influenza-centre)  (<https://www.crick.ac.uk/research/worldwide-influenza-centre>), and the [United States](http://www.cdc.gov/flu/) (CDC in Atlanta, Georgia)

Europe:

The most up-to-date influenza information from Europe is available from [WHO/Europe](http://www.flunewseurope.org/) and the [European Centre for Disease Prevention and Control](http://www.flunewseurope.org/)  (<http://www.flunewseurope.org/>).

Public Health Agency of Canada:

The most up-to-date influenza information from Canada is available in [Canada’s weekly FluWatch report](http://www.phac-aspc.gc.ca/fluwatch/)  (<http://www.phac-aspc.gc.ca/fluwatch/>).

Public Health England:

The most up-to-date influenza information from the United Kingdom is available from [Public Health England](http://www.hpa.org.uk/Topics/InfectiousDiseases/InfectionsAZ/SeasonalInfluenza/)  (<http://www.hpa.org.uk/Topics/InfectiousDiseases/InfectionsAZ/SeasonalInfluenza/>).

Any links provided to non-Federal organizations are provided solely as a service to our users. These links do not constitute an endorsement of these organizations or their programs by CDC or the Federal Government, and none should be inferred. CDC is not responsible for the content of the individual organization web pages found at these links.

A description of the CDC influenza surveillance system, including methodology and detailed descriptions of each data component is available on the [surveillance methods](http://www.cdc.gov/flu/weekly/overview.htm) (<http://www.cdc.gov/flu/weekly/overview.htm>) page.