

SUPPLEMENTARY TABLE S2. MULTIVITAMIN USE PREVALENCE ESTIMATES BY PREGNANCY STATUS  
AND REGRESSION MODEL PARAMETER ESTIMATES

Variable	MV use category	Year (prevalence%)							Regression model		
		2006	2007	2008	2009	2010	2011	2012	Parameter estimate	95% CI	p
Trying to get pregnant	None	38.9	43.6	50.5	46.7	40.0	65.5	54.7	0.061	0.025 to 0.097	<0.001
	≥7 times/week	40.2	25.4	35.1	24.9	39.9	21.9	38.3	-0.031	-0.084 to 0.021	0.19
Not pregnant nor trying to get pregnant	None	45.5	46.0	53.8	49.4	52.6	56.0	57.3	0.035	0.023 to 0.047	<0.001
	≥7 times/week	31.3	31.6	27.9	32.9	29.8	23.0	21.3	-0.044	-0.063 to -0.025	<0.001
Pregnant	None	37.0	8.1	30.5	34.5	22.2	21.5	8.7	-0.091	-0.189 to 0.007	0.07
	≥7 times/week	53.8	78.2	36.1	65.0	77.9	59.2	71.0	0.031	-0.014 to 0.076	0.21
Overall	None	44.3	42.6	51.4	48.4	50.8	55.2	55.6	0.036	0.025 to 0.047	<0.001
	≥7 times/week	32.7	34.0	30.3	33.5	31.7	24.8	24.3	-0.041	-0.058 to -0.024	<0.001

Trends are only reported for the “None” and “≥7” MV use categories due to low sample size of other categories.  
MV, multivitamin.