

Supplementary Data

Systematic Review Citations

- S1. Sandoval E, Smith S, Walter J, et al. A comparison of frequent and infrequent visitors to an urban emergency department. *J Emerg Med* 2008;38:115–121.
- S2. Blank F, Li H, Henneman PL, Smithline HA, Santoro JS, Provost D, et al. A descriptive study of heavy emergency department users at an academic emergency department reveals heavy ED users have better access to care than average users. *J Emerg Nurs* 2005;31:139–144.
- S3. Rosenthal MB, Alidina S, Friedberg MW, et al. A difference-in-difference analysis of changes in quality, utilization and cost following the Colorado multi-payer patient-centered Medical home pilot. *J Gen Intern Med* 2016;31:289–296.
- S4. Simpson GG, Hahn HR, Powel AA, et al. A patient-centered emergency department management strategy for sickle-cell disease super-utilizers. *West J Emerg Med* 2017;18:335–339.
- S5. Bell J, Mancuso D, Krupski T, et al. A randomized controlled trial of king county care partners' rethinking care intervention: health and social outcomes up to two years post-randomization. Seattle, WA: Center for Healthcare Improvement for Additions, Mental Illness and Medically Vulnerable Populations, University of Washington, Harborview Medical Center, 2012.
- S6. United States Government Accountability Office. Medicaid: a small share of enrollees consistently accounted for a large share of expenditures: GA-15-460. Washington, DC: United States Government Accountability Office, 2015.
- S7. Powers, BW, Chaguturu, SK. ACOs and high-cost patients. *N Engl J Med* 2016;374:203–205.
- S8. DuBard CA, Jackson CT. Active redesign of a medicaid care management strategy for greater return on investment: predicting impactability. *Popul Health Manag* 2018;21:102–109.
- S9. Rask KJ, Williams MV, McNagny SE, et al. Ambulatory health care use by patients in a public hospital emergency department. *J Gen Intern Med* 1998;13:614–620.
- S10. Division of Healthcare Services. AMCCI provider program overview. Anchorage, AK: Alaska Department of Health and Social Services. <http://dhss.alaska.gov/dhcs/Pages/amcci/providers.aspx> Accessed November 9, 2017.
- S11. Lucas RH, Sanford SM. An analysis of frequent users of emergency care at an urban university hospital. *Ann Emerg Med* 1998;32:563–568.
- S12. Raven MC, Doran KM, Kostrowski S, Gillespie CC, Elbel BD. An intervention to improve care and reduce costs for high-risk patients with frequent hospital admissions: a pilot study. *BMC Health Serv Res* 2011; 11:270.
- S13. Ruger JP, Richter CJ, Spitznagel EL, Lewis LM. Analysis of costs, length of stay, and utilization of emergency department services by frequent users: implications for health policy. *Acad Emerg Med* 2004;11: 1311–1317.
- S14. Napoles TM, Burke NJ, Shim JK, Davis E, Moskowitz D, Yen IH. Assessing patient activation among high-need, high-cost patients in urban safety net care settings. *J Urban Health* 2017;94:803–813.
- S15. Riggs JE, Davis SM, Hobbs GR, et al. Association between early returns and frequent ED visits at a rural academic medical center. *Am J Emerg Med* 2003;21: 30–31.
- S16. Griswold SK, Nordstrom CR, Clark S, et al. Asthma exacerbations in North American adults. *Chest* 2005;127: 1579–1586.
- S17. Miller A, Cunningham M, Ali N. Bending the cost curve and improving quality of care in America's poorest city. *Popul Health Manag* 2013;16:S17–S19.
- S18. Bronsky ES, McGraw C, Johnson R, et al. Cares—a community-wide collaboration identifies super-utilizers and reduces their 9-1-1 call emergency department and hospital visit rates. *Prehosp Emerg Care* 2017;21:693–699.
- S19. Hong CS, Siegel AL, Ferris TG. Caring for high-need, high-cost patients: what makes for a successful care management program? Issue brief (Commonwealth Fund) 2014. pp. 1–19.
- S20. Hardie L, Polek C, Wheeler E, et al. Characterising emergency department high-frequency users in a rural hospital. *Emerg Med J* 2015;32: 21–25.
- S21. Milbrett P, Halm M. Characteristics and predictors of frequent utilization of emergency services. *J Emerg Nurs* 2009;35:191–198.
- S22. Jiang HJ, Weiss AJ, Barrett ML. Characteristics of emergency department visits for super-utilizers by payer, 2014: statistical brief #221. Rockville, MD: Healthcare Cost and Utilization Project (HCUP) Statistical Briefs, 2017.
- S23. Markham D, Graudins A. Characteristics of frequent emergency department presenters to an Australian emergency medicine network. *BMC Emerg Med* 2011;11:21.
- S24. Yamamoto LG, Zimmerman KR, Butts RJ, et al. Characteristics of frequent pediatric emergency department users. *Pediatr Emerg Care* 1995;11:340–346.
- S25. Hunt KA, Weber EJ, Showstack JA, et al. Characteristics of frequent users of emergency departments. *Ann Emerg Med* 2006;48:1–8.
- S26. Harris LJ, Graetz I, Podila PSB, et al. Characteristics of hospital and emergency care super-utilizers with multiple chronic conditions. *J Emerg Med* 2016;50:e203–e214.
- S27. Jiang HJ, Barrett ML, Sheng M. Characteristics of hospital stays for nonelderly medicaid super-utilizers, 2012: statistical brief #184. Rockville, MD: Healthcare Cost and Utilization Project (HCUP) Statistical Briefs, 2014.
- S28. Jiang HJ, Weiss AJ, Barrett ML, Sheng M. Characteristics of hospital stays for super-utilizers by payer, 2012: statistical brief #190. Rockville, MD: Healthcare Cost and Utilization Project (HCUP) Statistical Briefs, 2015.
- S29. Zuckerman S, Shen YC. Characteristics of occasional and frequent emergency department users: do insurance coverage and access to care matter? *Med Care* 2004; 42:176–182.
- S30. Freitag FG, Kozma CM, Slaton T, et al. Characterization and prediction of emergency department use in chronic daily headache patients. *Headache* 2005;45:891–898.

- S31. Kim H, Charlesworth C. Characterizing Oregon's Medicaid super-utilizers: descriptive analysis. Portland, OR: Oregon Health Authority, 2016.
- S32. Waxmonsky JA, Giese AA, McGinnis GF, et al. Colorado access' enhanced care management for high-cost, high-need medicaid members: preliminary outcomes and lessons learned. *J Ambul Care Manage* 2011;34:183–191.
- S33. Association of State and Territorial Health Officials. Colorado case study: super-utilizers. Arlington, VA: Association of State and Territorial Health Officials, 2016.
- S34. Perla J. Colorado super-utilizer activities: a primer. Denver, CO: Colorado Department of Health Care Policy and Financing, 2013.
- S35. Shumway M, Boccellari A, O'Brien K, et al. Cost-effectiveness of clinical case management for ED frequent users: results of a randomized trial. *Am J Emerg Med* 2008;26:155–164.
- S36. Hardin L, Kilian A, Muller L, Callison K, Olgren M. Cross-Continuum Tool Is Associated with Reduced Utilization and Cost for Frequent High-Need Users. *West J Emerg Med* 2017;18:189–200.
- S37. Moe J, Bailey AL, Oland R, Levesque L, Murray H. Defining, quantifying, and characterizing adult frequent users of a suburban Canadian emergency department. *CJEM* 2013;15:214–226.
- S38. Okuyemi KS, Frey B. Describing and predicting frequent users of an emergency department. *J Assoc Acad Minor Phys* 2001;12:1191–1123.
- S39. Huang JA, Lai CS, Tsai WC, et al. Determining factors of patient satisfaction for frequent users of emergency services in a medical center. *J Chin Med Assoc* 2004;67:403–410.
- S40. Oberste, A. Determining patient barriers affecting emergency room overutilization at Northeast Georgia Medical Center (doctoral dissertation). Atlanta, GA: Emory University, 2014.
- S41. Surbhi S, Munshi KD, Bell PC, Bailey JE. Drug therapy problems and medication discrepancies during care transitions in super-utilizers. *J Am Pharm Assoc* 2016;56:633–642.
- S42. Stamy CD, MacKinney AC. Emergency department super-utilizer programs: rural health systems analysis and technical assistance project. Iowa City, IA: Rural Health Value, 2013.
- S43. Lasser KE, Kronman AC, Cabral H, Samet JH. Emergency department use by primary care patients at a safety-net hospital. *Arch Intern Med* 2012;172:278–280.
- S44. Mandelberg JH, Kuhn RE, Kohn MA. Epidemiologic analysis of an urban, public emergency department's frequent users. *Acad Emerg Med* 2000;7:637–646.
- S45. Shah R, Chen C, O'Rourke S, et al. Evaluation of care management for the uninsured. *Med Care* 2011;49:166–171.
- S46. Huang JA, Tsai WC, Chen YC, Hu WH, Yang DY. Factors associated with frequent use of emergency services in a medical center. *J Formos Med Assoc* 2003;102:222–228.
- S47. New York State Department of Health. DSRIP—Medicaid accelerated exchange (MAX) series program: final report: improving care for super utilizers. Albany, NY: New York State Department of Health, 2017.
- S48. Phipps EJ, Singletary SB, Cooblall CA, Hares HD, Braitman LE. Food insecurity in patients with high hospital utilization. *Popul Health Manag* 2016;19:414–420.
- S49. Wright BJ, Vartanian KB, Li HF, Royal N, Matson JK. Formerly homeless people had lower overall health care expenditures after moving into supportive housing. *Health Aff* 2016;35:20–27.
- S50. Byrne M, Murphy AW, Plunkett PK, et al. Frequent attenders to an emergency department: a study of primary health care use, medical profile, and psychosocial characteristics. *Ann Emerg Med* 2003;41:309–318.
- S51. Liu SW, Nagurney JT, Chang Y, et al. Frequent ED users: are most visits for mental health, alcohol, and drug-related complaints? *Am J Emerg Med* 2013;31:1512–1515.
- S52. Kne T, Young R, Spillane L. Frequent ED users: patterns of use over time. *Am J Emerg Med* 1998;16:648–652.
- S53. Minassian A, Vilke GM, Wilson MP. Frequent emergency department visits are more prevalent in psychiatric, alcohol abuse, and dual diagnosis conditions than in chronic viral illnesses such as hepatitis and human immunodeficiency virus. *J Emerg Med* 2013;45:520–525.
- S54. Hansagi H, Olsson M, Sjöberg S, Tomson Y, Goransson S. Frequent use of the hospital emergency department is indicative of high use of other health care services. *Ann Emerg Med* 2001;37:561–567.
- S55. Doupe MB, Palatnick W, Day S, et al. Frequent users of emergency departments: developing standard definitions and defining prominent risk factors. *Ann Emerg Med* 2012;60:24–32.
- S56. Chan BT, Ovens HJ. Frequent users of emergency departments: do they also use family physicians' services? *Can Fam Physician* 2002;48:1654–1660.
- S57. Fuda KK, Immekus R. Frequent users of Massachusetts emergency departments: a statewide analysis. *Ann Emerg Med* 2006;48:9–16.
- S58. Vinton DT, Capp R, Rooks SP, Abbott JT, Ginde AA. Frequent users of US emergency departments: characteristics and opportunities for intervention. *Emerg Med J* 2014;31.
- S59. Salzberg CA, Hayes SL, McCarthy D, et al. Health system performance for the high-need patient: a look at access to care and patient care experiences. Boston, MA: Commonwealth Fund, 2016.
- S60. Lee NS, Whitman N, Vakharia N, Taksler GB, Rothberg MB. High-cost patients: hot-spotters don't explain the half of it. *J Gen Intern Med* 2017;32:28–34.
- S61. Hayes S, Salzberg C, McCarthy D, et al. High-need, high-cost patients: who are they and how do they use health care? New York, NY: The Commonwealth Fund, 3rd ed and Philadelphia, PA: Wolters Kluwer, 2016:3–16.
- S62. Belcher JV, Alexy B. High-resource hospital users in an integrated delivery system. *J Nurs Adm* 1999;29:30–36.
- S63. Peikes D, Peterson G, Brown RS, Graff S, Lynch JP. How changes in Washington University's Medicare co-ordinated care demonstration pilot ultimately achieved savings. *Health Aff* 2012;31:1216–1226.
- S64. Lynch CS, Wajnberg A, Jervis R, et al. Implementation science workshop: a novel multidisciplinary primary care program to improve care and outcomes for super-utilizers. *J Gen Intern Med* 2016;31:797–802.

- S65. Bryk J, Fischer GS, Lyons A. Improvement in quality metrics by the UPMC enhanced care program: a novel super-utilizer program. *Popul Health Manag* 2018;21: 217–221.
- S66. Koch KL, Karafin MS, Simpson P, Field JJ. Intensive management of high-utilizing adults with sickle cell disease lowers admissions. *Am J Hematol* 2015;90:215–219.
- S67. New York State Department of Health. Managing care for super utilizers. Albany, NY: New York State Department of Health, 2016.
- S68. Office of the Inspector General. MassHealth’s and the health safety net’s management of healthcare and healthcare costs for super-utilizers. Boston, MA: Office of the Inspector General, 2016.
- S69. Shane DM, Nguyen-Hoang P, Bentler SE, Damiano PC, Momany ET. Medicaid health home reducing costs and reliance on emergency department: evidence from Iowa. *Med Care* 2016;54:752–757.
- S70. Baillargeon J, Thomas CR, Williams B, et al. Medical emergency department utilization patterns among uninsured patients with psychiatric disorders. *Psychiatr Serv* 2008;59:808–811.
- S71. Zulman DM, Ezeji-Okoye SC, Shaw JG, et al. Partnered research in healthcare delivery redesign for high-need, high-cost patients: development and feasibility of an intensive management patient-aligned care team (IMPACT). *J Gen Intern Med* 2014;29 Suppl 4(Suppl 4): 861–869.
- S72. Fleming MD, Shim JK, Yen IH, et al. Patient engagement at the margins: Health care providers’ assessments of engagement and the structural determinants of health in the safety-net. *Soc Sci Med* 2017;183:11–18.
- S73. Singh K, Drouin K, Newmark LP, et al. Patient-facing mobile apps to treat high-need, high-cost populations: a scoping review. *JMIR Mhealth Uhealth* 2016;4:e136.
- S74. Ford JG, Meyer IH, Sternfels P, et al. Patterns and predictors of asthma-related emergency department use in Harlem. *Chest* 2001;120:1129–1135.
- S75. LeDuc K, Rosebrook H, Rannie M, et al. Pediatric emergency department recidivism: demographic characteristics and diagnostic predictors. *J Emerg Nurs* 2006;32:131–138.
- S76. Rhodes KV, Basseyn S, Gallop R, et al. Pennsylvania’s medical home initiative: reductions in healthcare utilization and cost among Medicaid patients with medical and psychiatric comorbidities. *J Gen Intern Med* 2016;31:1373–1381.
- S77. Szymkowiak D, Montgomery AE, Johnson EE, Manning T, O’Toole TP. Persistent super-utilization of acute care services among subgroups of veterans experiencing homelessness. *Med Care* 2017;55:893–900.
- S78. Pennsylvania Health Care Cost Containment Council. PHC4 releases recent findings on PA’s “super-utilizers” of hospital care. Harrisburg, PA: Pennsylvania Health Care Cost Containment Council, 2017.
- S79. Leininger LJ, Friedsam D, Voskuil K, et al. Predicting high-need cases among new Medicaid enrollees. *Am J Manag Care* 2014;20:e399–e407.
- S80. Sun BC, Burstin HR, Brennan TA. Predictors and outcomes of frequent emergency department users. *Acad Emerg Med* 2003; 10:320–328.
- S81. Pines JM, Buford K. Predictors of frequent emergency department utilization in southeastern Pennsylvania. *J Asthma* 2006;43:219–223.
- S82. Graven PF, Meath THA, Mendelson A, et al. Preventable acute care spending for high-cost patients across payer types. *J Health Care Finance* 2016;42.
- S83. Mulder BJ, Tzeng HM, Vecchioni ND. Preventing avoidable rehospitalizations by understanding the characteristics of “frequent fliers.” *J Nurs Care Qual* 2012; 27:77–82.
- S84. Brown KE, Levine JM, Fiellin DA, O’Connor P, Sledge WH. Primary intensive care: pilot study of a primary care-based intervention for high-utilizing patients. *Dis Manag* 2005;8:169–177.
- S85. Brown MD, Reeves MJ, Meyerson K, et al. Randomized trial of a comprehensive asthma education program after an emergency department visit. *Ann Allergy Asthma Immunol* 2006;97:44–51.
- S86. Van Der Linden MC, Van Den Brand CL, Van Der Linden N, Rambach AH, Brumsen C. Rate, characteristics, and factors associated with high emergency department utilization. *Int J Emerg Med* 2014;7:9.
- S87. Michigan Department of Community Health. Recommendations for addressing the needs of high utilizer/super utilizer patients in Michigan. Lansing, MI: Michigan Department of Community Health, 2014.
- S88. Cook LJ, Knight S, Junkins EP Jr, et al. Repeat patients to the emergency department in a statewide database. *Acad Emerg Med* 2004;11:256–263.
- S89. Washington State Health Care Authority. Report to the legislature: Emergency department utilization: Update on assumed savings from best practices implementation. Olympia, WA: Washington State Health Care Authority, 2014.
- S90. Ondler C, Hegde GG, Carlson JN. Resource utilization and health care charges associated with the most frequent ED users. *Am J Emerg Med* 2014;32:1215–1219.
- S91. McCormack RP, Hoffman LF, Wall SP, et al. Resource-limited, collaborative pilot intervention for chronically homeless, alcohol-dependent frequent emergency department users. *Am J Public Health* 2013;103(suppl): S221–S224.
- S92. Chakravarty S, Cantor JC, Walkup JT, Tong J. Role of behavioral health conditions in avoidable hospital use and cost. New Brunswick, NJ: Rutgers Center for State Health Policy, 2014.
- S93. Liptak GS, Shone LP, Auinger P, et al. Short-term persistence of high health care costs in a nationally representative sample of children. *Pediatrics* 2006;118: e1001–e1009.
- S94. Brown RS, Peikes D, Peterson G, Schore J, Razafindrakoto CM. Six features of Medicare coordinated care demonstration programs that cut hospital admissions of high-risk patients. *Health Aff* 2012;31:1156–1166.
- S95. Bieler G, Paroz S, Faouzi M, et al. Social and medical vulnerability factors of emergency department frequent users in a universal health insurance system. *Acad Emerg Med* 2012;19:63–68.
- S96. Coen SB. Statistical brief #448: differentials in the concentration of health expenditures across population subgroups in the U.S, 2012. Rockville, MD: Agency for Healthcare Research and Quality, 2014.
- S97. Bodenheimer T. Strategies to reduce costs and improve care for high-utilizing Medicaid patients: reflections on pioneering programs. San Francisco, CA: Center for Health Care Strategies, Inc., 2013.
- S98. Mann C. CMCS informational bulletin: targeting Medicaid super-utilizers to decrease costs and improve quality.

- Baltimore, MD: Centers for Medicare & Medicaid Services, 2013.
- S99. Curran GM, Sullivan G, Williams K, et al. The association of psychiatric comorbidity and use of the emergency department among persons with substance use disorders: an observational cohort study. *BMC Emerg Med* 2008;8:17.
- S100. Xu KT, Nelson BK, Berk S. The changing profile of patients who used emergency department services in the United States: 1996 to 2005. *Ann Emerg Med* 2009;54:805–810.e1–e7.
- S101. Coen SB. Statistical brief #455: the concentration of health care expenditures and related expenses for costly medical conditions, 2012. Rockville, MD: Agency for Healthcare Research and Quality, 2014.
- S102. Okin RL, Boccellari A, Azocar F, et al. The effects of clinical case management on hospital service use among ED frequent users. *Am J Emerg Med* 2000;18:603–608.
- S103. Dirmeyer VF. The frequent fliers of New Mexico: hospital readmissions among the homeless population. *Soc Work Public Health* 2016 Jul;31:288–298.
- S104. Stanton MW W, Rutherford MK. The high concentration of U.S. health care expenditures. Rockville, MD: Agency for Healthcare Research and Quality, 2005.
- S105. Durfee J, Johnson T, Batal H. The impact of tailored intervention services on charges and mortality for adult super-utilizers. *Healthc (Amst)* 2018;6:253–258.
- S106. Jackson TJ, Trygstad TK, DeWalt DA, DuBard AC. Transitional care cut hospital readmissions for North Carolina Medicaid patients with complex chronic conditions. *Health Affairs* 2013;32:1407–1415.
- S107. Friedman BW, Serrano D, Reed M, et al. Use of the emergency department for severe headache. A population-based study. *Headache* 2009;49:21–30.
- S108. Isasi F. Using data to better serve the most complex patients: highlights from NGA's intensive work with seven states. Washington, DC: National Governor's Association, 2016.
- S109. Huang A. Using SAS programing to identify super-utilizers and improve healthcare services. Cary, NC: The Proceedings of the SouthEast SAS Users Group, 2015.
- S110. Warning W, Wood J, Letcher A, et al. Working with the super-utilizer population: the experience and recommendations of Pennsylvania programs. Camden, NJ: South Central Pennsylvania High Utilizer Collaborative, 2014.
- S111. Johansson S, Fuchs F. Wyoming super-utilizer program (WySUP). Cheyenne, WY: Wyoming Department of Health, 2016.