

Morbidity and Mortality

Weekly
Report

PUBLIC HEALTH SERVICE

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE

Prepared by the NATIONAL OFFICE OF VITAL STATISTICS Executive 3-6300, Ext. 4744

October 30, 1959

Washington 25, D. C.

Vol. 7, No. 54

ANNUAL SUPPLEMENT

Reported Incidence of Notifiable Diseases in the United States, 1958

This supplement contains final figures on the reported incidence of notifiable diseases for 1958. The data are compiled from annual reports supplied by the State and Territorial health departments.

Trends in the nationwide incidence of notifiable diseases during the last 10 years are shown in table 1. In table 2, numbers of cases are distributed by month, usually by month of report; in table 3, age distributions for 7 diseases are shown. In the other tables, distributions of cases for States and outlying areas and the numbers of cases of certain diseases not regularly reported are shown. The numbers of cases of 4 diseases reported in 1953, 1957, and 1958 appear in figure 1.


For some of the diseases, there was a substantial decrease in the figures reported for 1958 compared with 1957. These include amebiasis, anthrax, botulism, diphtheria, malaria,

nonparalytic poliomyelitis, psittacosis, typhoid fever, endemic typhus fever, and syphilis. The decrease in the number of cases of anthrax was accompanied by a change in the usual geographic distribution of the disease, with more cases reported in the West South Central States than in the New England and Middle Atlantic States. An anthrax vaccine evaluation program underway in industry may account for the decrease in cases reported in the latter areas. The reported incidence of malaria continues to decline steadily. Practically all the cases have origin in areas outside the United States, although this information is not shown in reports from a number of States.

Among the diseases for which the incidence in 1958 remained about the same as in 1957 are brucellosis, meningococcal infections, rheumatic fever, Rocky Mountain spotted fever, salmonellosis, tetanus, trichinosis, and tuberculosis.

Figure 1

REPORTED CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES, 1953, 1957, AND 1958



COMMUNICABLE DISEASE CENTER
LIBRARY
ATLANTA 22, GEORGIA

Morbidity and Mortality Weekly Report


Substantial increases were reported for encephalitis, infectious hepatitis, shigellosis, measles, other meningitis, paralytic poliomyelitis, pertussis, scarlet fever, and gonorrhoea. A number of States are now reporting cases of aseptic meningitis, and these are included in the category "Meningitis, other." This probably accounts for the larger total reported in 1958 for this category. The total of 763,094 cases of measles reported in 1958 is the largest figure since 1941. Since the disease was epidemic in States with large populations, the total number of cases is not unexpectedly high. However, it is probable that only a small fraction of the cases are reported.

Poliomyelitis

The total number of cases of all forms of poliomyelitis reported in 1958 was 5,787, an increase of about 5 percent over the 5,485 reported in 1957. However, the increase in

Figure 2

**PARALYTIC POLIOMYELITIS — PERCENT OF CASES
BY AGE: UNITED STATES, 1956-58**



paralytic cases was almost 50 percent—3,697 cases in 1958 compared with 2,499 in 1957. This increase interrupted the downward trend in cases since 1954. About 64 percent of the total cases were classified as paralytic in final figures for 1958, whereas in 1957 the percentage was 46. Also there was some change between the final figures and the provisional figures derived from weekly morbidity reports made during the year. They compare as follows:

	Provisional	Final
Total-----	6,061	5,787
Paralytic-----	3,140	3,697
Nonparalytic-----	2,023	1,941
Unspecified-----	898	149

These figures indicate there was considerable followup of reported cases. A large number of cases initially reported as unspecified apparently were changed to the paralytic classification. The smaller number of nonparalytic cases on final analysis indicates the greater use of laboratory facilities and a reluctance to report such cases without laboratory confirmation.

Twenty-seven States reported more cases of paralytic poliomyelitis in 1958 than in 1957, and paralytic case totals for all geographic divisions except the Pacific Division were larger in 1958. The States reporting an increase of 50 percent or more in 1958 were Rhode Island (no cases in 1957), Connecticut, New York, New Jersey, Pennsylvania, Ohio, Michigan, Iowa, Missouri, North Dakota, Delaware, Virginia, West Virginia, Florida, Montana, Nevada, and Washington.

The age distribution of the paralytic cases shows that about 50 percent of the cases occurred in children under 5 years of age, with the largest number among 3-year-old children. Figure 2 shows that since 1956, when the age distribution was first tabulated, there has been a relative increase in the number of cases in preschool children, whereas the number of cases in older school age groups has remained the same or decreased. This probably reflects the effects of vaccination programs in which school children were the first to receive the vaccine. The ratio of paralytic cases to total cases was highest in 1-year-old children, 88 percent of all cases in these children being paralytic.


There were several localized epidemics reported during 1958. The largest occurred in Detroit, Michigan, chiefly among children in the lower socioeconomic groups living in slum areas where few were vaccinated. Localizations also occurred in New Jersey, some Texas cities, contiguous areas of Virginia and West Virginia, an Indian reservation in Montana, and in Hawaii, mainly among dependents of military personnel.

Infectious hepatitis

In 1958 there was an increase of about 9 percent in the number of reported cases of infectious and serum hepatitis over the 1957 figure—16,294 cases in 1958 compared with 14,922 in 1957. However, the number of cases reported during the last 6 months of 1958 is about 32 percent greater than the number reported during the comparable period of 1957. Much of the increase occurred in the Mountain Division, where all the States except Wyoming and New Mexico reported more cases in 1958. In Arizona, there was an outbreak of more than 300 cases on an Indian reservation reported early in the year. The number of cases and the number of deaths reported since 1952 is shown in figure 3. The number of cases has decreased greatly from the peak in 1954, but the number of deaths has remained relatively constant, with a slight increase in 1957. The number of deaths for 1958 is not available.

Figure 3

INFECTIOUS HEPATITIS, UNITED STATES: REPORTED CASES, 1952-58, AND REGISTERED DEATHS, 1952-57



Diphtheria

For the first time since records have been kept, the national total number of cases of diphtheria has dropped below 1,000. The total, 918, is 24 percent less than the 1957 figure of 1,211. The age distribution of cases was similar to that reported in 1956 and 1957. Two-thirds of the cases were reported in the South Atlantic and South Central States. Localized outbreaks were reported in Arkansas, Louisiana, Minnesota, and Mississippi.

Encephalitis

The number of reported cases of encephalitis in 1958 was about 21 percent higher than the number reported in 1957. Of the 2,587 cases, 344 were reported as arthropod-borne encephalitis. These were reported by 14 States, 11 of which were in the western half of the country. In the central and western parts of Kansas, there was an outbreak of 83 cases during August, September, and October. Both St. Louis and western equine encephalitis were confirmed in this outbreak, but the former was predominant. In California, the predominant type was western equine encephalitis. In the eastern half of the country, Alabama, Florida, and Illinois listed a number of arthropod-borne cases on their annual reports.

Typhoid fever

The number of cases of typhoid fever (1,043) dropped 15 percent from the 1,231 cases reported in 1957 and about 40 percent from the 1,700 in 1956. Almost 20 percent of the cases were reported in California and Texas. Several outbreaks were

reported, but only one, consisting of 30 cases, was reported as due to a food source.

Venereal diseases

Although there was a decrease of 15 percent in reported cases of syphilis compared with 1957, there was an increase of about 9 percent in the number of cases of gonorrhea. Consequently, the total number of cases of venereal diseases remained about the same as in 1957. The 1958 figure for gonorrhea (241,792) also exceeds that of 1956 and is comparable with the 1955 figure. Except for 1958 and 1954, there has been a steady decline in the incidence of gonorrhea since 1946, when more than 400,000 cases were reported.

The data in this report include cases on military reservations. Consequently, this data will vary in many instances from reports of the Venereal Disease Branch of the Communicable Disease Center, which do not include cases among military personnel.

EXPLANATION OF SYMBOLS USED IN TABLES

- Quantity zero ----- -
- Disease stated not notifiable ----- *
- No report made by State ----- **
- Data not available ----- ---

Table 1. REPORTED CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES, 1949-58

(Numbers after diseases are category numbers of the Seventh Revision of the International Lists, 1955)

DISEASE	1958	1957	1956	1955	1954	1953	1952	1951	1950	1949
Amebiasis-----046	4,380	5,031	3,689	3,348	3,523	4,444	4,280	3,550	4,568	5,543
Anthrax-----062	16	26	38	39	22	45	47	60	49	54
Botulism-----049.1	6	28	17	16	18	18	18	33	20	24
Brucellosis (undulant fever)-----044	924	983	1,300	1,444	1,823	2,032	2,537	3,139	3,510	4,235
Dengue-----090	-	-	2	1	6	8	5	16	26	46
Diphtheria-----055	918	1,211	1,568	1,984	2,041	2,355	2,960	3,983	5,796	7,969
Dysentery, bacillary (shigellosis)-----045	11,861	9,822	10,306	13,912	13,846	16,533	23,197	32,215	23,367	29,080
Encephalitis, acute infectious-----082	2,587	2,135	2,624	2,166	2,606	1,935	1,912	1,123	1,135	903
Hepatitis, infectious, and serum-----092, N998.5 pt.	16,294	14,922	19,234	31,961	50,093	33,700	17,428	7,349	2,820	2,027
Leprosy-----060	39	36	52	75	56	60	57	57	44	41
Leptospirosis-----072	55	47	44	24	48	42	62	9	30	17
Malaria-----110-117	85	132	234	522	715	1,310	7,023	5,600	2,184	4,151
Measles-----085	763,094	486,799	611,936	555,156	682,720	449,146	683,077	530,118	319,124	625,281
Meningococcal infections-----057	2,581	2,691	2,735	3,455	4,436	5,077	4,884	4,164	3,788	3,519
Plague-----058	-	-	1	-	-	-	-	1	3	3
Poliomyelitis, acute-----080	5,787	5,485	15,140	28,985	38,476	35,592	57,879	28,386	33,300	42,033
Paralytic-----080.0, 080.1	3,697	2,499	7,911	13,850	18,308	15,648	21,269	10,037	---	---
Nonparalytic-----080.2	1,941	2,826	6,555	12,453	13,221	12,144	12,802	5,470	---	---
Unspecified-----080.3	149	160	674	2,682	6,947	7,800	23,808	12,879	---	---
Psittacosis-----096.2	158	278	568	334	563	169	135	25	26	35
Rabies in man ¹ -----094	5	5	10	4	13	12	24	18	18	10
Rabies in animals ² -----	4,787	4,542	5,681	5,799	7,297	8,903	8,445	8,008	7,901	7,587
Rocky Mountain spotted fever-----104 pt.	243	240	293	295	294	313	327	347	464	570
Salmonellosis, except typhoid fever-----041, 042	6,363	6,693	6,704	5,447	5,375	3,946	2,596	1,773	1,233	1,243
Scarlet fever and streptococcal sore throat-----050, 051	264,097	226,973	176,392	147,502	147,785	132,935	113,677	84,151	64,494	87,220
Smallpox-----084	-	91	-	92	99	4	21	11	39	49
Tetanus-----061	445	447	468	462	524	506	484	506	486	579
Trichinosis-----128	176	178	262	264	277	395	367	393	327	353
Tuberculosis, all forms-----001-019	82,266	86,861	90,465	98,860	100,540	106,925	109,837	118,491	121,742	134,865
Tularemia-----059	587	601	522	584	681	601	668	702	927	1,179
Typhoid fever-----040	1,043	1,231	1,700	1,704	2,169	2,252	2,341	2,128	2,484	2,795
Typhus fever, endemic (murine)-----101	71	113	98	135	163	221	205	378	685	985
Venereal diseases: ⁴										
Gonorrhea-----030-034	241,792	222,048	233,593	244,279	249,883	246,311	253,839	254,057	286,746	317,950
Syphilis-----020-029	115,173	136,039	131,763	123,004	131,260	150,026	169,198	174,924	217,558	256,463
Other specified venereal diseases-----036-038	2,454	2,519	3,147	4,078	4,650	5,209	6,093	6,885	8,187	11,034
Whooping cough (pertussis)-----056	32,148	28,295	31,732	62,786	60,886	37,129	45,030	68,687	120,718	69,479

¹For 1949-57, figures represent registered deaths.

²For 1949-51, figures from Agricultural Research Service, U.S. Department of Agriculture.

³These cases do not fulfill the generally accepted criteria for a diagnosis of smallpox.

⁴For 1949-51, figures (civilian cases only) from the Venereal Disease Program, Public Health Service.

Table 2. REPORTED CASES OF SPECIFIED NOTIFIABLE DISEASES BY MONTH: UNITED STATES, 1958

(Numbers after diseases are category numbers of the Seventh Revision of the International Lists, 1955)

DISEASE	Total ¹	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
Amebiasis-----046	4,380	412	376	357	420	404	327	326	423	350	389	259	325
Anthrax-----062	16	-	-	1	-	1	2	5	3	-	-	3	1
Botulism-----049.1	6	-	-	-	-	3	1	-	-	1	1	-	-
Brucellosis (undulant fever)-----044	924	46	52	53	62	86	66	91	70	69	59	43	60
Cholera-----043	-	-	-	-	-	-	-	-	-	-	-	-	-
Dengue-----090	-	-	-	-	-	-	-	-	-	-	-	-	-
Diphtheria-----055	918	79	52	69	44	51	25	36	52	82	136	152	140
Dysentery, bacillary (shigellosis)-----045	11,861	619	678	591	851	1,170	915	817	1,092	1,260	1,402	1,411	1,051
Encephalitis, acute infectious-----082	2,587	86	91	167	196	208	220	286	357	417	246	168	144
Arthropod-borne-----082.0	344	5	5	10	5	12	23	53	49	96	44	32	10
Glanders-----064.2	1	-	-	-	-	-	1	-	-	-	-	-	-
Hepatitis, infectious, and serum-----092,N998.5 pt.	16,294	1,620	1,375	1,511	1,516	1,383	1,051	1,103	1,112	1,244	1,339	1,255	1,770
Leprosy-----080	39	-	4	1	2	12	4	1	2	4	5	3	1
Leptospirosis-----072	55	3	1	3	-	1	3	7	6	3	8	5	3
Malaria-----110-117	85	5	5	-	7	9	12	11	13	6	3	6	5
Measles-----085	763,094	39,963	65,387	128,714	166,965	170,039	96,707	33,915	10,284	4,750	8,412	13,520	22,818
Meningococcal infections-----057	2,581	267	276	303	219	208	144	163	155	204	190	192	258
Meningitis, other-----340	5,149	270	254	280	232	229	264	451	919	890	575	410	375
Plague-----058	-	-	-	-	-	-	-	-	-	-	-	-	-
Poliomyelitis, acute-----080	5,787	63	58	51	60	139	231	572	1,309	1,505	1,000	502	295
Paralytic-----080.0,080.1	3,697	42	38	34	43	78	143	327	761	907	686	399	238
Nonparalytic-----080.2	1,941	16	15	16	13	49	80	218	531	578	283	90	51
Unspecified-----080.3	149	5	5	1	4	12	8	27	17	20	31	13	6
Psittacosis-----096.2	158	9	13	20	14	10	15	17	21	10	9	5	10
Rabies in man-----094	5	-	1	-	-	-	-	-	-	-	3	-	1
Rabies in animals-----	4,787	437	383	534	559	381	395	469	355	320	333	269	344
Relapsing fever (louse-borne)-----071.0	-	-	-	-	-	-	-	-	-	-	-	-	-
Rheumatic fever-----400-402	6,889	562	619	640	695	575	573	604	445	513	586	470	586
Rocky Mountain spotted fever-----104 pt.	243	1	1	-	8	30	65	52	54	21	5	3	3
Salmonellosis, except typhoid fever-----041,042	6,363	475	307	447	495	486	493	663	599	514	613	601	656
Scarlet fever and streptococcal sore throat-----050,051	264,097	21,802	25,758	36,838	33,771	28,941	17,686	11,980	13,103	12,866	17,850	20,648	22,587
Smallpox-----084	-	-	-	-	-	-	-	-	-	-	-	-	-
Tetanus-----061	445	20	29	25	27	34	43	45	50	64	40	32	36
Trichinosis-----128	176	11	14	12	6	11	27	13	22	7	18	21	14
Tuberculosis, all forms-----001-019	82,266	-	-	-	-	-	-	-	-	-	-	-	-
Tularemia-----059	587	55	43	32	30	48	65	57	54	32	40	41	89
Typhoid fever-----040	1,043	48	51	55	66	78	63	117	166	133	127	68	63
Typhus fever, endemic (murine)-----101	71	4	3	3	3	5	14	10	7	11	2	2	7
Typhus fever, epidemic-----100	-	-	-	-	-	-	-	-	-	-	-	-	-
Veneral diseases-----020-029,030-034,036-038	359,419	-	-	-	-	-	-	-	-	-	-	-	-
Whooping cough (pertussis)-----056	32,148	1,723	1,580	2,005	2,549	2,980	2,961	3,248	2,844	3,021	2,723	3,022	3,259
Yellow fever-----091	-	-	-	-	-	-	-	-	-	-	-	-	-

¹Cases not tabulated by month included in "Total" but not distributed.

Morbidity and Mortality Weekly Report

Table 3. REPORTED CASES OF SPECIFIED NOTIFIABLE DISEASES BY AGE: TOTAL OF REPORTING STATES, 1958

(Numbers after diseases are category numbers of the Seventh Revision of the International Lists, 1955)

DISEASE	Number of States ¹	Total cases	Under 1 year	1-4 years	5-9 years	10-14 years
Diphtheria-----055	46	909	17	246	260	131
Encephalitis, acute infectious-----082	42	2,334	117	414	605	298
Arthropod-borne-----082.0	41	344	31	40	63	36
Hepatitis, infectious, and serum-----092,N998.5 pt.	38	12,835	48	518	2,318	1,831
Meningococcal infections-----057	38	2,063	422	737	277	138
Meningitis, other-----340	38	4,621	795	977	850	496
Poliomyelitis, acute-----080	46	5,487	282	1,856	1,158	542
Paralytic-----080.0,080.1	46	3,494	239	1,468	634	233
Nonparalytic-----080.2	46	1,845	34	341	495	298
Unspecified-----080.3	45	148	9	47	29	11
Typhoid fever-----040	44	999	13	122	197	135

DISEASE	15-19 years	20-24 years	25-34 years	35-44 years	45 years and over	Not stated
Diphtheria-----055	74	33	35	34	39	40
Encephalitis, acute infectious-----082	120	106	218	161	235	60
Arthropod-borne-----082.0	21	13	27	31	77	5
Hepatitis, infectious, and serum-----092,N998.5 pt.	1,253	1,092	1,711	1,102	1,426	1,536
Meningococcal infections-----057	109	37	56	41	167	79
Meningitis, other-----340	274	227	437	178	306	81
Poliomyelitis, acute-----080	397	384	575	204	75	14
Paralytic-----080.0,080.1	209	198	333	130	44	6
Nonparalytic-----080.2	176	172	228	72	22	7
Unspecified-----080.3	12	14	14	2	9	1
Typhoid fever-----040	78	57	94	85	177	43

¹The District of Columbia is not included in "Number of States" but is represented in data shown for each disease.

Morbidity and Mortality Weekly Report

Table 4. REPORTED CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES, EACH DIVISION AND STATE, 1958
(Numbers under diseases are category numbers of the Seventh Revision of the International Lists, 1955)

AREA	Amebiasis 046	Brucel- losis (undulant fever) 044	Diph- theria 055	Dysentery, bacillary (shigel- losis) 045	ENCEPHALITIS, ACUTE INFECTIOUS		Hepatitis, infectious, and serum 092, N988.5 pt.	MALARIA 110-117	
					Total 082	Arthro- pod-borne 082.0		Total ¹	Acquired outside United States
UNITED STATES-----	4,380	924	918	11,861	2,587	344	16,294	85	37
NEW ENGLAND-----	94	16	7	202	118	-	636	2	-
Maine-----	-	1	-	1	6	-	87	-	-
New Hampshire-----	1	-	-	1	2	-	8	-	-
Vermont-----	-	5	-	-	-	-	35	-	-
Massachusetts-----	5	7	6	133	60	-	304	-	-
Rhode Island-----	24	1	-	-	42	-	80	-	-
Connecticut-----	64	2	1	67	8	-	122	2	-
MIDDLE ATLANTIC-----	1,671	28	34	311	313	-	2,278	10	-
New York-----	1,169	13	12	202	228	-	1,564	8	-
New Jersey-----	315	4	5	14	23	-	193	1	-
Pennsylvania-----	187	11	17	95	62	-	521	1	-
EAST NORTH CENTRAL-----	383	133	46	1,471	540	88	2,802	11	-
Ohio-----	13	6	14	399	85	-	844	4	-
Indiana-----	14	8	14	35	126	-	313	4	-
Illinois-----	227	83	12	193	218	88	706	3	-
Michigan-----	128	24	6	735	91	-	688	-	-
Wisconsin-----	1	12	-	109	20	-	251	-	-
WEST NORTH CENTRAL-----	172	501	135	537	253	93	1,359	1	-
Minnesota-----	2	38	75	265	12	7	213	-	-
Iowa-----	78	283	12	37	19	-	201	-	-
Missouri-----	8	41	12	163	7	1	281	-	-
North Dakota-----	-	10	3	14	34	-	267	-	-
South Dakota-----	-	23	21	8	9	-	28	-	-
Nebraska-----	1	35	10	5	27	2	96	1	-
Kansas-----	83	71	2	45	145	83	273	-	-
SOUTH ATLANTIC-----	366	62	283	816	284	24	1,249	15	6
Delaware-----	-	-	-	4	2	-	62	-	-
Maryland-----	8	1	3	82	92	-	201	-	-
District of Columbia-----	2	-	2	64	9	-	25	-	-
Virginia-----	54	21	33	130	25	-	286	4	-
West Virginia-----	1	-	10	5	5	-	168	-	-
North Carolina-----	18	7	32	83	53	-	75	6	6
South Carolina-----	7	3	39	-	35	-	42	-	-
Georgia-----	19	25	101	324	39	-	148	-	-
Florida-----	257	5	63	124	24	24	242	5	-
EAST SOUTH CENTRAL-----	181	41	104	173	89	17	1,315	8	6
Kentucky-----	33	12	7	-	18	-	668	3	3
Tennessee-----	60	19	12	87	23	-	331	3	3
Alabama-----	25	2	53	15	17	17	226	-	-
Mississippi-----	63	8	32	71	31	-	90	2	-
WEST SOUTH CENTRAL-----	308	69	229	5,386	205	46	1,127	6	1
Arkansas-----	23	19	48	81	18	-	105	1	-
Louisiana-----	102	15	89	86	2	*	26	-	-
Oklahoma-----	38	5	28	58	37	-	163	5	1
Texas-----	145	30	64	5,161	148	46	833	-	-
MOUNTAIN-----	185	28	64	513	213	16	2,456	8	-
Montana-----	-	3	16	12	9	-	424	-	-
Idaho-----	8	1	1	41	4	-	219	1	-
Wyoming-----	-	2	2	-	7	1	15	-	-
Colorado-----	18	5	14	238	113	8	327	-	-
New Mexico-----	51	3	27	119	41	-	365	-	-
Arizona-----	104	2	4	95	5	-	814	6	-
Utah-----	3	10	-	7	26	7	183	-	-
Nevada-----	1	2	-	1	8	-	109	1	-
PACIFIC-----	1,020	46	16	2,452	572	60	3,072	24	24
Washington-----	6	3	3	411	21	2	513	-	-
Oregon-----	25	8	7	78	26	5	455	-	-
California-----	989	35	6	1,963	525	53	2,104	24	24

¹Includes cases with source of contraction not stated.

Morbidity and Mortality Weekly Report

Table 4. REPORTED CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES, EACH DIVISION AND STATE, 1958—Continued

(Numbers under diseases are category numbers of the Seventh Revision of the International Lists, 1955)

AREA	Measles 085	Meningo- coccal infec- tions 057	Menin- gitis, other 340	POLIOMYELITIS, ACUTE			Psitta- cosis 096.2	Rabies in man 094	
				Total	Paralytic	Non- paralytic			Unspeci- fied
				080	080.0, 080.1	080.2			080.3
UNITED STATES-----	763,094	2,581	5,149	5,787	3,697	1,941	149	158	5
NEW ENGLAND-----	75,072	146	232	90	56	34	-	8	-
Maine-----	4,476	20	37	4	4	-	-	2	-
New Hampshire-----	4,711	13	6	6	1	5	-	1	-
Vermont-----	2,582	3	35	6	5	1	-	1	-
Massachusetts-----	39,047	59	99	29	14	15	-	3	-
Rhode Island-----	7,468	9	55	3	3	-	-	-	-
Connecticut-----	16,988	42	-	42	29	13	-	1	-
MIDDLE ATLANTIC-----	131,452	329	412	684	507	177	-	29	-
New York-----	64,350	139	412	282	210	72	-	18	-
New Jersey-----	27,546	69	-	266	186	80	-	2	-
Pennsylvania-----	39,556	121	-	136	111	25	-	9	-
EAST NORTH CENTRAL-----	189,748	633	1,323	1,996	1,107	887	2	37	2
Ohio-----	31,444	123	301	368	276	92	-	3	2
Indiana-----	19,510	56	152	139	102	37	-	-	-
Illinois-----	24,413	175	734	204	134	70	-	7	-
Michigan-----	41,448	228	91	1,228	557	671	-	5	-
Wisconsin-----	72,933	51	45	57	38	17	2	22	-
WEST NORTH CENTRAL-----	24,264	166	292	428	284	143	1	33	1
Minnesota-----	1,346	36	133	26	23	3	-	22	-
Iowa-----	12,723	15	60	73	35	38	-	6	-
Missouri-----	3,640	49	25	195	157	38	-	2	-
North Dakota-----	4,734	29	-	51	34	17	-	-	-
South Dakota-----	302	-	8	15	10	5	-	-	1
Nebraska-----	31,519	24	19	34	3	30	1	2	-
Kansas-----	*	13	47	34	22	12	-	1	-
SOUTH ATLANTIC-----	80,684	436	685	870	563	215	92	12	1
Delaware-----	567	5	7	29	18	11	-	-	-
Maryland-----	6,391	33	88	28	23	5	-	1	-
District of Columbia-----	1,496	21	108	10	8	2	-	1	-
Virginia-----	22,420	98	272	159	140	18	1	4	-
West Virginia-----	13,066	37	34	216	159	56	1	-	-
North Carolina-----	4,590	90	1	74	37	37	-	2	-
South Carolina-----	12,600	26	27	32	25	6	1	1	-
Georgia-----	7,040	31	148	70	56	14	-	2	1
Florida-----	12,514	95	-	252	97	66	89	1	-
EAST SOUTH CENTRAL-----	60,229	296	343	380	240	126	14	5	-
Kentucky-----	20,530	55	66	88	70	18	-	-	-
Tennessee-----	29,748	67	170	119	76	43	-	4	-
Alabama-----	8,402	115	-	60	52	8	-	1	-
Mississippi-----	1,549	59	107	113	42	57	14	-	-
WEST SOUTH CENTRAL-----	97,307	240	359	791	538	252	1	2	-
Arkansas-----	3,404	19	32	33	31	2	-	-	-
Louisiana-----	317	99	*	78	58	20	*	-	-
Oklahoma-----	7,500	38	112	61	26	34	1	-	-
Texas-----	86,086	84	215	619	423	196	-	2	-
MOUNTAIN-----	46,927	81	280	203	111	53	39	5	-
Montana-----	8,250	8	17	68	49	18	1	-	-
Idaho-----	4,138	3	12	13	3	4	6	2	-
Wyoming-----	1,383	5	95	13	5	2	6	1	-
Colorado-----	9,867	20	75	18	13	5	-	-	-
New Mexico-----	8,764	12	27	39	15	5	19	1	-
Arizona-----	10,400	21	30	34	17	16	1	-	-
Utah-----	3,921	7	18	12	5	2	5	1	-
Nevada-----	204	5	6	6	4	1	1	-	-
PACIFIC-----	57,411	254	1,223	345	291	54	-	27	1
Washington-----	11,428	36	105	40	32	8	-	1	-
Oregon-----	9,752	22	91	40	27	13	-	9	-
California-----	36,231	196	1,027	265	232	33	-	17	1

²Not notifiable in New York City.³Includes rubella (German measles) (086).

Morbidity and Mortality Weekly Report

9

Table 4. REPORTED CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES, EACH DIVISION AND STATE, 1958—Continued

(Numbers under diseases are category numbers of the Seventh Revision of the International Lists, 1955)

AREA	Rabies in animals	Rheumatic fever 400-402	Rocky Mountain spotted fever 104 pt.	Salmonel- losis, except typhoid fever 041,042	Scarlet fever and strepto- coccal sore throat 050,051	Tetanus 061	Trichino- sis 128	Tuber- culosis, all forms 001-019	Tularemia 059
UNITED STATES-----	4,787	6,889	243	6,363	264,097	445	176	82,266	587
NEW ENGLAND-----	1	35	3	775	10,473	11	22	2,915	2
Maine-----	-	3	-	20	826	3	1	211	-
New Hampshire-----	-	6	-	3	711	-	-	85	-
Vermont-----	-	-	-	5	539	-	-	153	-
Massachusetts-----	-	-	3	613	3,687	3	12	1,505	2
Rhode Island-----	-	24	-	31	1,618	2	1	278	-
Connecticut-----	1	*	-	103	3,092	3	8	683	-
MIDDLE ATLANTIC-----	324	-	26	747	19,516	39	91	17,978	5
New York-----	261	*	7	⁴ 399	10,044	16	55	8,073	-
New Jersey-----	-	-	9	52	5,225	5	11	2,790	-
Pennsylvania-----	63	-	10	296	4,247	18	25	7,115	5
EAST NORTH CENTRAL-----	790	4,441	27	1,140	31,119	57	17	19,256	117
Ohio-----	163	2,100	8	153	8,931	12	14	4,410	1
Indiana-----	332	63	10	68	4,528	13	-	1,754	51
Illinois-----	32	894	9	459	4,501	24	3	6,297	61
Michigan-----	38	1,320	-	335	5,380	4	-	5,587	1
Wisconsin-----	225	64	-	125	7,779	4	-	1,208	3
WEST NORTH CENTRAL-----	1,071	200	8	519	7,006	24	21	4,165	76
Minnesota-----	461	96	-	225	1,369	8	10	1,151	1
Iowa-----	230	15	-	68	2,030	2	-	539	2
Missouri-----	165	49	3	99	743	9	-	1,534	36
North Dakota-----	74	39	2	44	2,009	-	11	91	-
South Dakota-----	75	-	3	11	203	-	-	164	-
Nebraska-----	55	-	-	17	72	-	-	212	4
Kansas-----	11	1	-	55	580	5	-	474	33
SOUTH ATLANTIC-----	869	325	108	913	28,925	93	5	12,156	90
Delaware-----	-	11	3	16	113	-	-	173	-
Maryland-----	8	-	15	61	612	5	2	1,583	5
District of Columbia-----	-	1	-	38	143	-	-	1,053	-
Virginia-----	252	-	32	126	7,105	9	-	1,851	36
West Virginia-----	121	57	3	2	3,368	4	3	944	-
North Carolina-----	39	-	25	213	1,406	9	-	1,635	12
South Carolina-----	119	195	4	8	4,865	3	-	990	6
Georgia-----	274	61	26	244	8,574	23	-	1,701	27
Florida-----	56	-	-	205	2,739	40	-	2,226	4
EAST SOUTH CENTRAL-----	836	521	33	128	43,727	85	5	7,186	96
Kentucky-----	463	127	7	34	9,090	5	-	1,575	48
Tennessee-----	131	123	21	44	26,852	26	5	2,566	36
Alabama-----	223	246	4	28	5,101	29	-	2,068	6
Mississippi-----	19	25	1	22	2,684	25	-	977	6
WEST SOUTH CENTRAL-----	678	212	13	503	45,321	85	2	7,345	132
Arkansas-----	135	13	11	65	683	16	-	1,151	80
Louisiana-----	31	99	-	226	174	31	1	1,357	28
Oklahoma-----	29	36	2	69	6,825	6	1	963	17
Texas-----	483	64	-	143	37,639	32	-	3,874	7
MOUNTAIN-----	46	682	22	269	51,956	8	5	3,151	52
Montana-----	-	77	1	8	2,134	-	-	294	-
Idaho-----	-	31	2	3	2,496	3	-	100	6
Wyoming-----	-	277	5	-	2,485	-	-	123	24
Colorado-----	7	135	12	145	13,552	2	-	863	4
New Mexico-----	12	47	1	⁵ 17	11,527	-	-	670	1
Arizona-----	27	94	-	35	13,369	3	4	852	-
Utah-----	-	21	-	57	6,337	-	1	127	17
Nevada-----	-	-	1	4	56	-	-	122	-
PACIFIC-----	172	475	3	1,369	26,054	43	8	8,114	17
Washington-----	-	271	-	169	⁶ 8,735	1	-	1,219	4
Oregon-----	-	59	3	58	2,274	-	2	572	7
California-----	172	145	-	1,142	15,045	42	6	6,323	6

⁴Only paratyphoid fever (041) notifiable in New York City.

⁶Includes erysipelas (052).

⁵Excludes paratyphoid fever (041).

Morbidity and Mortality Weekly Report

Table 4. REPORTED CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES, EACH DIVISION AND STATE, 1958--Continued

(Numbers under diseases are category numbers of the Seventh Revision of the International Lists, 1955)

AREA	Typhoid fever 040	Typhus fever, endemic (murine) 101	VENEREAL DISEASES						Whooping cough (pertussis) 056
			Chancroid 036	Gonorrhea 030-034	Granuloma inguinale 038	Lympho-granuloma venereum 037	Syphilis		
							Total 020-029	Primary, secondary, ⁷ 021,028.0	
UNITED STATES-----	1,043	71	1,682	241,792	322	450	115,173	23,949	32,148
NEW ENGLAND-----	18	1	9	4,059	2	5	3,346	829	2,931
Maine-----	1	-	-	97	-	-	45	9	503
New Hampshire-----	2	-	-	42	-	1	84	9	239
Vermont-----	-	-	-	106	-	-	14	5	233
Massachusetts-----	8	-	4	2,069	1	2	2,211	535	1,270
Rhode Island-----	1	-	3	321	-	1	371	53	185
Connecticut-----	6	1	2	1,424	1	1	621	218	501
MIDDLE ATLANTIC-----	104	-	160	28,511	25	44	26,846	4,621	3,320
New York-----	37	-	^a 126	17,021	10	^a 37	15,031	2,725	1,659
New Jersey-----	23	-	13	5,852	6	5	6,077	826	239
Pennsylvania-----	44	-	21	5,638	9	2	5,738	1,070	1,422
EAST NORTH CENTRAL-----	111	1	251	41,401	12	24	19,904	3,764	6,084
Ohio-----	40	-	25	7,892	1	8	5,840	1,067	1,383
Indiana-----	21	-	5	2,222	1	7	1,457	243	1,181
Illinois-----	28	1	43	21,522	8	8	7,355	1,558	947
Michigan-----	18	-	178	8,621	2	1	4,267	752	1,233
Wisconsin-----	4	-	-	1,144	-	-	985	144	1,340
WEST NORTH CENTRAL-----	75	1	91	12,134	6	3	7,064	725	1,006
Minnesota-----	3	-	-	987	-	-	294	35	313
Iowa-----	12	-	-	921	-	-	1,286	34	94
Missouri-----	33	-	75	6,599	6	2	3,782	467	299
North Dakota-----	4	-	1	232	-	-	41	10	72
South Dakota-----	8	-	-	570	-	-	167	29	13
Nebraska-----	3	-	1	932	-	-	196	14	30
Kansas-----	12	1	14	1,893	-	1	1,298	136	185
SOUTH ATLANTIC-----	178	15	593	68,433	183	215	26,198	6,550	2,643
Delaware-----	5	-	5	597	-	10	1,312	271	14
Maryland-----	14	-	17	7,377	9	4	2,516	393	85
District of Columbia-----	6	-	40	10,077	13	52	2,019	549	18
Virginia-----	43	1	22	7,249	12	13	4,046	652	613
West Virginia-----	22	-	7	1,419	1	-	1,149	152	464
North Carolina-----	17	2	109	12,125	33	21	4,298	1,001	401
South Carolina-----	13	6	37	5,433	16	13	4,038	1,023	449
Georgia-----	33	3	159	13,827	52	58	3,634	1,567	135
Florida-----	25	3	197	10,329	47	44	3,186	942	464
EAST SOUTH CENTRAL-----	131	16	177	26,493	55	61	5,521	1,505	4,221
Kentucky-----	44	-	4	3,127	-	2	1,890	123	2,428
Tennessee-----	36	4	43	11,997	14	16	1,507	445	1,347
Alabama-----	26	11	35	3,668	18	22	1,286	729	231
Mississippi-----	25	1	95	7,701	23	21	838	208	215
WEST SOUTH CENTRAL-----	239	33	232	32,005	28	63	14,870	2,838	4,347
Arkansas-----	32	-	8	4,906	5	3	2,988	531	156
Louisiana-----	85	3	89	5,847	6	29	6,614	828	19
Oklahoma-----	16	-	21	5,114	1	2	1,932	284	152
Texas-----	106	30	114	16,138	16	29	3,336	1,195	4,020
MOUNTAIN-----	78	-	54	7,277	1	6	3,911	992	2,203
Montana-----	8	-	-	296	-	-	200	42	169
Idaho-----	7	-	-	269	-	-	62	19	57
Wyoming-----	4	-	-	61	-	-	55	10	5
Colorado-----	9	-	20	1,190	-	-	380	89	197
New Mexico-----	^b 31	-	7	2,151	-	-	1,254	322	495
Arizona-----	9	-	18	2,733	1	5	1,655	441	1,145
Utah-----	-	-	-	214	-	-	169	28	105
Nevada-----	10	-	9	363	-	1	136	41	30
PACIFIC-----	109	4	115	21,479	10	29	7,513	2,125	5,393
Washington-----	3	-	5	1,635	1	-	600	62	1,018
Oregon-----	10	-	3	916	1	-	716	80	538
California-----	96	4	107	18,928	8	29	6,197	1,983	3,837

⁷For complete title, see table 7.^aNotifiable in New York City only.^bIncludes paratyphoid fever (041).

Morbidity and Mortality Weekly Report

Table 5. CASES OF INFREQUENTLY REPORTED NOTIFIABLE DISEASES BY STATE: UNITED STATES, 1958

(Numbers under diseases are category numbers of the Seventh Revision of the International Lists, 1955)

STATES REPORTING CASES	Anthrax 062	Botulism 049.1	Glanders 064.2	Leprosy 060	Lepto- spirosis 072	Q fever 108 pt.
Arkansas-----	7	-	-	-	4	-
California-----	-	1	-	12	8	-
Colorado-----	-	-	-	-	1	-
Connecticut-----	-	-	-	-	1	-
Iowa-----	-	-	-	-	15	-
Kansas-----	-	-	1	-	-	-
Kentucky-----	-	2	-	-	2	-
Louisiana-----	2	-	-	4	1	-
Maine-----	-	-	-	-	1	-
Maryland-----	-	-	-	-	1	1
Massachusetts-----	2	-	-	1	-	-
Michigan-----	-	-	-	-	4	-
Mississippi-----	1	-	-	-	1	-
Missouri-----	-	-	-	-	1	-
Nebraska-----	-	-	-	-	6	-
Nevada-----	-	-	-	1	-	-
New Jersey-----	-	-	-	-	1	-
New York-----	-	-	-	16	2	-
North Carolina-----	-	1	-	-	-	-
Ohio-----	-	-	-	-	1	-
Oklahoma-----	-	1	-	-	1	-
Oregon-----	-	-	-	1	-	-
Pennsylvania-----	4	-	-	-	2	6
Texas-----	-	-	-	14	2	-
Washington-----	-	1	-	-	-	1

¹Notifiable in New York City only.

Morbidity and Mortality Weekly Report

Table 6. CASES OF SPECIFIED DISEASES REPORTED IN CERTAIN STATES: UNITED STATES, 1958

(Numbers under diseases are category numbers of the Seventh Revision of the International Lists, 1955)

AREA	Actino- mycosis 132	Ancy- losto- miasis (hook- worm) 129	Ascari- asis 130.0	Chicken- pox 087	Coccidi- o- mycosis (Valley fever) 133	Diarrhea of newborn 764	Erysip- elas 052	Histo- plas- mosis 134.2	Influenza 480-483	Kerato- conjunc- tivitis, infecti- ous 096.1
UNITED STATES-----	14	14,113	12	187,467	873	1,550	290	142	138,935	27
NEW ENGLAND:										
Maine-----	**	**	**	2,542	**	**	**	**	515	**
New Hampshire-----	**	**	**	1,031	**	**	**	**	13	**
Vermont-----	**	**	**	1,787	**	**	**	**	397	**
Massachusetts-----	-	**	**	15,662	**	**	**	**	**	**
Rhode Island-----	**	**	**	1,327	**	**	**	**	**	**
Connecticut-----	-	100	*	9,159	*	-	13	*	2,355	*
MIDDLE ATLANTIC:										
New York-----	**	¹ 956	**	26,667	**	134	**	**	¹ 802	1
New Jersey-----	*	*	*	*	*	6	*	*	**	*
Pennsylvania-----	1	120	**	**	**	**	**	**	**	**
EAST NORTH CENTRAL:										
Ohio-----	**	**	**	12,534	**	864	9	5	**	**
Indiana-----	**	**	**	3,902	**	**	1	15	**	5
Illinois-----	1	-	*	11,631	*	159	140	-	581	-
Michigan-----	1	-	-	11,356	-	213	45	-	474	7
Wisconsin-----	**	**	**	33,430	**	**	**	**	355	**
WEST NORTH CENTRAL:										
Minnesota-----	**	2	5	2,534	2	43	2	14	**	2
Iowa-----	1	**	**	4,855	**	**	**	15	**	**
Missouri-----	**	**	**	1,897	**	1	2	16	1,462	**
North Dakota-----	**	**	**	2,570	**	**	8	**	2,526	**
South Dakota-----	**	**	**	210	**	**	2	**	**	**
Nebraska-----	**	**	**	297	**	**	**	**	**	**
Kansas-----	5	**	**	**	**	3	1	**	**	**
SOUTH ATLANTIC:										
Delaware-----	-	-	-	331	-	-	-	-	**	-
Maryland-----	-	-	-	1,629	-	1	-	2	**	-
District of Columbia-----	-	-	-	386	-	5	2	-	289	7
Virginia-----	**	**	**	**	**	**	**	**	**	**
West Virginia-----	**	**	**	3,196	**	18	1	1	**	**
North Carolina-----	**	**	**	**	**	**	**	**	**	**
South Carolina-----	**	**	**	**	**	**	**	**	**	**
Georgia-----	**	9,037	**	**	**	**	**	**	2,733	**
Florida-----	**	3,183	**	4,456	**	64	**	**	**	**
EAST SOUTH CENTRAL:										
Kentucky-----	-	6	4	6,482	-	-	2	21	24,912	-
Tennessee-----	**	**	**	**	**	**	**	**	69,584	**
Alabama-----	1	**	**	1,557	**	**	**	**	**	**
Mississippi-----	**	696	**	378	**	**	**	16	**	**
WEST SOUTH CENTRAL:										
Arkansas-----	-	10	3	542	-	-	*	33	**	-
Louisiana-----	1	*	*	*	*	*	*	4	*	*
Oklahoma-----	*	*	*	*	*	*	*	*	**	*
Texas-----	**	**	**	**	**	**	**	**	**	**
MOUNTAIN:										
Montana-----	**	**	**	3,591	**	**	**	**	**	**
Idaho-----	-	-	-	901	-	-	3	-	-	-
Wyoming-----	1	-	-	651	**	**	**	**	**	**
Colorado-----	*	*	*	*	*	*	*	*	2,613	*
New Mexico-----	-	-	-	3,495	2	**	4	-	16,100	-
Arizona-----	1	-	-	3,562	647	**	33	**	**	**
Utah-----	**	**	**	2,076	**	1	**	**	**	**
Nevada-----	**	**	**	255	**	**	**	**	115	**
PACIFIC:										
Washington-----	*	1	*	10,511	1	1	(³)	-	**	-
Oregon-----	1	2	-	² 77	-	16	22	*	13,109	5
California-----	*	*	*	*	221	21	*	*	**	*

¹Notifiable in New York City only.²Persons 16 years of age and over.³Included in scarlet fever and streptococcal sore throat (050,051).

Morbidity and Mortality Weekly Report

13

Table 6. CASES OF SPECIFIED DISEASES REPORTED IN CERTAIN STATES: UNITED STATES, 1958—Continued

(Numbers under diseases are category numbers of the Seventh Revision of the International Lists, 1955)

AREA	Lympho- cytic choriomen- ingitis, acute 082.1	Malignant neoplasms (cancer) 140-199	Mononu- cleosis, infectious (glandular fever) 093	Mumps 089	Pellagra 281	Pneu- monias 490-493, 763	Rubella (German measles) 086	Taeniasis (tape- worm) 126 pt.	Vincent's angina 070 pt.
UNITED STATES-----	55	50,488	3,022	160,028	4	20,150	172,411	6	301
NEW ENGLAND:									
Maine-----	(4)	**	**	3,730	**	492	1,054	**	**
New Hampshire-----	**	**	6	1,768	**	95	913	**	**
Vermont-----	**	**	145	1,480	**	**	**	**	**
Massachusetts-----	2	**	**	9,617	**	**	12,921	**	**
Rhode Island-----	**	2,788	27	630	**	**	1,282	**	2
Connecticut-----	*	*	485	4,348	*	1,386	11,304	*	*
MIDDLE ATLANTIC:									
New York-----	**	5 ⁵ 24,017	**	14,656	**	1 ² 2,677	22,297	**	**
New Jersey-----	**	*	*	*	*	**	*	*	*
Pennsylvania-----	5	**	149	**	**	**	**	**	**
EAST NORTH CENTRAL:									
Ohio-----	**	**	31	9,508	**	**	6,854	**	3
Indiana-----	**	**	119	4,373	**	**	5,853	**	41
Illinois-----	2	*	501	9,538	*	3,587	15,431	*	134
Michigan-----	-	**	189	15,209	-	**	5,073	-	-
Wisconsin-----	**	**	4	21,926	**	**	44,953	**	**
WEST NORTH CENTRAL:									
Minnesota-----	3	**	789	190	1	**	**	5	**
Iowa-----	1	**	20	6,262	**	**	4,139	**	**
Missouri-----	**	**	120	1,172	**	**	**	**	**
North Dakota-----	**	**	**	4,461	**	4,269	**	**	**
South Dakota-----	**	**	13	51	**	33	**	**	1
Nebraska-----	**	**	**	245	**	**	(a)	**	**
Kansas-----	3	**	**	**	**	**	**	**	**
SOUTH ATLANTIC:									
Delaware-----	-	1,231	-	122	-	**	250	-	-
Maryland-----	4	**	-	569	-	**	2,281	-	1
District of Columbia-----	-	**	10	76	-	734	392	-	-
Virginia-----	**	**	**	**	**	**	**	**	**
West Virginia-----	**	**	4	2,407	**	**	2,569	**	**
North Carolina-----	**	**	**	**	**	**	**	**	**
South Carolina-----	**	3,581	15	**	**	**	**	**	**
Georgia-----	**	**	**	**	**	**	**	**	**
Florida-----	**	2,485	**	2,702	**	**	1,005	1	**
EAST SOUTH CENTRAL:									
Kentucky-----	-	**	132	7,586	-	1,411	3,392	-	52
Tennessee-----	**	**	**	**	**	**	**	**	**
Alabama-----	**	6,815	**	1,190	2	**	2,065	**	**
Mississippi-----	11	**	62	171	**	**	1,549	**	**
WEST SOUTH CENTRAL:									
Arkansas-----	9	1,242	24	362	-	**	3,883	**	**
Louisiana-----	2	5,149	*	*	*	*	*	*	*
Oklahoma-----	*	*	*	*	*	**	*	*	*
Texas-----	**	**	**	**	**	**	**	**	**
MOUNTAIN:									
Montana-----	**	**	**	**	**	**	**	**	**
Idaho-----	-	1,447	26	1,038	-	**	653	-	8
Wyoming-----	**	557	7	35	**	**	**	**	**
Colorado-----	*	*	*	*	*	*	1,434	*	*
New Mexico-----	-	7482	12	3,184	-	3,118	3,496	-	3
Arizona-----	5	**	43	3,046	1	**	4,298	**	**
Utah-----	2	680	30	1,013	**	**	1,076	**	2
Nevada-----	**	14	2	182	**	**	**	**	6
PACIFIC:									
Washington-----	*	*	57	9,910	-	**	5,837	-	48
Oregon-----	6	*	*	*	*	2,348	6,157	*	*
California-----	*	*	*	17,271	*	*	*	*	*

⁴Included in meningitis, other (340).

⁵Not notifiable in New York City.

⁶Included in measles (085).

⁷Includes neoplasms of lymphatic and hematopoietic tissues (200-204).

NOTE.—Cases of other diseases reported are: Colorado tick fever (096.9 pt.)—Colorado 79, Oregon 26; echinococcosis (hydatid disease) (125)—New York City 4; relapsing fever (tick-borne) (071.1)—California 2, Oregon 4.

Morbidity and Mortality Weekly Report

Table 7. REPORTED CASES OF SPECIFIED NOTIFIABLE DISEASES: ALASKA, HAWAII, PUERTO RICO, VIRGIN ISLANDS, GUAM, AND THE CANAL ZONE, 1958

(Numbers after diseases are category numbers of the Seventh Revision of the International Lists, 1955. Only diseases with frequencies are listed. For a complete list of diseases, see table 2)

DISEASE	Alaska	Hawaii	Puerto Rico	Virgin Islands	Guam	Canal Zone
Amebiasis-----046	-	1	11	3	6	5
Anthrax-----062	-	-	-	-	4	-
Brucellosis (undulant fever)-----044	1	1	1	-	2	-
Diphtheria-----055	-	-	55	-	1	10
Dysentery, bacillary (shigellosis)-----045	7	246	-	56	91	11
Encephalitis, acute infectious-----082	-	2	2	-	1	-
Hepatitis, infectious, and serum-----092,N998.5 pt.	74	71	185	-	65	18
Leprosy-----060	-	18	1	-	3	-
Leptospirosis-----072	-	6	1	-	-	-
Malaria-----110-117	-	1	-	-	-	103
Measles-----085	1,386	485	3,347	-	55	-
Meningococcal infections-----057	-	2	4	-	6	1
Meningitis, other-----340	19	-	58	2	-	-
Poliomyelitis, acute-----080	2	78	48	-	-	1
Paralytic-----080.0,080.1	1	78	46	-	-	1
Nonparalytic-----080.2	1	-	2	-	-	---
Unspecified-----080.3	-	-	-	-	-	---
Rabies in animals-----	-	-	17	-	-	-
Rheumatic fever-----400-402	5	3	2	-	24	2
Salmonellosis, except typhoid fever-----041,042	-	152	-	6	3	2
Scarlet fever and streptococcal sore throat-----050,051	413	147	343	-	322	3
Tetanus-----061	-	-	116	-	-	-
Trichinosis-----128	-	1	-	-	2	-
Tuberculosis, all forms-----001-019	429	463	2,800	9	148	28
Typhoid fever-----040	-	1	22	1	-	-
Typhus fever, endemic (murine)-----101	-	2	-	-	-	-
Venereal diseases:						
Chancroid-----036	-	15	53	-	7	2
Gonorrhoea-----030-034	207	570	3,646	73	198	68
Granuloma inguinale-----038	-	-	-	-	-	-
Lymphogranuloma venereum-----037	1	1	2	-	-	7
Syphilis, primary, secondary, and early latent-----021,028.0	8	10	380	---	-	---
Syphilis, other forms-----020,022-027,028.1,028.2,029	12	146	933	---	3	---
Whooping cough (pertussis)-----056	46	5	739	119	3	5