



COVID-19

COVID-19 Vaccine Boosters

Updated Apr. 19, 2022

What You Need to Know

- As with vaccines for other diseases, people are best protected against infection with the virus that causes COVID-19 when they stay up to date with vaccinations.
- COVID-19 vaccines continue to protect people from getting seriously ill, being hospitalized, and even dying—especially people who have received a booster.
- CDC recommends everyone ages 12 years and older receive a COVID-19 vaccine booster after completing their primary COVID-19 vaccination series. Some people can receive two boosters.
- People who are moderately or severely immunocompromised have specific COVID-19 vaccine recommendations, including recommendations for a booster. Learn more about COVID-19 vaccine recommendations for people who are moderately or severely immunocompromised.

Choosing Your COVID-19 Booster

Three COVID-19 vaccines are authorized or approved for use in the United States to prevent COVID-19. Pfizer-BioNTech or Moderna (COVID-19 mRNA vaccines) are preferred. You may get Johnson & Johnson's Janssen COVID-19 vaccine in some situations.

Who Can Get a Booster

PRIMARY SERIES COVID-19 VACCINE
Pfizer-BioNTech

Who should get one booster:

Everyone 12 years and older

Who can get a second booster:

Adults 50 years and older

When to get your booster:

At least 5 months after completing your primary COVID-19 vaccination series

If eligible for a second booster, at least 4 months after your first booster

Which booster should you get?

- Adults 18 years and older should get an mRNA COVID-19 vaccine (Pfizer-BioNTech or Moderna) for the first booster in most* situations
- The second booster must be an mRNA COVID-19 vaccine
- Teens 12–17 years old may only get a Pfizer-BioNTech COVID-19 vaccine booster

PRIMARY SERIES COVID-19 VACCINE

Moderna

Who should get one booster:

Adults 18 years and older

Who can get a second booster:

Adults 50 years and older

When to get your booster:

At least 5 months after completing your primary COVID-19 vaccination series

If eligible for a second booster, at least 4 months after your first booster

Which booster should you get?

For the first booster, an mRNA COVID-19 vaccine (Pfizer-BioNTech or Moderna) is preferred in most* situations

The second booster must be an mRNA COVID-19 vaccine

PRIMARY SERIES COVID-19 VACCINE

Johnson & Johnson's Janssen*

Who should get a booster:

Adults 18 years and older

Who can get a second booster:

Anyone who received a J&J/Janssen COVID-19 vaccine for both their primary dose and booster

Adults 50 years and older who first received a J&J/Janssen COVID-19 vaccine, regardless of what type of booster they received

When to get a booster:

At least 2 months after receiving your J&J/Janssen COVID-19 vaccination

If eligible for a second booster, at least 4 months after your first booster

Which booster should you get?

For the first booster, an mRNA COVID-19 vaccine (Pfizer-BioNTech or Moderna) is preferred in most* situations

The second booster must be an mRNA COVID-19 vaccine

*Although mRNA vaccines are preferred for the first booster, J&J/Janssen COVID-19 vaccine may be considered in some situations.

Scheduling Your Booster

If you need help scheduling your booster, contact the location that set up your previous appointment. If you need to get your booster in a location different from where you received your previous vaccination, there are several ways you can find a vaccine provider.

Find a COVID-19 vaccine or booster: Search vaccines.gov, text your ZIP code to 438829, or call 1-800-232-0233 to find locations near you.

What to Expect during and after Your Booster Shot Appointment

- Bring your CDC COVID-19 Vaccination Record card to your booster appointment so your provider can fill in the information about your booster dose. If you did not receive a card at your first appointment, contact the vaccination site where you got your first dose or your state health department to find out how you can get a card.
- You may experience side effects after getting a COVID-19 vaccine. These are normal signs that your body is building protection against COVID-19.
- If you have enrolled in **v-safe**, enter your booster dose in your **v-safe** account, the system will send you daily health check-ins. You will be able to tell CDC about any side effects.
- A person is considered "boosted" and up to date right after getting their first booster. Getting a second booster is not necessary to be considered up to date at this time.

Related Pages

- > COVID-19 Vaccine Safety and Monitoring
- > Understanding How COVID-19 Vaccines Work
- > Ensuring COVID-19 Vaccines Work
- Frequently Asked Questions about COVID-19 Vaccination
- > COVID-19 Vaccines for Moderately to Severely Immunocompromised People
- > People Vaccinated Outside the United States



For Healthcare and Public Health

Considerations for Use of a COVID-19 Vaccine Booster Dose

More Information

ACIP Presentation Slides, December 16, 2021

ACIP Presentation Slides, November 19, 2021

ACIP Presentation Slides. October 21, 2021

ACIP Presentation Slides, September 22–23, 2021

Last Updated Apr. 19, 2022