

THE LANCET Infectious Diseases


Supplementary webappendix

This webappendix formed part of the original submission and has been peer reviewed. We post it as supplied by the authors.

Supplement to: Sáez-Llorens X, Clemens R, Leroux-Roels G, et al. Immunogenicity and safety of a novel monovalent high-dose inactivated poliovirus type 2 vaccine in infants: a comparative, observer-blind, randomised, controlled trial. *Lancet Infect Dis* 2015; published online Dec 21. [http://dx.doi.org/10.1016/S1473-3099\(15\)00488-0](http://dx.doi.org/10.1016/S1473-3099(15)00488-0).

APPENDIX

Figure : Reverse cumulative distribution of type 2 poliovirus shedding index after mOPV2 challenge





Vaccine Type	AGE of Admin	N° DE DOSIS	INTERVAL BETWEEN DOSES	DOSE, Administration way
Hepatitis B (1)	*Newborns before 12 hours	1	-	0.5 c.c intramuscular anterolateral in Thigh
B.C.G (2)	Newborns	1	-	0.05 l.c.c .01 cc Intradermal Left Deltoid
Pneumococcal Conjugate	2-4 months	2	4-8 Weeks	0.5 c.c. intramuscular anterolateral in Thigh
Oral Polio	2-4-6 months	3	8 Weeks	2 oral drops.
Inactivated Polio (Intramuscular) (3)	2-4-6 months	3	4-8 Weeks	0.5 c. c intramuscular anterolateral in Thigh
Hexavalent (4)	2-4-6 months	3	4-8 Weeks	0.5 cc intramuscular anterolateral in Thigh
Pentavalent (DPT-Hib-Hep B) (5)	2-4-6 meses	3	8 Weeks	0.5 c.c intramuscular anterolateral in Thigh
Influenza (6)	6-11 meses	2	4 Weeks	0.25 c.c intramuscular anterolateral in Thigh or deltoids
Rotavirus (7)	2-8 meses	2	4-8 Weeks	1 c.c oral

1. If New born has more than 12 hours DO NOT ADMINISTER.
2. Administer in all newborns no matter their weight, it can be administered until 15 years of age.
3. Only will administered in immunosupressed subjects or with suspected HIV infection.
4. **To be administered in children born after april 2014. Do Not Administer Oral Polio**, Can be administered until 24 months of age, if the child appears at 15 months without previous administration.
5. **Only to be administered in children that already received a Penta Vaccine dose.**
6. First time 2 doses will be administered in the child younger than 12 months begin administration at 6 months. If Pneumo or Penta are administered, please administered influenza on Deltoid area.
7. Avoid administering in children older than 8 months. can be administered in mothers HIV + and Immunosupressed subjects.