



HHS Public Access

Author manuscript

Am J Prev Med. Author manuscript; available in PMC 2023 January 01.

Published in final edited form as:

Am J Prev Med. 2022 January ; 62(1): e45–e55. doi:10.1016/j.amepre.2021.06.021.

Intimate Partner and Sexual Violence Prevention Among Youth: A Community Guide Systematic Review

Ramona K.C. Finnie, DrPH¹, Devon L. Okasako-Schmucker, MPH¹, Leigh Buchanan, PhD², Denise Carty, PhD³, Holly Wethington, PhD¹, Shawna L. Mercer, MSc, PhD⁴, Kathleen C. Basile, PhD⁵, Sarah DeGue, PhD⁵, Phyllis Holditch Nolon, PhD⁵, Jennifer Bishop, ScD, MPH⁶, Tisha Titus, MD, MPH⁷, Samia Noursi, PhD⁸, Shavon Artis Dickerson, DrPH⁶, Daniel Whitaker, PhD⁹, Susan Swider, PhD¹⁰, Patrick Remington, MD, MPH¹¹, The Community Preventive Services Task Force

¹Community Guide Office, Office of the Associate Director for Policy and Strategy, Centers for Disease Control and Prevention, Atlanta, Georgia

²Cherokee Nation Assurance, Arlington, Virginia

³Office of Women's Health, Centers for Disease Control and Prevention, Atlanta, Georgia

⁴Office of the Director, Center for Surveillance, Epidemiology, and Laboratory Sciences, Centers for Disease Control and Prevention, Atlanta, Georgia

⁵Division of Violence Prevention, National Center for Injury Prevention and Control, Centers for Disease Control and Prevention, Atlanta, Georgia

⁶Office of Women's Health, Health and Human Services, Washington, District of Columbia

⁷American College of Preventive Medicine, Washington, District of Columbia

⁸Office of Research on Women's Health, National Institutes of Health, Bethesda, Maryland

⁹School of Public Health, Georgia State University, Atlanta, Georgia

Address correspondence to: Ramona K.C. Finnie, DrPH, Community Guide Office, Office of the Associate Director for Policy and Strategy, Centers for Disease Control and Prevention, 1600 Clifton Road, Mailstop V25-5, Atlanta GA 30329. vng7@cdc.gov.

No financial disclosures were reported by the authors of this paper.

CREDIT AUTHOR STATEMENT

Ramona K.C. Finnie: Data Curation, Formal analysis, Investigation, Methodology, Validation, Visualization, Writing - Original Draft, Writing - Review & Editing. Devon L. Okasako-Schmucker: Formal analysis, Investigation, Methodology, Validation, Visualization, Writing - Original Draft, Writing - Review & Editing. Leigh Buchanan: Investigation, Validation, Writing - Review & Editing. Denise Carty: Investigation, Validation, Writing - Review & Editing. Holly Wethington: Investigation, Validation, Writing - Review & Editing. Shawna L. Mercer: Conceptualization, Supervision, Writing - Review & Editing. Kathleen C. Basile: Conceptualization, Subject Matter Input, Writing - Review & Editing. Sarah DeGue: Conceptualization, Subject Matter Input, Writing - Review & Editing. Phyllis Holditch Nolon: Conceptualization, Subject Matter Input, Writing - Review & Editing. Jennifer Bishop: Conceptualization, Writing - Review & Editing. Tisha Titus: Conceptualization, Writing - Review & Editing. Samia Noursi: Conceptualization, Writing - Review & Editing. Shavon Artis Dickerson: Conceptualization, Writing - Review & Editing. Daniel Whitaker: Conceptualization, Writing - Review & Editing. Susan Swider: Conceptualization, Supervision, Writing - Review & Editing. Patrick Remington: Conceptualization, Supervision, Writing - Review & Editing. The Community Preventive Services Task Force: Supervision.

SUPPLEMENTAL MATERIAL

Supplemental materials associated with this article can be found in the online version at <https://doi.org/10.1016/j.amepre.2021.06.021>.

Names and affiliations of the Community Preventive Services Task Force members can be found at: www.thecommunityguide.org/about/task-force-members.html.

¹⁰College of Nursing, Rush University, Chicago, Illinois

¹¹School of Medicine and Public Health, University of Wisconsin-Madison, Madison, Wisconsin

Abstract

Introduction: Intimate partner violence and sexual violence are widespread and often occur early in life. This systematic review examines the effectiveness of interventions for primary prevention of intimate partner violence and sexual violence among youth.

Methods: Studies were identified from 2 previous systematic reviews and an updated search (January 2012–June 2016). Included studies were implemented among youth, conducted in high-income countries, and aimed to prevent or reduce the perpetration of intimate partner violence or sexual violence. In 2016–2017, *Guide to Community Preventive Services (Community Guide)* methods were used to assess effectiveness as determined by perpetration, victimization, or bystander action. When heterogeneity of outcomes prevented usual *Community Guide* methods, the team systematically applied criteria for favorability (statistically significant at $p < 0.05$ or approaching significance at $p < 0.10$) and consistency (75% of results in the same direction).

Results: A total of 28 studies (32 arms) met inclusion and quality of execution criteria. Interventions used combinations of teaching healthy relationship skills, promoting social norms to protect against violence, or creating protective environments. Overall, 18 of 24 study arms reported favorable results on the basis of the direction of effect for decreasing perpetration; however, favorability for bystander action diminished with longer follow-up. Interventions did not demonstrate consistent results for decreasing victimization. A bridge search conducted during Fall 2020 confirmed these results.

Discussion: Interventions for the primary prevention of intimate partner violence and sexual violence are effective in reducing perpetration. Increasing bystander action may require additional follow-up as effectiveness diminishes over time. Findings may inform researchers, school personnel, public health, and other decision makers about effective strategies to prevent intimate partner violence and sexual violence among youth.

INTRODUCTION

Intimate partner violence (IPV) and sexual violence (SV) are major public health problems that occur across every stage of life but often begin during adolescence.¹ IPV, including dating violence, is committed by a current or former intimate partner and includes physical violence or SV, stalking, or psychological aggression.² SV may be committed by a current or former intimate partner or by someone else; it includes sexual acts (e.g., kissing, touching, intercourse) committed or attempted without consent or against an individual who is unable to consent or refuse.³ In 2019, a total of 10.8% of American high-school students reported experiencing some sort of SV; 8.2% of students who had dated in the previous 12 months reported experiencing sexual dating violence, 8.2% reported experiencing physical dating violence, and 3.0% experienced both in the past year.⁴ Prevalence of IPV and SV was higher among female than among male youth, with 16.4% of female and 8.2% of male high schoolers experiencing any dating violence.⁴ Lesbian, gay, bisexual, or questioning students reported a significantly greater prevalence of any dating violence and SV than

heterosexual students.⁴ Prevalence of any dating violence did not vary significantly by race or ethnicity⁴; however, some research suggests that racial and ethnic minority youth may be at higher risk.⁵ Experiences of IPV and SV have consequences for youth, including physical injury, substance abuse, poor mental health, and low academic achievement.^{6,7} Preventing perpetration (rather than general awareness, risk reduction, victimization response, and other types of secondary and tertiary prevention) has the greatest potential to reduce population rates of violence and its health consequences.^{8–10} Thus, adolescence is a critical time to promote attitudes and behaviors that could prevent IPV and SV across the lifespan.¹¹

The public health approach to IPV and SV is to prevent or reduce a person's risk of perpetrating IPV and SV.¹⁰ Primary prevention interventions may be geared toward potential perpetrators or bystanders—those who can challenge violence-supportive norms by directly reducing risk (e.g., noticing a risky social situation and intervening) or by indirectly reducing risk (e.g., challenging hostile attitudes toward women such as offensive jokes or objectifying language)^{12–15}—and may also reduce victimization (an act that makes someone a victim).^{2,3} IPV and SV technical packages,^{13,15} developed by the Centers for Disease Control and Prevention Division of Violence Prevention, compile and prioritize primary prevention strategies and actions to help states, local communities, and organizations reduce IPV and SV.^{13,15} This *Guide to Community Preventive Services (Community Guide)* systematic review aligns with these technical packages and builds on 2 existing systematic reviews^{16,17} to examine the evidence of effectiveness on perpetration, victimization, and bystander action of primary prevention interventions designed to reduce IPV or SV among youth aged 12–24 years.

Methods

Community Guide methods were used for this review.¹⁸ The review coordination team (called the team in the remaining part of this paper) was composed of subject-matter experts in IPV or SV from various agencies and institutions along with systematic review experts from the Community Guide Office at the Centers for Disease Control and Prevention. The team worked under the oversight of the independent, nonpartisan, nonfederal, unpaid Community Preventive Services Task Force.

Conceptual Approach

The team defined interventions for the primary prevention of IPV and SV as those that aim to prevent or reduce the perpetration of IPV and SV and promote healthier relationships between peers and partners. Interventions must take place among youth aged 12–24 years.¹⁹ The interventions included in this review provided educational information about how to recognize IPV or SV, the warning signs, or the consequences of IPV or SV. The interventions could also focus on 1 of the following strategies: teaching healthy relationship skills, promoting social norms that protect against violence, and creating protective environments (e.g., improving school climate and safety) (Appendix Table 1, available online). Interventions were implemented in schools, homes, or communities or in a combination of settings. They either targeted the general population or high-risk groups for violence, which may have included youth who previously experienced IPV or SV as a victim or perpetrator.

Interventions may modify social norms around violence by increasing awareness and knowledge of IPV and SV, improving attitudes toward gender equity, and decreasing acceptance of IPV and SV. Interventions implemented by policymakers may increase access to available resources and support within communities to create protective environments. Interventions may lead to improved relationship skills, increased self-efficacy, and improved conflict resolution skills, leading to decreased risk behavior. Through these pathways, primary prevention interventions may reduce perpetration and victimization and decrease morbidity, mortality, and disparities. In addition, these interventions may increase bystander action, which may also reduce victimization and perpetration. Potential effect modifiers include peer influence, risk behaviors, structural factors (e.g., racism, poverty), and population characteristics, including race, age, sex, and SES. Primary prevention interventions may also have the additional benefits of increasing school achievement and decreasing peer violence, such as bullying.^{13,15}

Search for Evidence

The search for evidence consisted of 3 steps. The first step involved searching for existing systematic reviews on the effectiveness of IPV and SV interventions. Two existing systematic reviews were identified: the Whitaker et al.¹⁷ systematic review focused on IPV (search period through 2013) and the DeGue and colleagues¹⁶ systematic review focused on SV (search period through 2014). The second step involved combining IPV and SV interventions into a single review because intervention strategies and outcomes of interest were in alignment. The third step was updating the search, merging the search terms used in both reviews. The updated literature search was from January 2012 to June 2016. Searches were conducted in PsycNET, PsycExtra, PubMed, ERIC, Sociological Abstracts, MEDLINE, Web of Knowledge, Dissertation Abstracts International, and Google Scholar. Reference lists in retrieved articles were also reviewed. The search is available on the *Community Guide* website under IPV/SV Supporting Materials.

Inclusion and Exclusion Criteria

Inclusion and exclusion criteria aligned with those of Whitaker et al.¹⁷ and DeGue and colleagues¹⁶ so that studies were included if they evaluated the primary prevention of IPV or SV among youth aged 12–24 years. *Community Guide* methods include a range of study designs to better assess the effectiveness of public health interventions. For this review, studies were included if they had concurrent comparison groups. The team further restricted to studies that also met the following criteria: (1) reported 1 of the following behavioral outcomes: perpetration, victimization, or bystander action; (2) conducted in a very high human development Index country, as classified by the UN Development Program (for comparability to U.S. populations)²⁰; (3) peer-reviewed manuscripts; and (4) published in English.

Studies that included interventions to prevent victimization but did not address perpetration (e.g., self-defense or other interventions to modify the potential victim's behavior) were excluded because they did not focus on changing the behavior of potential perpetrators, which is also consistent with the Whitaker et al.¹⁷ and DeGue and colleagues¹⁶ reviews.

Studies that combined intervention groups or compared one intervention with another intervention without including an untreated control group were excluded.

Outcomes of Interest

Effectiveness outcomes were assessed using self-reported perpetration, victimization, and bystander action as measured below:

Perpetration and victimization were assessed using self-reported standard scales such as the Conflict in Adolescent Dating Relationships Inventory,²¹ Peer Rejection Questionnaire,^{22–24} Revised Conflict Tactics Scale,²⁵ Safe Dates Dating Violence Scale,²⁶ Sexual Experiences Survey,^{27–29} and Sexual Harassment Survey.³⁰ A decrease *in perpetration and victimization* was defined as favorable.

Bystander action was measured using a variety of scales, including the Bystander Behavior Scale,^{31,32} Sexual Social Norms Inventory,³³ and Reactions to Offensive Language and Behavior Scale.³⁴ An increase in bystander action was favorable.

Assessing and Summarizing the Body of Evidence on Effectiveness

Each included study was independently abstracted by 2 reviewers. Abstraction was based on a standardized abstraction form³⁵ that included information on study quality, intervention components, participant demographics, and outcomes. Disagreements between reviewers were resolved by team consensus. Threats to validity were used to characterize studies as having good (0–1 limitation), fair (2–4), or limited (5) quality of execution.¹⁸ These included internal and external threats to validity such as poor description of the intervention, population, or sampling frame; poor measurement of exposure or outcome; poor reporting of analytic methods; loss to follow-up; or intervention and comparison groups not being comparable at baseline. Studies with limited quality of execution (5 limitations) were excluded from the analyses.

Calculation of the Effect Estimates for Qualifying Studies

Effect estimates were calculated for each study when possible.¹⁸ The formula for calculating effect estimates was carried out using 1 of 2 methods, depending on study design and variability of the outcome. The preferred method included nontreated comparison (C) and intervention (I) groups, the basic units for the calculation, with measurements made before (pre) and after (post) the intervention. For studies with multiple intervention arms meeting inclusion criteria and a single nontreated comparison arm, effect estimates for each intervention arm were calculated using the same comparison arm. The team calculated absolute percentage point difference using the following formula:

$$(I_{\text{post}} - I_{\text{pre}}) - (C_{\text{post}} - C_{\text{pre}}).$$

To pool data from studies reporting different measures to assess the same outcome, relative percent change was calculated using the following formula:

$$\left(\frac{I_{\text{post}} - I_{\text{pre}}}{I_{\text{pre}}}\right) - \left(\frac{C_{\text{post}} - C_{\text{pre}}}{C_{\text{pre}}}\right) * 100.$$

Interquartile intervals (IQIs) were calculated when independent effect estimates were available for at least 5 studies; otherwise, the range of estimates was displayed. For studies with multiple publications, the publication with the latest data point was used in the analysis. In addition, this review stratified results by short-term follow-up (< 6 months) and longer-term follow-up (>6 months). For studies with multiple follow-ups, the team looked at the latest data point in both stratifications. Effect estimates that could not be combined on a scatterplot were described narratively.

Overall Determination of Favorability

Owing to the heterogeneity of outcome measures, effect estimates could not be pooled quantitatively because they typically are for *Community Guide* reviews. For example, many studies reported standardized and unstandardized β -coefficients that could not be combined. Therefore, the team ensured a systematic synthesis process by employing criteria and decision rules for favorability and consistency. First, the team assessed all studies (both those summarized quantitatively and qualitatively) for direction of effect. The result was considered favorable if the effect estimate was either statistically significant at $p < 0.05$ or approaching significance at $p < 0.10$ in favor of the intervention. The same criteria were applied to results that were in the unfavorable direction. Summary of the outcome was considered consistent if 75% of the study arms were in the same direction. Overall direction was determined by team consensus regardless of statistical significance.

RESULTS

A total of 3,153 citations were screened: 2,996 from the database search, 18 from included studies in Whitaker et al.,¹⁷ and 140 from those included in DeGue and colleagues.¹⁶ Full-text screening was conducted for 44 publications; 31 studies^{32,36–51,52–64} met inclusion criteria (Figure 1). Two studies were reported in 1 publication,⁴² 1 study was reported in 4 publications,^{26,52,65,66} 1 study was reported in 3 publications,^{36,67,68} and 2 studies were each reported over 2 publications.^{38,59,69,70} For the 4 studies that are represented by multiple publications, the publication with the latest data point was chosen as the main publication.^{36,38,52,59} Summary evidence tables for all included studies can be found at <https://www.thecommunityguide.org/sites/default/files/assets/SET-Violence-IPV-SV.pdf>.

Appendix Figure 1 (available online) displays the quality of execution assessment for included studies. A total of 8 studies^{38,42,47,52,61–63} had good quality of execution (< 1 limitation), 28 studies^{32,36,37,39–41,42–46,48,49,53–60} had fair quality of execution (2–4 limitations), and 3 studies^{50,51,64} were excluded owing to limited quality of execution (> 5 limitations). The most common limitations were for sampling (used convenience sampling, 16 studies)^{32,39–41,44,46–49,52–54,57–60} and loss to follow-up (15 studies).^{32,36–39,42,43,45,48,54,56,57,59,62,63}

Study Characteristics

Most included studies were conducted in the U.S.,^{32,36,38–49,52–61} whereas 2 others were in Canada,^{62,63} and 1 was in The Netherlands.³⁷ Almost half of the studies were implemented on college campuses,^{32,36,42,43,46,48,49,54–57} and nearly half took place in middle schools,^{38,45,47,60,62} high schools,^{37,40,41,58,59,63} or both.⁵² Two studies were implemented in the home,^{39,53} and 2 were implemented in community centers or agencies.^{44,62} Of the 17 studies reporting population density, most took place in urban areas^{36,40,41,44,45,47,58,59,61,62} or a mixture of urban and suburban,^{39,42,43,46,53,60,63} whereas 1 study⁵² took place in a rural area.

Population Characteristics

Study participants in included studies had a median age of 15.5 years^{32,38,39,41–44,46,47,49,52,54,58,62,63}; the median age of participants in studies implemented on college campuses was 19.4 years, and the median age of participants in studies implemented in middle and high schools was 13.9 years. A total of 17^{32,36–47,49,52–63} studies included participants identifying as either male or female: 48.1% were male, and 51.9% were female. A total of 7 studies focused on 1 sex: 6 studies^{37,49,54,55,57,59} included male participants only, and 1 study⁴⁴ included female participants only. Most study participants identified as White (median=69.9%),^{38,39,41–46,48,49,52–54,56,57,59–62} whereas the median proportion identifying as Black or African American was 16.1%,^{38,39,41,45,47,49,52,54,56,57,59–62} the proportion identifying as Hispanic or Latino was 12.6%,^{38,40–43,45,47–49,53,54,56–62} the proportion identifying as Asian was 6.9%,^{41,43,48,49,54,56,59–62} the proportion identifying as American Indian or Alaska Native was 2.2%,^{41,43,48,56,57,59} and the proportion identifying as other was 9.9%.^{36–39,41–43,45,47–49,52–54,56–61} One study⁴⁰ had an exclusively Hispanic or Latino population.

Intervention Characteristics

All studies included strategies that provided information on IPV or SV. Two study arms provided information but did not include any additional prevention strategies.^{58,60} In addition to providing information, 19 study arms in 17 studies^{36–41,44,45,47,52–55,60–63} also taught healthy relationship skills, and 21 arms in 20 studies^{23,25–39,56,59,60,64} also promoted social norms that protect against violence. Furthermore, 15 arms in 15 studies implemented bystander approaches.^{32,36,40,42,43,45,46,48,49,54,55,57,59,61} A total of 3 arms in 2 studies^{45,61} created protective environments (e.g., improving school climate by identifying hotspots and increasing staff presence).

Outcomes

Perpetration.—A total of 24 study arms from 21 studies^{36–41,44,45,47,49,52–55,57–63} reported perpetration. A total of 3 arms^{40,44,57} reported a median absolute decrease of 6.7 percentage points (ranging from –7.3 to –5.2 percentage points). A total of 3 arms^{52,54,59} reported a median relative decrease of 10.1% (ranging from –61.7% to 31.4%). A total of 10 arms from 8 studies^{37,38,41,47,49,53,61,63} reported a median decrease in odds of perpetration (OR=0.6, IQI=0.4, 0.8). A total of 9 arms reported data that could not be combined to

calculate a median: 5 arms^{36,45,53,55,62} reported decreases in perpetration, 1 arm⁶⁰ reported no change, and 3 arms^{39,58,60} reported increases in perpetration.

Overall, for perpetration, 18 of 24 arms from 17 studies^{36–38,40,41,44,45,47,49,52,53,55,57,59,61–63} reported favorable results on the basis of direction (i.e., decrease) of the effect estimate. There was no difference when stratified by length of follow-up time ≤6 months or >6 months or whether the intervention included a bystander approach. Table 1 lists the strategy combinations employed in addition to providing information (teaching healthy relationship skills, promoting social norms that protect against violence, and creating protective environments) that were favorable and consistent along with examples of interventions. Strategy combinations not included in Table 1 had inconsistent results across studies or too few studies to draw conclusions about perpetration. Effectiveness by combinations of strategies along with their corresponding approaches could not be determined.

Victimization.—A total of 18 arms from 15 studies^{36,38–41,44,45,47,52,53,56,58,60–62} reported on victimization. A total of 7 arms from 5 studies^{38,41,47,53,61} reported a median decrease in odds of victimization (OR=0.9, IQI=0.3, 1.0). A total of 12 arms reported data that could not be combined on a scatterplot: 7 arms^{36,40,44,45,52,53,62} reported decreases in victimization, 3 arms in 2 studies^{56,60} reported no change, and 2 arms^{39,58} reported increases in victimization. Overall, 11 of 18 arms^{36,38,40,41,44,45,47,52,53,61,62} reported favorable results on the basis of direction (decreased) of the effect estimate. There was no difference by length of follow-up time ≤6 months or >6 months. Table 1 provides a list of strategy combinations that were favorable and consistent. Strategy combinations not included in Table 1 had inconsistent results across studies or too few studies to draw conclusions about victimization. Effectiveness by combinations of strategies along with their corresponding approaches could not be determined.

Bystander action.—A total of 10 arms from 9 studies^{32,42,43,46,48,49,57,59} reported on bystander action. A total of 8 arms from 7 studies^{32,42,43,48,57,59} reported a median relative increase of 2.5% for bystander action (IQI= −5.5%, 22.3%). Two arms reported data that could not be combined to calculate a median. One arm²⁵ reported significant increases in the percentage of male undergraduate students who reported intervening behaviors (e.g., expressing disapproval when a peer is verbally abusive toward women, attempting to stop a peer who tries to be coercive or violent), and 1 arm⁴⁶ reported significant increases among first-year university students in helping behavior for a friend but no change in helping behavior for a stranger. Of the 10 arms, 6 arms^{42,46,48,49,59} reported favorable results on the basis of increased bystander action, 1 arm⁵⁷ reported no change, and 3 arms^{32,43} reported unfavorable results.

When stratified by the length of follow-up, 8 arms^{32,42,43,48,57,59} reported a median relative increase in bystander action of 17.9% (IQI=2.8%, 34.6%) within 6 months of completing the intervention (Appendix Figure 2, available online). However, 4 arms in 3 studies^{32,57,59} reported decreases 6 months after intervention completion. Interventions reporting solely on bystander action included strategies to promote social norms—specifically protecting

against violence through bystander education and empowerment, engaging men and boys as allies in prevention, or both (Table 1).

Applicability.—A total of 4 studies reported results by race or ethnicity,^{40,44,47,58} 4 reported results for low-SES populations,^{38,39,44,47} and 3 reported results for youth in high-risk settings.^{44,52,62} Of studies reporting results by race or ethnicity,^{40,44,47,58} 1 study⁴⁷ stratified results by race or ethnicity of the sample, 1 study⁴⁴ targeted Black or African American adolescent girls, and 2 studies^{40,58} targeted Hispanic adolescents (1 study⁴⁰ was exclusively Hispanic, and 1 study⁵⁸ was majority Hispanic). None of the included studies stratified bystander outcomes by race, ethnicity, or SES. Interventions had favorable results for reduced perpetration and victimization among Black students^{44,47} and mixed for both perpetration and victimization for Hispanic students.^{40,47,58} A total of 4 studies^{38,39,44,47} reported outcomes for low-SES populations measured as the majority of the population eligible for a free or reduced-price school lunch program, on public assistance, or with annual household income < \$10,000. All interventions had favorable results for reduced perpetration and victimization among low-SES populations. A total of 3 studies targeted youth in high-risk settings (i.e., youth in foster care system,⁶² pregnant and parenting adolescent girls,⁴⁴ or youth that had experienced violence as a perpetrator or victim⁵²) and reported favorable results, whereas 1 study that targeted youth that had experienced violence³⁹ reported unfavorable results for perpetration and victimization.

Results were consistently favorable for decreasing perpetration among high-school-aged youth and middle-school-aged youth, and results were mixed for perpetration among college-aged youth. A total of 8^{37,39–41,44,59,62,63} of 10 arms^{37,39–41,44,53,58,59,62,63} among high-school-aged youth were favorable for decreasing perpetration; of these, 1 arm⁵⁹ also reported favorable results for bystander action. A total of 6^{38,45,47,61,69} of 8 arms^{38,45,47,60,61,69} among middle-school-aged youth were favorable for decreasing perpetration; none measured bystander action. One study arm⁵² evaluated a program that started with middle-school students and followed them through high school. Results showed that the program was effective for decreasing perpetration and victimization at first follow-up (1 month) and remained effective as the students moved into high school (at 3-year follow-up). A total of 4^{36,49,55,57} of 6 arms^{36,49,54–57} among college-aged youth were favorable for decreasing perpetration; of these, 1 arm⁴⁹ also reported favorable results for bystander action.

DISCUSSION

Summary of Findings

This review found sufficient evidence that primary prevention interventions are effective in reducing the perpetration of IPV and SV among youth. Specifically, those interventions that used the following strategies were consistent and favorable across studies: (1) teaching healthy relationship skills, (2) promoting social norms that protect against violence, and (3) creating protective environments. Other strategy combinations had inconsistent results across studies or too few studies to draw conclusions about perpetration or victimization. In addition, interventions that promote social norms to protect against violence through

bystander education and empowerment, engage men and boys as allies in prevention, or do both were found to be effective in increasing bystander action in the short term. Two studies in the review^{49,70} that examined the effects of the intervention on both bystander action and perpetration reported favorable results for both outcomes, suggesting that increased bystander action may be associated with decreased perpetration. For studies that reported bystander action, intervention effects appeared to diminish over time, possibly indicating the need for booster sessions or extended interventions.

Similar to this *Community Guide* review, previous reviews reported findings that were favorable but often did not reach statistical significance on the effectiveness of interventions to prevent perpetration and victimization. Previous reviews and this *Community Guide* review highlighted the need for more interventions focused on creating protective environments, changing social norms, and equipping young people to safely intervene when they witness behaviors that can result in dating violence or SV.^{11,71,72} The *Community Guide* review differs because it systematically assesses heterogeneous data to identify effective combinations of intervention strategies that can help inform decision makers regarding the best intervention to implement for their population. In addition, the findings from this review provide the basis for a Community Preventive Services Task Force recommendation for primary prevention interventions to prevent or reduce the perpetration of IPV or SV among youth.⁷³

The review team conducted a bridge search in November 2020 to identify studies published after this review's search period. Two systematic reviews that focused on bystander interventions reported similar results, although inclusion criteria differed slightly.^{74,75} Two systematic reviews focused on dating violence prevention among adolescents in high- and low-income countries.^{76,77} Findings from studies in high-income countries aligned with this *Community Guide* review. Each review also reported the need for more research identifying the specific combination of strategies or components that work together to prevent or reduce dating violence. One additional study not captured in either review reported promising results for the effectiveness of a dating violence program on reducing any SV among middle-school youth.⁷⁸

Evidence Gaps

Included studies consisted of various combinations of intervention strategies and their corresponding approaches, making it difficult to determine which combinations were the most effective, and many combinations included too few studies to draw any conclusions. Specifically, more studies are needed that evaluate interventions aimed at creating protective environments, such as policy change in health, economic, educational, and social sectors. The review also lacked studies that were conducted in rural settings or among youth identifying as 2 spirit, lesbian, gay, bisexual, transgender, queer, intersex, asexual, or questioning or youth with intellectual or developmental disabilities. Moreover, better consensus on the best scales to use to increase comparability across studies and increase the ability to synthesize evidence is needed.⁷⁹ More studies are needed that measure morbidity-related outcomes, which were reported by 1 study⁶² in this review. Finally, participants in included studies in middle schools reported violence at baseline,^{38,45,47,60,61,69} suggesting

that some students are experiencing and perpetrating IPV and SV in or before middle school. Therefore, age-appropriate interventions for elementary school students may need to be developed and tested for immediate and later impact on IPV and SV outcomes.

Limitations

This review has several limitations. First, included articles were from peer-reviewed literature; therefore, there is potential publication bias. However, not all published studies found favorable effects. Second, outcomes are based on self-reported data; therefore, there is potential for recall bias and social desirability bias. However, the included studies used validated scales to minimize these biases. Finally, owing to the use of numerous different outcome measures for perpetration and victimization, many studies could not be combined into pooled estimates. Instead, systematic methods were developed and used to explore effectiveness across these highly heterogeneous, self-reported data by requiring 75% of studies for each outcome to show a consistent effect in the favorable direction, regardless of statistical significance.

Conclusions

Adolescence is a critical time to promote attitudes and behaviors to prevent violence. Primary prevention interventions to prevent or reduce IPV and SV can be effective ways to decrease the perpetration of both IPV and SV among youth aged 12–24 years as well as to increase bystander action in the short term. Findings from this review can inform researchers, school personnel, public health decision makers, and parents and other caregivers about effective strategies to prevent violence among youth.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

The authors acknowledge staff from the Centers for Disease Control and Prevention library for assistance with the systematic literature search, and Carrie Mulford (National Institute of Justice) and Cierra Olivia Thomas-Williams (Indiana Coalition Against Domestic Violence) for their participation on the coordination team and assistance in conceptualizing the review.

The findings and conclusions in this paper are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention/the Agency for Toxic Substances and Disease Registry. The content expressed in this paper is solely the responsibility of the authors and does not necessarily represent the official views of NIH or HHS.

The work of DLOS was supported with funds from the Oak Ridge Institute for Science and Education. This review was partially supported by funds from the HHS Office of Women's Health. Patrick Remington is an AJPM Editor and was not involved in the peer review protocol for this manuscript.

REFERENCES

1. The National Intimate Partner and Sexual Violence Survey (NISVS). National Center for Injury Prevention and Control, Centers for Disease Control and Prevention. <https://www.cdc.gov/violenceprevention/data-sources/nisvs/index.html>. Updated July 19, 2021. Accessed February 1, 2018.

2. Breiding MJ, Basile BK, Smith SG, Black MC, Mahendra RR. Intimate partner violence surveillance: uniform definitions and recommended data elements, version 2.0. Atlanta, GA: National Center for Injury Prevention and Control, Centers for Disease Control and Prevention; 2015. <https://www.cdc.gov/violenceprevention/pdf/ipv/intimatepart-nerviolence.pdf>. Published 2015. Accessed February 1, 2018.
3. Basile KC, Smith SS, Breiding MJ, Black MC, Mahendra RR. Sexual violence surveillance: uniform definitions and recommended data elements, version 2.0. Atlanta, GA: National Center for Injury Prevention and Control, Centers for Disease Control and Prevention; Published 2014. https://www.cdc.gov/violenceprevention/pdf/sv_surveillance_-_definitions-2009-a.pdf. Accessed February 1, 2018.
4. Basile KC, Clayton HB, DeGue S, et al. Interpersonal violence victimization among high school students - Youth Risk Behavior Survey, United States, 2019. *MMWR Suppl.* 2020;69(1):28–37. 10.15585/mmwr.su6901a4. [PubMed: 32817605]
5. Eaton AA, Stephens DP. Chapter 10 - adolescent dating violence among ethnically diverse youth. In: Wolfe DA, Temple JR, eds. *Adolescent Dating Violence: Theory, Research, and Prevention*. Miami, FL: Elsevier Academic Press, 2018:233–260. 10.1016/b978-0-12-811797-2.00010-4.
6. Exner-Cortens D, Eckenrode J, Rothman E. Longitudinal associations between teen dating violence victimization and adverse health outcomes. *Pediatrics.* 2013;131(1):71–78. 10.1542/peds.2012-1029. [PubMed: 23230075]
7. Offenauer P, Buchalter A. Teen dating violence: a literature review and annotated bibliography. Washington, DC: Federal Research Division Library of Congress. <https://www.ncjrs.gov/pdffiles1/nij/grants/235368.pdf>. Published July 2011. Accessed June 14, 2021.
8. Cox PJ, Ortega S, Cook-Craig PG, Conway P. Strengthening systems for the primary prevention of intimate partner violence and sexual violence: CDC's DELTA and EMPOWER Programs. *J Fam Soc Work.* 2010;13(4):287–296. 10.1080/10522158.2010.492565.
9. DeGue S, Simon TR, Basile KC, Yee SL, Lang K, Spivak H. Moving forward by looking back: reflecting on a decade of CDC's work in sexual violence prevention, 2000–2010. *J Womens Health (Larchmt).* 2012;21(12):1211–1218. 10.1089/jwh.2012.3973. [PubMed: 23140201]
10. McMahon PM. The public health approach to the prevention of sexual violence. *Sex Abuse.* 2000;12(1):27–36. 10.1177/107906320001200104. [PubMed: 10729957]
11. Lundgren R, Amin A. Addressing intimate partner violence and sexual violence among adolescents: emerging evidence of effectiveness. *J Adolesc Health.* 2015;56(1 suppl):S42–S50. 10.1016/j.jadohealth.2014.08.012. [PubMed: 25528978]
12. Banyard VL. *Toward the Next Generation of Bystander Prevention of Sexual and Relationship Violence: Action Coils to Engage Communities*. Heidelberg, Germany: Springer, 2015. 10.1007/978-3-319-23171-6.
13. Basile KC, DeGue S, Jones K, et al. STOP SV: a technical package to prevent sexual violence. Atlanta, GA: National Center for Injury Prevention and Control, Centers for Disease Control and Prevention; Published 2016. <https://www.cdc.gov/violenceprevention/pdf/sv-prevention-technical-package.pdf>. Accessed February 1, 2018.
14. McCauley HL, Tancredi DJ, Silverman JG, et al. Gender-equitable attitudes, bystander behavior, and recent abuse perpetration against heterosexual dating partners of male high school athletes. *Am J Public Health.* 2013;103(10):1882–1887. 10.2105/AJPH.2013.301443. [PubMed: 23947324]
15. Niolon PH, Kearns M, Dills J, et al. Preventing intimate partner violence across the lifespan: a technical package of programs, policies, and practices. Atlanta, GA: National Center for Injury Prevention and Control, Centers for Disease Control and Prevention; 2017. <https://www.cdc.gov/violenceprevention/pdf/ipv-technicalpackages.pdf>. Published 2017. Accessed February 1, 2018.
16. DeGue S, Valle LA, Holt MK, Massetti GM, Matjasko JL, Tharp AT. A systematic review of primary prevention strategies for sexual violence perpetration. *Aggress Violent Behav.* 2014;19(4):346–362. 10.1016/j.avb.2014.05.004. [PubMed: 29606897]
17. Whitaker DJ, Murphy CM, Eckhardt CI, Hodges AE, Cowart M. Effectiveness of primary prevention efforts for intimate partner violence. *Partner Abuse.* 2013;4(2):175–195. 10.1891/1946-6560.4.2.175.

18. Briss PA, Zaza S, Pappaioanou M, et al. Developing an evidence-based Guide to Community Preventive Services—methods. The Task Force on Community Preventive Services. *Am J Prev Med.* 2000;18(1 suppl):35–43. 10.1016/S0749-3797(99)00119-1. [PubMed: 10806978]
19. Recognizing adolescence. WHO. <https://apps.who.int/adolescent/second-decade/section2/page1/recognizing-adolescence.html>. Accessed August 10, 2021.
20. United Nations Development Programme. Human development report 2016: human development for everyone. New York, NY: United Nations Development Programme. http://hdr.undp.org/sites/default/files/2016_human_development_report.pdf. Published 2016. Accessed April 3, 2019.
21. Wolfe DA, Scott K, Reitzel-Jaffe D, Wekerle C, Grasley C, Straatman AL. Development and validation of the Conflict in Adolescent Dating Relationships Inventory. *Psychol Assess.* 2001;13(2):277–293. 10.1037/1040-3590.13.2.277. [PubMed: 11433803]
22. Prinstein MJ, Boergers J, Spirito A, Little TD, Grapentine WL. Peer functioning, family dysfunction, and psychological symptoms in a risk factor model for adolescent inpatients' suicidal ideation severity. *J Clin Child Psychol.* 2000;29(3):392–405. 10.1207/S15374424JCCP2903_10. [PubMed: 10969423]
23. Prinstein MJ, Boergers J, Vernberg EM. Overt and relational aggression in adolescents: social-psychological adjustment of aggressors and victims. *J Clin Child Psychol.* 2001;30(4):479–491. 10.1207/S15374424JCCP3004_05. [PubMed: 11708236]
24. Vernberg EM, Jacobs AK, Hershberger SL. Peer victimization and attitudes about violence during early adolescence. *J Clin Child Psychol.* 1999;28(3):386–395. 10.1207/S15374424jccp280311. [PubMed: 10446688]
25. Straus MA, Hamby SL, Boney-McCoy S, Sugarman DB. The revised Conflict Tactics Scales (CTS2): development and preliminary psychometric data. *J Fam Issues.* 1996;17(3):283–316. 10.1177/019251396017003001.
26. Foshee VA, Bauman KE, Arriaga XB, Helms RW, Koch GG, Linder GF. An evaluation of Safe Dates, an adolescent dating violence prevention program. *Am J Public Health.* 1998;88(1):45–50. 10.2105/AJPH.88.1.45. [PubMed: 9584032]
27. Johnson SM, Murphy MJ, Gidycz CA. Reliability and validity of the Sexual Experiences Survey-Short Forms Victimization and Perpetration. *Violence Vict.* 2017;32(1):78–92. 10.1891/0886-6708.VV-D-15-00110. [PubMed: 28234199]
28. Koss MP, Abbey A, Campbell R, et al. Revising the SES: a collaborative process to improve assessment of sexual aggression and victimization. *Psychol Women Q.* 2007;31(4):357–370. 10.1111/j.1471-6402.2007.00385.x.
29. Koss MP, Gidycz CA. Sexual experiences survey: reliability and validity. *J Consult Clin Psychol.* 1985;53(3):422–423. 10.1037/0022-006X.53.3.422. [PubMed: 3874219]
30. Espelage DL, Low S, De La Rue L. Relations between peer victimization subtypes, family violence, and psychological outcomes during early adolescence. *Psychol Violence.* 2012;2(4):313–324. 10.1037/a0027386.
31. Banyard VL, Moynihan MM, Cares AC, Warner R. How do we know if it works? Measuring outcomes in bystander-focused abuse prevention on campuses. *Psychol Violence.* 2014;4(1):101–115. 10.1037/a0033470.
32. Banyard VL, Moynihan MM, Plante EG. Sexual violence prevention through bystander education: an experimental evaluation. *J Commun Psychol.* 2007;35(4):463–481. 10.1002/jcop.20159.
33. Bruner JB. Measuring Rape-Supportive Attitudes, Behaviors, and Perceived Peer Norms Among College Students: Validation of a Social Norms Survey[dissertation]. Greeley: University of Northern Colorado, 2002.
34. Loh C, Gidycz CA, Lobo TR, Luthra RA. A prospective analysis of sexual assault perpetration: risk factors related to perpetrator characteristics. *J Interpers Violence.* 2005;20(10):1325–1348. 10.1177/0886260505278528. [PubMed: 16162492]
35. Zaza S, Wright-De Agüero LK, Briss PA, et al. Data collection instrument and procedure for systematic reviews in the Guide to Community Preventive Services. *Am J Prev Med.* 2000;18(1 suppl 1):44–74. 10.1016/S0749-3797(99)00122-1. [PubMed: 10806979]

36. Coker AL, Bush HM, Fisher BS, et al. Multi-college bystander intervention evaluation for violence prevention. *Am J Prev Med.* 2016;50 (3):295–302. 10.1016/j.amepre.2015.08.034. [PubMed: 26541099]
37. de Graaf I, de Haas S, Zaagsma M, Wijzen C. Effects of Rock and Water: an intervention to prevent sexual aggression. *J Sex Aggress.* 2016;22(1):4–19. 10.1080/13552600.2015.1023375.
38. Espelage DL, Low S, Polanin JR, Brown EC. Clinical trial of Second Step middle-school program: impact on aggression & victimization. *J Appl Dev Psychol.* 2015;37:52–63. 10.1016/j.appdev.2014.11.007.
39. Foshee VA, Benefield T, Dixon KS, et al. The effects of Moms and Teens for Safe Dates: a dating abuse prevention program for adolescents exposed to domestic violence. *J Youth Adolesc.* 2015;44(5):995–1010. 10.1007/s10964-015-0272-6. [PubMed: 25776110]
40. Gonzalez-Guarda RM, Guerra JE, Cummings AA, Pino K, Becerra MM. Examining the preliminary efficacy of a dating violence prevention program for Hispanic adolescents. *J Sch Nurs.* 2015;31(6):411–421. 10.1177/1059840515598843. [PubMed: 26260135]
41. Joppa MC, Rizzo CJ, Nieves AV, Brown LK. Pilot investigation of the Katie Brown educational program: a school-community partnership. *J Sch Health.* 2016;86(4):288–297. 10.1111/josh.12378. [PubMed: 26930241]
42. Jouriles EN, McDonald R, Rosenfield D, et al. TakeCARE, a video bystander program to help prevent sexual violence on college campuses: results of two randomized, controlled trials. *Psychol Violence.* 2016;6(3):410–420. 10.1037/vio0000016. [PubMed: 27867694]
43. Kleinsasser A, Jouriles EN, McDonald R, Rosenfield D. An online bystander intervention program for the prevention of sexual violence. *Psychol Violence.* 2015;5(3):227–235. 10.1037/a0037393. [PubMed: 26240776]
44. Langhinrichsen-Rohling J, Turner LA. The efficacy of an intimate partner violence prevention program with high-risk adolescent girls: a preliminary test. *Prev Sci.* 2012;13(4):384–394. 10.1007/s11121-011-0240-7. [PubMed: 21842333]
45. Miller S, Williams J, Cutbush S, Gibbs D, Clinton-Sherrod M, Jones S. Evaluation of the Start Strong initiative: preventing teen dating violence and promoting healthy relationships among middle school students. *J Adolesc Health.* 2015;56(2 suppl 2):S14–S19. 10.1016/j.jadohealth.2014.11.003.
46. Moynihan MM, Banyard VL, Cares AC, Potter SJ, Williams LM, Stapleton JG. Encouraging responses in sexual and relationship violence prevention: what program effects remain 1 year later? *J Interpers Violence.* 2015;30(1):110–132. 10.1177/0886260514532719. [PubMed: 24850763]
47. Peskin MF, Markham CM, Shegog R, Baumler ER, Addy RC, Tortolero SR. Effects of the it's your game... keep it real program on dating violence in ethnic-minority middle school youths: a group randomized trial. *Am J Public Health.* 2014;104(8):1471–1477. 10.2105/AJPH.2014.301902. [PubMed: 24922162]
48. Peterson K, Sharps P, Banyard V, et al. An evaluation of two dating violence prevention programs on a college campus. *J Interpers Violence.* 2018;33(23):3630–3655. 10.1177/0886260516636069. [PubMed: 26976433]
49. Salazar LF, Vivolo-Kantor A, Hardin J, Berkowitz A. A web-based sexual violence bystander intervention for male college students: randomized controlled trial. *J Med Internet Res.* 2014;16(9):e203. 10.2196/jmir.3426. [PubMed: 25198417]
50. Borsky AE, McDonnell K, Turner MM, Rimal R. Raising a red flag on dating violence: evaluation of a low-resource, college-based bystander behavior intervention program. *J Interpers Violence.* 2018;33 (22):3480–3501. 10.1177/0886260516635322. [PubMed: 26965908]
51. Daigneault I, Hébert M, McDuff P, et al. Effectiveness of a sexual assault awareness and prevention workshop for youth: A 3-month follow-up pragmatic cluster randomization study. *Can J Hum Sex.* 2015;24(1):19–30. 10.3138/cjhs.2626.
52. Foshee VA, Bauman KE, Ennett ST, Suchindran C, Benefield T, Linder GF. Assessing the effects of the dating violence prevention program “Safe Dates” using random coefficient regression modeling. *Prev Sci.* 2005;6(3):245–258. 10.1007/s11121-005-0007-0. [PubMed: 16047088]

53. Foshee VA, McNaughton Reyes HL, Ennett ST, Cance JD, Bauman KE, Bowling JM. Assessing the effects of Families for Safe Dates, a family-based teen dating abuse prevention program. *J Adolesc Health*. 2012;51(4):349–356. 10.1016/j.jadohealth.2011.12.029. [PubMed: 22999835]
54. Foubert JD. The longitudinal effects of a rape-prevention program on fraternity men's attitudes, behavioral intent, and behavior. *J Am Coll Health*. 2000;48(4):158–163. 10.1080/07448480009595691. [PubMed: 10650733]
55. Foubert JD, Newberry JT, Tatum J. Behavior differences seven months later: effects of a rape prevention program. *J Stud Aff Res Pract*. 2007;44(4):728–749. 10.2202/1949-6605.1866.
56. Gidycz CA, Layman MJ, Rich CL, et al. An evaluation of an acquaintance rape prevention program: impact on attitudes, sexual aggression, and sexual victimization. *J Interpers Violence*. 2001;16(11):1120–1138. 10.1177/088626001016011002.
57. Gidycz CA, Orchowski LM, Berkowitz AD. Preventing sexual aggression among college men: an evaluation of a social norms and bystander intervention program. *Violence Against Women*. 2011;17(6):720–742. 10.1177/1077801211409727. [PubMed: 21571742]
58. Jaycox LH, McCaffrey D, Eiseman B, et al. Impact of a school-based dating violence prevention program among Latino teens: randomized controlled effectiveness trial. *J Adolesc Health*. 2006;39(5):694–704. 10.1016/j.jadohealth.2006.05.002. [PubMed: 17046506]
59. Miller E, Tancredi DJ, McCauley HL, et al. One-year follow-up of a coach-delivered dating violence prevention program: a cluster randomized controlled trial. *Am J Prev Med*. 2013;45(1):108–112. 10.1016/j.amepre.2013.03.007. [PubMed: 23790995]
60. Taylor B, Stein N, Burden F. The effects of gender violence/harassment prevention programming in middle schools: a randomized experimental evaluation. *Violence Vict*. 2010;25(2):202–223. 10.1891/0886-6708.25.2.202. [PubMed: 20514817]
61. Taylor BG, Stein ND, Mumford EA, Woods D. Shifting Boundaries: an experimental evaluation of a dating violence prevention program in middle schools. *Prev Sci*. 2013;14(1):64–76. 10.1007/s11121-012-0293-2. [PubMed: 23076726]
62. Wolfe DA, Wekerle C, Scott K, Straatman AL, Grasley C, Reitzel-Jaffe D. Dating violence prevention with at-risk youth: a controlled outcome evaluation. *J Consult Clin Psychol*. 2003;71(2):279–291. 10.1037/0022-006x.71.2.279. [PubMed: 12699022]
63. Wolfe DA, Crooks C, Jaffe P, et al. A school-based program to prevent adolescent dating violence: a cluster randomized trial. *Arch Pediatr Adolesc Med*. 2009;163(8):692–699. 10.1001/archpediatrics.2009.69. [PubMed: 19652099]
64. Stephens KA, George WH. Rape prevention with college men: evaluating risk status. *J Interpers Violence*. 2009;24(6):996–1013. 10.1177/0886260508319366. [PubMed: 18591366]
65. Foshee VA, Bauman KE, Greene WF, Koch GG, Linder GF, MacDougall JE. The Safe Dates program: 1-year follow-up results. *Am J Public Health*. 2000;90(10):1619–1622. 10.2105/ajph.90.10.1619. [PubMed: 11029999]
66. Foshee VA, Linder GF, Bauman KE, et al. The Safe Dates Project: theoretical basis, evaluation design, and selected baseline findings. *Am J Prev Med*. 1996;12(5 suppl):39–47. 10.1016/S0749-3797(18)30235-6.
67. Coker AL, Cook-Craig PG, Williams CM, et al. Evaluation of Green Dot: an active bystander intervention to reduce sexual violence on college campuses. *Violence Against Women*. 2011;17(6):777–796. 10.1177/1077801211410264. [PubMed: 21642269]
68. Coker AL, Fisher BS, Bush HM, et al. Evaluation of the Green Dot bystander intervention to reduce interpersonal violence among college students across three campuses. *Violence Against Women*. 2015;21(12):1507–1527. 10.1177/1077801214545284. [PubMed: 25125493]
69. Espelage DL, Low S, Polanin JR, Brown EC. The impact of a middle school program to reduce aggression, victimization, and sexual violence. *J Adolesc Health*. 2013;53(2):180–186. 10.1016/j.jadohealth.2013.02.021. [PubMed: 23643338]
70. Miller E, Tancredi DJ, McCauley HL, et al. Coaching boys into men”: a cluster-randomized controlled trial of a dating violence prevention program. *J Adolesc Health*. 2012;51(5):431–438. 10.1016/j.jadohealth.2012.01.018. [PubMed: 23084163]

71. De Koker P, Mathews C, Zuch M, Bastien S, Mason-Jones AJ. A systematic review of interventions for preventing adolescent intimate partner violence. *J Adolesc Health*. 2014;54(1):3–13. 10.1016/j.jadohealth.2013.08.008. [PubMed: 24125727]
72. De La Rue L, Polanin JR, Espelage DL, Pigott TD. A meta-analysis of school-based interventions aimed to prevent or reduce violence in teen dating relationships. *Rev Educ Res*. 2017;87(1):7–34. 10.3102/0034654316632061.
73. Violence prevention: primary prevention interventions to reduce perpetration of intimate partner violence and sexual violence among youth. Community Preventive Services Task Force. <https://www.the-communityguide.org/findings/violence-prevention-primary-prevention-interventions-reduce-perpetration-intimate-partner-violence/sexual-violence-among-youth>. Updated February 17, 2021. Accessed June 14, 2021.
74. Storer HL, Casey E, Herrenkohl T. Efficacy of bystander programs to prevent dating abuse among youth and young adults: a review of the literature. *Trauma Violence Abuse*. 2016;17(3):256–269. 10.1177/1524838015584361. [PubMed: 25951840]
75. Mujal GN, Taylor ME, Fry JL, Gochez-Kerr TH, Weaver NL. A systematic review of bystander interventions for the prevention of sexual violence. *Trauma Violence Abuse*. 2021;22(2):381–396. 10.1177/1524838019849587. [PubMed: 31204606]
76. Lee C, Wong JS. Examining the effects of teen dating violence prevention programs: a systematic review and meta-analysis. *J Exp Criminol*. Online September 25, 2020 In press. 10.1007/s1129-2020-09442-x.
77. McNaughton Reyes HL, Graham LM, Chen MS, et al. Adolescent dating violence prevention programmes: a global systematic review of evaluation studies. *Lancet Child Adolesc Health*. 2021;5(3):223–232. 10.1016/S2352-4642(20)30276-5. [PubMed: 33220790]
78. DeGue S, Niolon PH, Estefan LF, et al. Effects of Dating Matters® on sexual violence and sexual harassment outcomes among middle school youth: a cluster-randomized controlled trial [published correction appears in *Prev Sci*. 2020;22(2):186]. *Prev Sci*. 2021;22(2):175–185. 10.1007/s11121-020-01152-0. [PubMed: 32844328]
79. Fellmeth GL, Heffernan C, Nurse J, Habibula S, Sethi D. Educational and skills-based interventions for preventing relationship and dating violence in adolescents and young adults. *Cochrane Database Syst Rev*. 2013(6):CD004534. 10.1002/14651858.CD004534.pub3. [PubMed: 23780745]




Figure 1.
PRISMA flowchart.

Table 1.

Summary of Strategy Combinations That Had Favorable and Consistent Results

Strategy	Result	Example
Teach healthy relationship skills	Perpetration: 5 study arms ^{37,44,47,61,69} 3 measures of effect, Favorable effects: 4 study arms ^{37,44,47,69} (80%) Victimization: 4 study arms ^{44,47,61,69} 3 measures of effect, Favorable effects: 3 study arms ^{44,47,69} (75%)	Exercises in social resilience aimed at body language, feeling, setting and respecting boundaries, intuition, making contact, standing up for oneself, and communication skills. Conflict management skills for dating.
Promote social norms that protect against violence	Perpetration: 3 study arms ^{49,57,59} 3 measures of effect Favorable effects: 3 study arms ^{49,57,59} (100%)	Web portal modules that include interactivity, didactic activities, and episodes of a serial drama Bystander education and empowerment.
Teach healthy relationship skills + Promote social norms that protect against violence	Victimization: 8 study arms ^{36,39-41,52-53,61-62} 3 measures of effect Favorable effects: 6 study arms ^{36,52-53,40-41,62} (75%)	Socioemotional learning programs to teach healthy dating skills (conflict resolution). Interactive activities that address dating violence norms, gender stereotyping, conflict resolution.
Teach healthy relationship skills + Promote social norms that protect against violence + Create protective environments	Perpetration: 2 study arms ^{45,61} 2 measures of effect Favorable effects: 2 study arms ^{45,61} (100%) Victimization: 2 study arms ^{45,61} 2 measures of effect Favorable effects: 2 study arms ^{45,61} (100%)	Identification of hotspots coupled with an increase in staff presence in those areas. Social marketing strategies. School-based teen dating violence prevention curricula to enhance skills and attitudes consistent with promotion of healthy relationships and reduction of teen dating violence.
Promote social norms that protect against violence	Bystander Action <6 months of completing the intervention: 8 ^a study arms ^{32,42-43,48,57,59} 2 measures of effect Favorable effects: 7 ^a study arms ^{32,42-43,57,59} (88%)	Protecting against violence through bystander education and empowerment Engaging men or boys as allies in prevention.

Note: Percentages represent the proportion of favorable studies for each strategy combination; favorable effects included studies with statistically significant effects ($p < 0.05$).

^aStudies included multiple study arms.