

COVID-19 Vaccine

Interim COVID-19 Immunization Schedule for 6 Months of Age and Older



The table below provides guidance for COVID-19 vaccination schedules based on age and medical condition. Scheduling considerations include:

- Administer the appropriate vaccine product based on the recipient's age and the product's age indications.
- COVID-19 vaccines may be administered on the same day as other vaccines.
- Doses administered at any time after the intervals outlined below are valid.

Detailed information can be found in CDC's Interim Clinical Considerations for Use of COVID-19 Vaccines Currently Approved or Authorized in the United States, see: www.cdc.gov/vaccines/covid-19/clinical-considerations/covid-19-vaccines-us.html

Table 1. Immunization Schedule for Children 6 Months through 17 Years of Age

Type	Product*	Recipient Age	For Most People		Those Who ARE Moderately or Severely Immunocompromised	
			Doses	Interval Between Doses ^{†‡}	Doses	Interval Between Doses ^{†‡}
mRNA vaccine	Moderna (Blue vial cap with magenta-bordered label)	6 months through 5 years	Total doses: 2 doses		Total doses: 3 doses	
			Dose 1 to 2	At least 4–8 weeks [‡]	Dose 1 to 2	At least 4 weeks
					Dose 2 to 3	At least 4 weeks
	Pfizer-BioNTech (Maroon vial cap with maroon-bordered label)	6 months through 4 years	Total number: 3 doses		Total number: 3 doses	
			Dose 1 to 2	At least 3–8 weeks [‡]	Dose 1 to 2	At least 3 weeks
			Doses 2 and 3	At least 8 weeks	Dose 2 to 3	At least 8 weeks
	Pfizer-BioNTech (Orange vial cap with orange-bordered label)	5 through 11 years	Total number: 3 doses		Total number: 4 doses	
			Dose 1 to 2	At least 3–8 weeks [‡]	Dose 1 to 2	At least 3 weeks
			Dose 2 to 3	At least 5 months	Dose 2 to 3	At least 4 weeks
					Dose 3 to 4	At least 3 months
	Pfizer-BioNTech (Purple vial cap with a purple-bordered label or gray vial cap with gray-bordered label)	12 years through 17 years	Total number: 3 doses		Total number: 5 doses	
			Dose 1 to 2	At least 3-8 weeks [‡]	Dose 1 to 2	At least 3 weeks
			Dose 2 to 3	At least 5 months	Dose 2 to 3	At least 4 weeks
					Dose 3 to 4	At least 3 months
					Dose 4 to 5	At least 4 months

* Complete the primary series with same product. If the vaccine product previously administered cannot be determined or is no longer available, any age-appropriate mRNA COVID-19 vaccine product may be administered at least 28 days after the first dose. Any COVID-19 vaccine product (age appropriate) may be administered for a booster dose. It does not need to be the same product used for the primary series.

[†] Persons with a recent SARS-CoV-2 infection may consider delaying a primary series or booster dose by 3 months from symptom onset or positive test (if infection was asymptomatic).

[‡] Some studies in adolescents and adults have shown the small risk of myocarditis associated with mRNA COVID-19 vaccines might be reduced and peak antibody responses and vaccine effectiveness may be increased with an interval longer than 4 weeks. An 8-week interval may be optimal for people who are not moderately or severely immunocompromised and ages 5–64 years, especially for males ages 12–39 years.

COVID-19 Vaccine

Interim COVID-19 Immunization Schedule
for 6 Months of Age and Older



Table 2. Immunization Schedule for Persons 18 Years of Age

Type	Product*	Recipient Age	For Most People		Those Who ARE Moderately or Severely Immunocompromised	
			Doses	Interval Between Doses ^{†‡}	Doses	Interval Between Doses ^{†‡}
mRNA vaccine	Moderna (Red vial cap with a blue-bordered label)	18 years and older	Total number: 3 or 4 doses		Total number: 5 doses	
			Dose 1 to 2	At least 4–8 weeks [‡]	Dose 1 to 2	At least 4 weeks
			Dose 2 to 3	At least 5 months	Dose 2 to 3	At least 4 weeks
			Dose 3 to 4	At least 4 months for persons ages 50 years and older	Dose 3 to 4	At least 3 months
					Dose 4 to 5	At least 4 months
	Pfizer-BioNTech (Purple vial cap with a purple-bordered label or gray vial cap with gray-bordered label)	18 years and older	Total number: 3 or 4 doses		Total number: 5 doses	
			Dose 1 to 2	At least 3-8 weeks [‡]	Dose 1 to 2	At least 3 weeks
			Dose 2 to 3	At least 5 months	Dose 2 to 3	At least 4 weeks
			Dose 3 to 4	At least 4 months for persons ages 50 years and older	Dose 3 to 4	At least 3 months
					Dose 4 to 5	At least 4 months
Viral vector vaccine	Janssen [§]	18 years and older	Total number: 2 or 3 doses		Total number: 4 doses	
			Dose 1 to 2	At least 8 weeks	Dose 1 to 2	At least 4 weeks (mRNA vaccine)
			Dose 2 to 3	At least 4 months for persons ages 50 years and older (mRNA vaccine) [¶]	Dose 2 to 3	At least 8 weeks (mRNA vaccine)
					Dose 3 to 4	At least 4 months (mRNA vaccine) ^{**}

* Complete the primary series with same product. If the vaccine product previously administered cannot be determined or is no longer available, any age-appropriate mRNA COVID-19 vaccine product may be administered at least 28 days after the first dose. Any COVID-19 vaccine product (age appropriate) may be administered for a booster dose. It does not need to be the same product used for the primary series. An mRNA COVID-19 vaccine is preferred.

† Persons with a recent SARS-CoV-2 infection may consider delaying a primary series or booster dose by 3 months from symptom onset or positive test (if infection was asymptomatic).

‡ Some studies in adolescents and adults have shown the small risk of myocarditis associated with mRNA COVID-19 vaccines might be reduced and peak antibody responses and vaccine effectiveness may be increased with an interval longer than 4 weeks. An 8-week interval may be optimal for people who are not moderately or severely immunocompromised and ages 5–64 years, especially for males ages 12–39 years.

§ Age-appropriate mRNA COVID-19 vaccines are preferred over the Janssen COVID-19 Vaccine for all vaccine doses for all vaccine-eligible people 18 years of age and older. Janssen COVID-19 vaccine should only be used in limited situations. See: www.cdc.gov/vaccines/covid-19/clinical-considerations/covid-19-vaccines-us.html

¶ People ages 18–49 years: Those who received Janssen COVID-19 Vaccine as both their primary series dose and booster dose may receive a second booster dose using an mRNA COVID-19 vaccine at least 4 months after the Janssen booster dose.

** mRNA vaccine must be used.

COVID-19 Vaccine

Interim COVID-19 Immunization Schedule
for 6 Months of Age and Older



Table 3. COVID-19 Vaccine Products Summary

Product*	Age Indications†	Diluent	Dosage (amount injected)	
Type: mRNA vaccine				
Moderna (Blue vial cap with magenta-bordered label)	6 months through 5 years	NONE	Any dose in the series	0.25 mL
Moderna (Red vial cap with blue- bordered label)	18 years and older	NONE	Doses 1, 2 and 3	0.5 mL
			Doses 4 and 5	0.25 mL
Pfizer-BioNTech (Maroon vial cap with maroon-bordered label)	6 months through 4 years	2.2 mL 0.9% sodium chloride (normal saline, preservative-free)	Any dose in the series	0.2 mL
Pfizer-BioNTech (Orange vial cap with orange-bordered label)	5 through 11 years	1.3 mL 0.9% sodium chloride (normal saline, preservative-free)	Any dose in the series	0.2 mL
Pfizer-BioNTech (Purple vial cap with a purple-bordered label)	12 years and older	1.8 mL 0.9% sodium chloride (normal saline, preservative-free)	Any dose in the series	0.3 mL
Pfizer-BioNTech (Gray vial cap with a gray- bordered label)	12 years and older	NONE	Any dose in the series	0.3 mL
Product	Age Indications	Diluent	Dosage (amount injected)	
Type:Viral Vector Vaccine				
Janssen [‡] (Blue Cap)	18 years and older	NONE	Any dose in the series	0.5 mL

* Complete the primary series using the same product. Every effort should be made to determine which vaccine product was received as the first dose. If the vaccine product previously administered cannot be determined or is no longer available, any age-appropriate mRNA COVID-19 vaccine product may be administered at least 28 days after the first dose.

† Administer the appropriate vaccine product based on the recipient's age and the vaccine product's age indications.

‡ An age-appropriate mRNA COVID-19 vaccine is preferred over the Janssen COVID-19 Vaccine for all vaccine doses for all vaccine-eligible people 18 years of age and older. Janssen COVID-19 vaccine should only be used in limited situations. An mRNA vaccine must be used for the 2nd booster dose. See: www.cdc.gov/vaccines/covid-19/clinical-considerations/covid-19-vaccines-us.html

CDC Resources

CDC COVID-19 vaccine clinical training and materials at: www.cdc.gov/vaccines/covid-19/info-by-product/index.html

CDC Interim Clinical Considerations for the Use of COVID-19 Vaccines Currently Approved or Authorized in the United States at: www.cdc.gov/vaccines/covid-19/clinical-considerations/covid-19-vaccines-us.html#primary-series

CDC Vaccine administration clinical materials at: www.cdc.gov/vaccines/hcp/admin/resource-library.html

CDC Vaccine Storage and Handling Toolkit at: www.cdc.gov/vaccines/hcp/admin/storage/toolkit/index.html