

Supplementary Material

Supplemental Table 1. Sensitivity analysis of the effectiveness of the *PASOS SALUDABLES* intervention on body mass index (primary outcome)

a. With 3-levels nested clustering		
Factor	Beta (standard error)	p-value
Month	0.03 (0.01)	<0.0001
Treatment	0.13 (0.37)	0.72
Month*treatment	-0.01 (0.01)	0.29
b. Visit number as a categorical variable		
Factor	Beta (standard error)	p-value
Visit #1	-0.04 (0.09)	0.68
2	0.01 (0.09)	0.96
3	0.37 (0.09)	<0.0001
4 (vs. baseline)	0.57 (0.10)	<0.0001
Treatment	0.22 (0.38)	0.56
Visit #1*treatment	-0.20 (0.12)	0.11
#2*treatment	-0.30 (0.13)	0.02
#3*treatment	-0.29 (0.13)	0.02
#4*treatment	-0.10 (0.14)	0.50
Overall test (df=4) for visit*treatment yields p=0.10		
c. Excluding participants with HbA1C ≥ 6.5% at baseline		
Factor	Beta (standard error)	p-value
Month	0.03 (0.01)	<0.0001
Treatment	0.17 (0.39)	0.66
Month*treatment	-0.01 (0.01)	0.26
d. Gender adjustment		
Factor	Beta (standard error)	p-value
Month	0.03 (0.01)	<0.0001
Treatment	0.18 (0.37)	0.63
Month*treatment	-0.01 (0.01)	0.29
Female	1.49 (0.37)	<0.0001