

Supplementary Appendix S1—This table provides a brief review of infectious etiologic agent, human symptoms, route of transmission, and mortality rate of each abortion-associated zoonoses evaluated in survey of 469 large animal veterinarians in the United States from June 10, 2019, through October 31, 2019.

Infectious agent(s)	Human symptoms	Modes of transmission	References
Brucellosis			
<i>Brucella melitensis</i> <i>B. suis</i> <i>B. abortus</i> <i>B. canis</i>	Fever, malaise, myalgia, anorexia, headache, lethargy	Direct contact, fomite, aerosol, food- or water-borne	23-25
Q fever			
<i>Coxiella burnetii</i>	Fever, headache, diarrhea, vomiting, lethargy, myalgia, cough	Direct contact, fomite, aerosol, food or water- borne, vector-borne	26-28
Salmonellosis			
<i>Salmonella spp.</i>	Diarrhea, vomiting, abdominal pain, fever	Direct contact, fomite, food- or water-borne	29-31
Leptospirosis			
<i>Leptospira interrogans</i> (Multiple serovars)	Fever, headache, myalgia, abdominal pain, diarrhea, jaundice, cough, skin rash	Direct contact, fomite, food- or water-borne	35,36
Campylobacteriosis			
<i>Campylobacter jejuni</i> <i>C. fetus</i> subsp. <i>fetus</i> <i>C. fetus</i> subsp. <i>venerealis</i>	Diarrhea, vomiting, abdominal pain, fever, headache, myalgia	Direct contact, fomite, food- or water-borne	32-34