

SUPPLEMENTARY TABLE. Influenza-attributable pneumonia and influenza diagnoses based on patients with severe acute respiratory illness and a positive influenza test at the National Influenza Centre — Chile, January–August 2022

Age group (yrs)/month	% SARI patients with influenza-positive test	No. admissions with P&I discharge diagnosis	Influenza-attributable P&I diagnoses*
All ages			
Jan	2.75	1,940	53.4
Feb	1.52	1,845	28.1
Mar	2.79	1,828	51.0
Apr	7.04	1,975	139.0
May	8.26	3,086	255.0
Jun	6.93	4,609	319.4
Jul	6.39	2,359	150.8
Aug	5.17	110	5.7
Total	—	17,752	1,002
<5			
Jan	4.72	295	13.9
Feb	2.94	144	4.2
Mar	1.15	135	1.6
Apr	4.00	402	16.1
May	3.17	1,032	32.7
Jun	2.44	2,091	51.1
Jul	1.46	800	11.7
Aug	2.90	12	0.3
Total	—	4,911	132
5–18			
Jan	0	82	0
Feb	3.85	43	1.7
Mar	7.14	58	4.1
Apr	5.00	123	6.2
May	6.67	224	14.9
Jun	10.59	291	30.8
Jul	20.00	101	20.2
Aug	12.00	7	0.8
Total	—	929	79
19–64			
Jan	2.04	510	10.4
Feb	1.52	454	6.9
Mar	5.60	452	25.3
Apr	9.82	428	42.0
May	15.91	518	82.4
Jun	15.98	597	95.4
Jul	10.67	361	38.5
Aug	4.92	22	1.1
Total	—	3,342	302
≥65			
Jan	2.72	1,053	28.7
Feb	1.16	1,204	14.0
Mar	1.21	1,183	14.3
Apr	9.52	1,022	97.3

May	11.30	1,312	148.3
Jun	8.16	1,630	133.1
Jul	7.32	1,097	80.3
Aug	6.67	69	4.6
Total	—	8,570	521

Abbreviations: P&I = pneumonia and influenza; SARI = severe acute respiratory illness.

* Age-specific numbers in this column do not sum to total because of rounding when calculating influenza-attributable P&I diagnoses from P&I hospitalizations and SARI data.