

Appendix A. Effects of Homicide Exposure on Alternative Indicators of Pregnancy Desire, Estimated from Linear Models with Fixed-Effects

	Pronatalism (-5/5)	Antinatalism pole (0/1)
Longer-term		
Ever exposed in any prior week	0.11* (0.05)	-0.02** (0.01)
Constant	6.61 (7.27)	-0.90 (1.26)
Cumulative		
Previously exposed once	0.10+ (0.05)	-0.02* (0.01)
Previously exposed twice	-0.04 (0.09)	-0.03+ (0.02)
Previously exposed more than twice	0.46*** (0.12)	-0.06** (0.02)
Constant	6.78 (7.27)	-0.94 (1.26)
<i>Post-estimation tests of equality</i>		
Exposed 1x v. 2x		
Exposed 1x v. >=3x		*
Exposed 2x v. >=3x	+	+

Note: In all models N=14,715 weeks from 387 women. All models adjust for time-varying age and previous exposure to police-involved fatalities during the RDSL study. Models further include month and year dummies and their interactions. Standard errors are shown in parentheses.

+p<.1, *p<.05, **p<.01, ***p<.001 (two-tailed tests).

Appendix B. Effects of Homicide Exposures on Pregnancy Desires, by Weeks Since Initial Exposure

	Any desire for pregnancy (0/1)					Strongest desire to avoid pregnancy (0/1)				
	M1: <=10 weeks	M2: 11-20 weeks	M3: 21-30 weeks	M4: 31-40 weeks	M5: 41-50 weeks	M1: <=10 weeks	M2: 11-20 weeks	M3: 21-30 weeks	M4: 31-40 weeks	M5: 41- 50 weeks
Nearby homicides										
Exposed <=.25 Miles	-.004 (.01)	.04** (.01)	.02 (.01)	.03* (.01)	.07*** (.02)	.01 (.01)	-.02+ (.01)	-.02 (.01)	-.01 (.02)	-.06*** (.02)
Constant	1.33 (1.28)	1.45 (1.31)	1.49 (1.23)	1.05 (1.2)	1.31 (1.31)	-.65 (1.44)	-.76 (1.46)	-.82 (1.38)	-.38 (1.45)	-.58 (1.47)
Observations (weeks)	12,608	12,394	12,388	12,192	12,123	12,608	12,394	12,388	12,192	12,123
Women	386	383	383	382	380	386	383	383	382	380
Faraway homicides										
Exposed 1-1.25 Miles	-.01 (.01)	-.0002 (.01)	-.002 (.01)	-.004 (.01)	-.02+ (.01)	-.004 (.01)	-.002 (.01)	-.01 (.01)	-.01 (.01)	.0003 (.01)
Constant	1.78 (1.65)	2.49 (1.69)	2.84+ (1.72)	2.68 (1.60)	3.20 (1.74)	-.72 (1.80)	-1.04 (1.84)	-1.08 (1.86)	-1.54 (1.75)	-1.38 (1.92)
Observations (weeks)	8,170	7,648	7,489	7,356	7,210	8,170	7,648	7,489	7,356	7,210
Women	384	374	372	372	370	384	374	372	372	370

Note: All models adjust for time-varying age and previous exposure to police-involved fatalities during the RDSL study.

Models further include month and year dummies and their interactions. Standard errors are shown in parentheses.

+p<.1, *p<.05, **p<.01, ***p<.001 (two-tailed tests).

Appendix C. Effects of Homicide Exposure *Within a Half Mile* on Pregnancy Desire,
Estimated from Linear Models with Fixed-Effects

	Desire to become pregnant		Desire to avoid pregnancy		Conflicting desires
	Desire for pregnancy	Any desire for pregnancy	Desire to avoid pregnancy	Strongest desire to avoid pregnancy	Indifferent
	(0/5)	(0/1)	(0-5)	(0/1)	(0/1)
Longer-term					
Ever exposed in any prior week	0.02 (0.02)	0.002 (0.01)	0.04 (0.02)	0.001 (0.01)	-0.01+ (0.00)
Constant	7.05+ (3.85)	1.62 (1.12)	1.00 (4.28)	-0.84 (1.23)	-0.19 (0.68)
Cumulative					
Previously exposed 1x	0.01 (0.02)	-0.00 (0.01)	0.05+ (0.03)	0.01 (0.01)	-0.01* (0.00)
Previously exposed 2x	0.04 (0.03)	0.01 (0.01)	0.00 (0.03)	-0.02+ (0.01)	-0.00 (0.01)
Previously exposed >=3x	0.02 (0.03)	0.00 (0.01)	-0.06 (0.04)	-0.03* (0.01)	0.00 (0.01)
Constant	7.04+ (3.86)	1.62 (1.12)	0.51 (4.29)	-0.98 (1.23)	-0.14 (0.68)
<i>Post-estimation tests of equality</i>					
Exposed 1x v. 2x		*		**	
Exposed 1x v. >=3x			***	***	*
Exposed 2x v. >=3x			*		

Note: In all models N=14,715 weeks from 387 women. All models adjust for time-varying age and previous exposure to police-involved fatalities during the RDSL study. Models further include month and year dummies and their interactions. Standard errors are shown in parentheses.

+p<.1, *p<.05, **p<.01, ***p<.001 (two-tailed tests).