



**Supplemental Table 1.** Medical conditions among 1477 Medicaid-insured women with preexisting type 2 diabetes mellitus diagnosed with breast cancer in New York State, 2004 to 2016

Variables	Chronic kidney disease	Coronary heart disease	Stroke	Obesity
	(n=404) n %	(n=609) n%	(n=167) n (%)	(n=958) n%
Metformin	152 (37.6)	248 (40.7)	61 (36.5)	469 (49.0)
Sulfonylurea	73 (18.1)	108 (17.7)	29 (17.4)	159 (16.6)
Insulin	97 (24.0)	121 (19.9)	40 (23.9)	148 (15.4)
Other	55 (13.6)	82 (13.5)	22 (13.2)	110 (11.5)
Combination <sup>a</sup>	27 (6.7)	50 (8.2)	15 (9.0)	72 (7.5)

<sup>a</sup> Combination: Combination of two separate drugs for specific treatments related to diabetes mellitus



**Supplemental Figure 1.** Cumulative incidence function for cancer mortality among Medicaid-insured women diagnosed with breast cancer with preexisting type 2 diabetes mellitus and without diabetes.



**Supplemental Figure 2.** Cumulative incidence function for cardiovascular mortality among Medicaid-insured women diagnosed with breast cancer with preexisting type 2 diabetes mellitus and without diabetes.