



HHS Public Access

Author manuscript

J Allergy Clin Immunol. Author manuscript; available in PMC 2024 March 01.

Published in final edited form as:

J Allergy Clin Immunol. 2023 March ; 151(3): 656–670. doi:10.1016/j.jaci.2022.11.025.

Environmental Justice and Allergic Disease: A Work Group Report of the AAAAI Environmental Exposure and Respiratory Health Committee and the *Diversity, Equity and Inclusion* Committee

Allison J. Burbank, MD¹, Michelle L. Hernandez, MD FAAAAI¹, Akilah Jefferson, MD, MSc², Tamara T. Perry, MD, FAAAAI², Wanda Phipatanakul, MD, MS, FAAAAI³, Jill Poole, MD, FAAAA⁴, Elizabeth C. Matsui, MD, MHS, FAAAAI⁵

¹Division of Pediatric Allergy and Immunology, University of North Carolina School of Medicine, Children's Research Institute, Chapel Hill, NC, USA

²University of Arkansas for Medical Sciences, Little Rock, AR; Arkansas Children's Research Institute, Little Rock, AR, USA

³Division of Asthma, Allergy and Immunology, Boston Children's Hospital, Harvard Medical School, Boston, MA, USA

⁴Department of Internal Medicine, Division of Allergy and Immunology, University of Nebraska Medical Center, Omaha, NE, USA

⁵Departments of Population Health and Pediatrics, Dell Medical School at University of Texas at Austin, Austin, TX, USA.

Abstract

Environmental justice is the concept that all people have the right to live in a healthy environment, to be protected against environmental hazards, and to participate in decisions affecting their communities. Communities of color and low-income populations live, work and play in environments with disproportionate exposure to hazards associated with allergic disease. This unequal distribution of hazards has contributed to health disparities and is largely the result of systemic racism that promotes segregation of neighborhoods, disinvestment in predominantly racial/ethnic minority neighborhoods, and discriminatory housing, employment and lending practices. The AAAAI Environmental Exposure and Respiratory Health (EERH) Committee and Diversity, Equity and Inclusion (DEI) Committee jointly developed this report to improve allergy/

Corresponding Author: Allison J. Burbank, MD, 116 Manning Dr, CB 7231, Chapel Hill, NC 27599, Allison_burbank@med.unc.edu, Fax: 919-962-4421, Phone: 919-962-5136.

Disclosure statement: J.A. Poole has received research support from AstraZeneca (no monies) and is involved in clinical research studies with Takeda and GlaxoSmithKline, all unrelated to this project. T.T. Perry and W. Phipatanakul are members of the AAAAI Board of Directors. W. Phipatanakul is a member of the American Board of Allergy & Immunology Board of Directors and is an associate editor for *Journal of Allergy and Clinical Immunology*. The remainder of the authors declare that they have no relevant conflicts of interest.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

immunology specialists' awareness of environmental injustice, its roots in systemic racism, and its impact on health disparities in allergic disease. We present evidence supporting the relationship between exposure to environmental hazards, particularly at the neighborhood level, and the disproportionately high incidence and poor outcomes from allergic diseases in marginalized populations. Achieving environmental justice requires investment in at-risk communities to increase access to safe housing, clean air and water, employment opportunities, education, nutrition, and healthcare. Through policies that promote environmental justice, we can achieve greater health equity in allergic disease.

Keywords

Environmental justice; race; ethnicity; systemic racism; segregation; health disparities; asthma; allergic rhinitis; atopic dermatitis; pollution; allergen; psychosocial stress; obesity; nutrition

Introduction

Exposure to environmental hazards is linked with poor health outcomes. Communities of color and low income populations experience higher rates of exposure to environmental hazards, the result of centuries-old structural racism and societal hierarchies, which shaped the built and social environments in which they live (1–3). Racial and ethnic minority and impoverished communities also experience disproportionately high prevalence of allergic diseases and worse outcomes compared to more affluent and White communities (4). Already vulnerable groups experience this “double-hit” of increased exposure to hazards that causes further deterioration in health and well-being.

The AAAAI Committee on the Underserved (COU), now the AAAAI Diversity, Equity and Inclusion Committee (DEI), in their recent Work Group Report described the available medical literature on atopic disease disparities in racial and ethnic minority and other underserved populations and identified areas where further work is needed to reduce health disparities (5). Building upon this report, the Environmental Exposure and Respiratory Health (EERH) Committee and DEI jointly present a report on the impact of disproportionate exposure to environmental hazards on allergic disease prevalence and severity in marginalized communities, including communities of color and those living in poverty. Further, we will discuss the origins of this uneven distribution of environmental hazards and the factors that continue to perpetuate this inequality, leading to poorer health in some while other groups are less affected. Our review of communities differentially impacted by environmental hazards is by no means comprehensive, but we have chosen to focus on groups in which there is a significant amount of scientific literature addressing the relationship between environment and allergic disease. The goal of this report is to improve practicing allergy/immunology specialists' awareness of environmental injustice, its roots in systemic racism, and its impact on health disparities in allergic disease.

Throughout this report, multiple terms are used to describe racial and ethnic identities that are not necessarily interchangeable. For example, an individual may identify as Black but not as African American or as Latinx but not Hispanic. Methods for classifying racial and ethnic identity differ across publications and are dependent on whether information

was obtained through self-report, interviewer observation, or electronic health records or databases. When summarizing the literature, we have included the descriptors of racial and ethnic identity reported in the referenced publication. Sometimes the term “person (or people) of color” is used to refer to non-white persons or groups, recognizing the shared experiences of racism within multiple racial and ethnic identity groups.

Disparities in Allergic Disease Prevalence and Outcomes

Marked disparities in asthma prevalence and morbidity have been recognized for decades, particularly among African American and multiracial populations who experience 1.5–2 fold greater prevalence of asthma and 2–3 fold greater risk of emergency department (ED) visits, hospitalization and death compared to non-Hispanic White populations (5–7). Asthma prevalence is also higher among people living below 100% of the federal poverty line (6). Black and Hispanic respondents to the 2018 National Health Interview Survey self-reported allergic rhinitis (AR) less frequently than non-Hispanic White respondents (8), yet AR among racial and ethnic minority populations may be significantly under-recognized and under-diagnosed. Studies of predominantly Black and Hispanic urban children found that they experienced higher AR symptom burden and lower quality of life compared to White children (9, 10). The prevalence (11) and incidence (8) of eczema or atopic dermatitis (AD) was higher in Non-Hispanic Black populations compared to Hispanic and non-Hispanic White individuals, and children living in poverty also had higher prevalence of AD compared to children from higher-income families (8).

Most environmental exposure literature relevant to allergy/immunology is focused on asthma, AR and, to a lesser degree, AD. The remainder of this report will focus on the differential exposure of certain groups to environmental hazards and the potential impact on allergic disease prevalence and outcomes.

Conceptual framework for racial/ethnic health and environmental exposure disparities

As with most health conditions, allergic disease is influenced by a combination of genetic and environmental factors. Many studies of health disparities fail to adequately account for contextual factors and the portion of the disparity that is explained by these often-unmeasured factors. In Figure 1, we present a conceptual model for understanding the different factors that contribute to racial and ethnic health disparities and their interconnectedness.

Genetics

While historical use of the term ‘race’ has implied an underlying distinct biology, race is a social construct and not a biologic or genetic classification (12). A worldwide study of human populations found that 95% of human genetic diversity is accounted for by *within*-population differences, with only a minor component of variability attributable to *between*-population differences (13). This means that genetic variability between racial groups is much smaller than the variability seen between individuals from the same racial group (14).

To date, there has been little evidence that genetic differences between racial and ethnic groups explain health disparities between the groups. There have been multiple studies of the association between differences in genetic ancestry and allergic disease prevalence and morbidity within racial and ethnic groups (15–22). Attempts to estimate the proportion of a health disparity *between* racial/ethnic groups that is explained by genetic ancestry is complicated by the fact that genetic associations observed within one population are not “portable” to another population (12, 14) and any observed differences between groups are confounded by sociocultural and economic differences that exist between groups. Even in genetic studies done within groups, special care must be taken to avoid assuming that associations between ancestry and disease risk are causal since ancestry is confounded by environmental and other contextual factors that are difficult to fully control for. Adjustment for contextual factors is often limited to crude measures of socioeconomic status such as income and educational attainment that may not capture other important predictors of disease susceptibility and health outcomes. Gene-environment interaction as a contributor to asthma morbidity among residents of poor urban neighborhoods is a burgeoning area of research (23, 24).

Built and social environments and the role of neighborhoods in health

Environmental exposures are often narrowly defined to include exposures to contaminants in air, water, and soil as well as chemicals through foods, personal care products, and other consumer products and materials. However, the environment defined by broader terms includes both the physical and social environments and encompasses the complex interplay between exposures that contributes to health inequities. Elements of the physical, or built, environment include location of residence (inner city, rural, suburban), housing quality, proximity to major roadways and neighborhood zoning among many others. Elements of the social environment include poverty, opportunities for employment, availability of transportation, access to quality healthcare, access to green spaces for recreation, availability of nutritious food, and exposure to neighborhood violence and crime. Neighborhoods represent the intersection of the built and social environments. The role of neighborhood-level exposure is increasingly recognized as an important contributor to health (25), in some cases independently of individual factors (26). Neighborhood-level exposures are often driven by factors like poverty and residential segregation by race and ethnicity. Living in highly segregated neighborhoods has been associated with higher prevalence and severity of allergic diseases like asthma (25) and AD (27). Evidence suggests that neighborhood allergic disease burden is not fully explained by income and racial and ethnic composition, though these can serve as markers of other environmental characteristics and contextual factors that may explain a larger proportion of health disparities (28).

The spatial patterning of racial and ethnic minority neighborhoods, the built and social environments created by concentration of poverty in these areas, and disproportionate exposure to pollution sources in predominantly low-income neighborhoods with high percentages of people of color all stem from specific structural forces at work in the U.S. for many years.

How structural racism created racially and economically segregated neighborhoods

Residential segregation and concentration of poverty in America's inner cities originated during "The Great Migration" of the early to mid-20th century, when millions of African Americans moved from the rural South to large industrialized cities in the Northeast and Midwest for better employment opportunities (29). To deal with housing shortages following the Great Depression, the Federal Housing Administration (FHA) enacted policies that promoted moving working-class White families out of depression-era inner city public housing developments and to the suburbs, while restricting African Americans from purchasing suburban homes (30, 31). Through the practice of "red-lining", racial and ethnic minority neighborhoods were color-coded to indicate areas of increased risk to insure mortgages, therefore ensuring that residents could not purchase homes with federally-insured loans. The elevated risk inferred by red-lining also restricted community investment, resulting in fewer businesses, fewer job opportunities, lower property values, less money for public schools, and reduced access to healthcare services. Unequal access to education and employment created environments of concentrated poverty, isolation and high crime rates that are still present today (30–32).

These activities were not restricted to inner cities. Restrictive covenants and red-lining pushed many people of color into unincorporated communities located outside of municipal boundaries, often lacking sanitation and water services and without elected officials to represent them (33). To this day, suburban housing developments frequently restrict or prohibit the construction of low-income housing, resulting in continued segregation and concentration of low-income individuals and families in less desirable neighborhoods (34). In 2019, only 47% of public housing was located in low to moderate poverty areas (35). Real estate agents influence buyers by showing them homes in certain neighborhoods while directing them away from others based on racial or ethnic composition through a practice called 'steering', and though outlawed by the Fair Housing Act of 1968, evidence suggests that the practice continues (36, 37). Neighborhoods inhabited by predominantly minority and low-income families often have a higher concentration of industrial zones compared to affluent, predominantly White neighborhoods. These disparities originated from practices meant to restrict undesirable activities from affluent areas with political power and influence, while rezoning existing residential areas with high proportions of people of color and low-income families to allow for more industrial use. The result is further deterioration in neighborhood value and increased exposure of residents to environmental hazards. Meanwhile, ongoing racial prejudice in employment, housing, and lending practices continues to pose a barrier to home ownership and wealth building for people of color (38, 39). This is reflected in the fact that as of 2019, approximately 1 in 3 Black and 1 in 4 Latinx children in the U.S. lived in poverty (40).

Effects of environmental exposures on allergic disease and the role of unequal distribution of environmental hazards on allergy-related health disparities

In this section, we will discuss categories of environmental hazards linked to allergic disease and present evidence that certain groups are more exposed to hazards than others (Table 1). While increased exposure to environmental hazards may be *associated* with or play a causal role in allergic disease prevalence and severity, there are very few studies that have examined whether or to what extent disparate exposure to hazards explains disparities in allergic disease. More research is needed to understand the extent to which exposure to hazards or groups of hazards explain health disparities and how these exposures might affect different populations in distinct ways.

AIR POLLUTION

In much of the U.S., people of lower socioeconomic status and racial and ethnic minority populations tend to live in areas with the poorest air quality (41–43). In this section, we will discuss the major sources of ambient air pollution relevant to human health and the disproportionate exposure of some groups to these hazards.

Pollution and allergic disease

Fixed Source Emissions

Exposure to environmental hazards associated with living near industrial sites is associated with higher all-cause (44, 45) and disease-specific mortality (46). Coal-fired power generation, steel production and refining of petroleum products release a host of hazardous byproducts including sulfur dioxide (SO₂), nitrogen dioxide (NO₂), and particulate matter (PM) (47–49) that have been extensively studied for their contribution to respiratory disease and have been linked with higher asthma prevalence (50), asthma exacerbations (51–55), lower lung function (48, 56–58) and all-cause and respiratory mortality (45). In rural areas, concentrated animal feeding operations, or CAFOs, are sources of PM and gaseous emissions as well as noxious odors from open waste lagoons and the practice of spraying waste on fields (59, 60). Residing near large scale industrial livestock operations has been associated with adverse health effects including higher all-cause and infant mortality and disease-specific mortality (anemia, kidney disease, tuberculosis, and sepsis) (46). Residing near swine CAFOs has been associated with increased prevalence of asthma (61–63) and nasal allergies (63) and with increased odds of uncontrolled asthma (63) and reduced lung function (59). Rural communities are also increasingly the sites of hydraulic “fracking” for extraction of oil and natural gas, which involves use of chemicals, many of which are toxic to humans, to facilitate drilling and extraction. Traffic exhaust from trucks and equipment used in extraction have an additional environmental impact. Because of their concentration in rural areas, many fracking sites are out of range of air quality monitoring sites (64).

Groups with disproportionate exposure to fixed source air pollution

A study examining air quality across the U.S. found that monitored counties with the worst airway quality in terms of PM less than 2.5 μm ($\text{PM}_{2.5}$) had higher percentages of non-Hispanic Black residents and people living in poverty compared to counties with the best air quality (43). A 2017 joint report prepared by the Clean Air Task Force, National Association for the Advancement of Colored People (NAACP) and the National Medical Association reported that Black people in the U.S. are exposed to 38% more air pollution than Whites (65). Historical land use patterns in the U.S. have resulted in siting of industrial facilities such as solid waste disposal sites (66) and power-generating facilities (67) in predominantly Black and Hispanic neighborhoods (68, 69). An examination of demographic characteristics of people residing near hazardous waste facilities in the U.S. found that as distance from the facility decreased, the proportion of residents of color increased; socioeconomic disparities were also identified but were less extensive than racial disparities (70).

In 2019, rural areas of the U.S. had higher rates of poverty than urban areas among all racial/ethnic groups but particularly among Black (30.7%) and American Indian or Alaskan Native (29.6%) populations, compared to White populations (13.3%) (71, 72). CAFOs are typically sited in rural areas and are often located in high poverty neighborhoods and communities with a high proportion of people of color (61). Ten times as many CAFOs in North Carolina are located within census blocks with the highest poverty rates and proportion of non-White residents, compared to census blocks with the lowest poverty and non-White populations (60, 73). The environmental impact of fracking, including the chemicals used, their persistence in the environment and effects on nearby residents are not well understood (64, 74–76) but are likely to disproportionately affect residents of rural areas.

Mobile Source Emissions:

Mobile sources of air pollution include cars, trucks, buses, construction equipment, airplanes, trains and watercraft. Traffic-related air pollution, or TRAP, is a major source of air pollution comprised of products of fossil fuel combustion, including gases (nitrogen oxides, NO_x , carbon monoxide, CO, carbon dioxide, CO_2) and particulate emissions (PM, including diesel exhaust particles (DEP) and black carbon). TRAP exposure has been linked to a variety of negative health outcomes, including respiratory morbidity, cancer, and heart disease (77). TRAP exposure has been linked with increased risk of allergic sensitization (78–80), AR (81), skin barrier dysfunction (82) and AD (83), asthma (84–97), and exacerbation of established asthma (98–101).

Children may be particularly susceptible to the harmful effects of TRAP, especially during the prenatal period and during the first few years of life (102). The relationship between prenatal and early-life TRAP exposure and risk of wheeze and asthma is somewhat controversial, with many studies showing a significant association (84–96) and others reporting no association (103–109). Multiple prospective studies have shown positive associations between early life exposure to TRAP and risk of reduced lung function (110) and asthma (111–113), AR and AD (93). Some of this discrepancy may be related to differences in the way exposure was assessed as well as the window of exposure, timing of outcomes, effects of weather or composition of air pollutants. Traffic-related pollution is

highly localized, and air quality monitoring stations may not accurately capture the burden of pollutants experienced by those that live closest to major roadways (114).

Groups with disproportionate exposure to TRAP

Racial and ethnic minority and low-income populations incur greater exposure to neighborhood traffic than White and higher-income populations (69, 97, 114, 115). Among the U.S. national Environmental Influences on Child Health Outcomes (ECHO) Consortium, Commodore et al. found that a larger proportion of non-Hispanic Black and Hispanic children were exposed to neighborhood traffic compared to non-Hispanic White children (39.5%, 34.9%, and 12.4%, respectively) and were also more likely to report asthma symptoms (40.1%, 31.5%, and 19.3%, respectively) (97). Approximately 40% of inner city children of predominantly racial and ethnic minority status participating in the School Inner-City Asthma Study lived within 100 m of major roadways, and more than half attended a school that was also located within this buffer (116). These children had higher odds of experiencing asthma symptom days, poor asthma control and healthcare utilization compared to children living/attending school more than 100 m from major roadways.

In urban centers, bus depots and major truck routes are often located in or near low-income neighborhoods with high percentages of people of color (117). Residents of these communities often spend significant amounts of time on city sidewalks for transportation, recreation and socializing and are therefore more at risk of exposure to traffic exhaust. Additionally, lack of central air conditioning, which is twice as prevalent among urban Black people compared to White people (118), forces residents to spend more time outdoors where they are exposed to higher levels of traffic-related pollution. Several studies have reported that neighborhood disadvantage strengthens the association between air pollution exposure and respiratory disease (119–121). For example, in one study of households from the Panel Study of Income Dynamics, neighborhood poverty was a significant moderator of the association between early life pollutant exposure and childhood asthma risk, with significant associations in moderate and high poverty neighborhoods but not in low poverty neighborhoods (122).

HOUSING AND THE INDOOR ENVIRONMENT

Characteristics of poor-quality housing include cracks in walls and doors that allow pests to enter (cockroaches, rodents), leaky pipes with resultant mold and mildew growth, inadequate ventilation, lack of air conditioning and/or heating, and exposure to volatile organic compounds from building materials. A secondary analysis of 33,201 households surveyed in the 2011 American Housing Survey found that poor quality housing was independently associated with diagnosis of asthma and with asthma-related ED visits (123). Single site studies in different U.S. urban centers showed associations between dilapidated housing or housing code violations and elevated asthma hospitalization rates (124, 125); remediation of housing conditions to limit indoor asthma triggers resulted in significant improvements in asthma-related healthcare utilization (126). In this section, we will discuss the indoor environment and the groups most affected by environmental hazards that come with poor quality, delapidated housing.

Allergen

Mouse and cockroach allergen sensitization and exposure are clearly linked with pediatric asthma morbidity, with the strongest evidence for these associations seen in studies of inner-city children (127–130). Older homes, mobile homes and high-rise apartments including public housing structures were associated with frequent reports of pests and higher concentrations of pest allergens (131). Exposure to cockroach allergen in sensitized individuals was associated with asthma prevalence in a dose-dependent manner (132–135). Children who were sensitized and exposed to cockroach allergen were more likely to be hospitalized for asthma (136) and missed more school days (130, 137). Mouse sensitization and exposure in urban children were associated with greater asthma severity (138, 139), increased symptom frequency (131, 140–142), asthma-related acute care visits (141, 143, 144), hospitalizations (131, 141, 142, 145), low lung function (140) and elevated fractional exhaled nitric oxide (FeNO) (143, 145). Mouse sensitization was also associated with rhinitis symptoms in urban children with asthma (146). Mouse allergen remediation interventions successfully reduced allergen levels, and this corresponded to reductions in asthma morbidity (147) and increased lung function growth (148) in mouse-sensitized children.

Mold is also a common allergen found in increased concentrations in dilapidated structures (149), often due to condensation from inadequate heating or improperly repaired roofs or plumbing leaks that create conditions favorable to fungal growth. Indoor fungal exposure was associated with increased respiratory symptoms in adults and children, including cough and wheeze (150–154) and rhinitis symptoms (155, 156). Moisture damage in school buildings was associated with increased odds of nocturnal cough, wheeze, nasal symptoms and school absences in children (157, 158). *Alternaria alternata*, found in indoor and outdoor environments, was associated with increased prevalence of asthma (159) and with increased asthma symptoms in sensitized children (160). In addition to allergic inflammation, inflammatory fungal components like β -1,3-D-glucan, mycotoxins and volatile compounds were associated with non-allergic cough and nasopharyngeal and ocular irritation (161).

Groups with disproportionate exposure to allergens associated with poor housing conditions

People of color and low-income families are more likely than others to live in poor housing conditions (162), with over twice as many non-Hispanic Black individuals reporting living in substandard housing as White individuals according to the American Housing Survey (163). An even larger racial disparity in housing was found in non-metropolitan areas of the U.S., with African American households three times as likely to live in low quality housing (164). Low income and non-White students are more likely to attend schools with dilapidated infrastructure (165). Racial/ethnic minority identity, low educational attainment, renting a home, and living in neighborhoods with high poverty were associated with increased odds of the presence of pests in the home (131, 166, 167). Multiple single site studies have shown higher levels of cockroach and mouse allergen in the homes of racial and ethnic minority families compared to White, English-speaking families (128,

168, 169) and higher rates of sensitization to pest allergens in Black and Hispanic populations (168). Exposure to cockroach and mouse allergens was also highly prevalent in the school environment among children living in socioeconomically disadvantaged urban neighborhoods (160, 170–175). Visible mold, water damage, and damp conditions were common in low-income homes with children enrolled in a Seattle, WA Healthy Homes project. African American children had over 2 times the odds of sensitization to *Alternaria alternata*, compared to White children (176), and mold sensitization was identified as a significant predictor of asthma-related ED visits in Black children (177).

Indoor pollutants

Common indoor sources of pollution include combustion products like NO₂ and PM from fossil fuel-powered stoves and heaters, secondhand smoke (SHS), and infiltration of polluted outdoor air (178, 179). Poor ventilation may contribute to concentration of pollutants within indoor spaces where people spend up to 90% of the day (180, 181). In cross-sectional studies, the presence of a gas stove in the home was associated with increased odds of current asthma (182) and wheezing in the last 12 months (183). Indoor NO₂ exposure was associated with a dose-dependent increase in asthma symptoms, rescue medication use and airflow obstruction in inner-city children (178, 184–186) and those living in multifamily housing units (187), and increased airway hyperresponsiveness (188, 189). Volatile organic compounds (VOCs) like formaldehyde are emitted from consumer products like furniture, insulation materials, combustion appliances and from cigarette smoking (179, 190) and were associated with asthma prevalence in cross-sectional studies; prospective studies examining this relationship are lacking (191).

SHS exposure is common, with 14% of the U.S. adult population identifying as current smokers (192) and 30% of all U.S. children living in a household with a smoker (193). SHS exposure during the prenatal period was associated with low birth weight (194–197), preterm birth (198, 199), reduced lung function shortly after birth (200), and increased incidence of childhood asthma or wheezing (201–203). Postnatal parental smoking during a child's early life was associated with increased odds of wheezing illness and asthma diagnosis in children (201, 204–206), low lung function during childhood (202, 207, 208), increased emergency department visits (190, 209), and increased risk of allergic sensitization (210). Additionally, a systematic review found that both active and passive smoking were associated with AR and AD in children and adolescents (211), while in adults active smoking was associated with increased odds of adult-onset AD, chronic rhinitis, but *decreased* prevalence of AR (212–214).

Groups with disproportionate exposure to indoor pollutants

African Americans are more likely than other groups to live in older, less energy-efficient homes (215) and are more likely to use natural gas appliances (216). African Americans in the U.S. are also less likely to own their homes (43%) compared to non-Hispanic White populations (65%), and landlords have fewer incentives to retrofit properties with more energy-efficient electric appliances (217, 218). People living in poor quality housing with insufficient insulation and other defects that contribute to energy inefficiency will sometimes

use their gas stoves or ovens to heat their homes in winter (181), increasing exposure to byproducts of combustion like NO₂. Racial/ethnic identity was significantly associated with VOC exposures in a secondary analysis of personal exposure monitoring data from the National Health and Nutrition Examination Survey (NHANES) 1999–2000, with Black and Hispanic participants having significantly higher exposures than non-Hispanic White participants (219).

People self-identifying as American Indian/Alaska Native (20.9%) or Other (19.7%) were more likely to be current smokers compared to non-Hispanic White (15.5%), non-Hispanic Black (14.9%), Hispanic (8.8%) and non-Hispanic Asian individuals (7.2%)(192). Low educational attainment and lower annual household income were also associated with higher prevalence of current smoking and with maternal smoking during pregnancy (220–222). SHS exposure among Black children and those from households with high poverty and low parental education were 3 times higher than for Hispanic children (the reference group), low poverty and high education households (193, 223). Living in multiunit housing was associated with higher rates of SHS exposure (assessed by serum cotinine levels) in children compared to living in detached homes, even amongst those without known SHS exposure, suggesting potential passage of tobacco smoke through ventilation systems or defects in walls (224).

OTHER ENVIRONMENTAL EXPOSURES RELEVANT TO ALLERGIC DISEASE

Psychosocial stress

Neighborhood safety and caregiver psychosocial stress have been linked to health outcomes in children (225–227). Increased caregiver psychosocial stress was associated with higher prevalence of childhood asthma (228), higher frequency of asthma symptoms (229, 230), lower bronchodilator reversibility (231) and poorer adherence to asthma medications (232). A meta-analysis of observational studies found that children of women who experienced psychological stress during pregnancy had higher prevalence of wheezing and asthma than children born to mothers who did not report psychological stress (228). Stressful life events were associated with increased risk of new-onset asthma in children from the Tucson Children's Respiratory Study (233), and a study of a nationally representative sample of children found a positive relationship between number of adverse childhood experiences and odds of asthma (234). A history of physical or sexual abuse was associated with increased odds of current asthma, asthma-related healthcare visits, and asthma medication use among Puerto Rican children (235). Exposure to community violence and low caregiver-perceived safety of the neighborhood were associated with higher odds of asthma (236) and poor asthma control in children (237). Neighborhood violence has also been linked with depression in children and caregivers (238), which may contribute to increased asthma-related ED visits (239) and difficulty using asthma medications correctly (232). Higher perceived neighborhood safety, lower caregiver stress and lower depressive symptoms were associated with lower rates of asthma symptoms in children (227).

People of color experience higher exposure to psychosocial stress and community violence than other groups (240, 241). Black adolescents and adults were at increased risk of violence resulting in homicide or physical injury compared to White adolescents and adults (242, 243). In-depth interviews of predominantly Black inner-city adolescents revealed that 42% had witnessed someone being shot or knifed and 22% had witnessed someone being killed (244). Black children and adults were more likely to report experiencing adverse childhood experiences compared to White children and adults (242, 245, 246). Discrimination based on racial/ethnic identity is a significant source of psychological distress linked to poor health in adults and children (247), including asthma (248). The experience of racism can vary with skin color, leading some investigators to include percent African ancestry as a potential modifier of the relationship between discrimination and disease outcomes. African American children from the Study of African Americans, Asthma, Genes, and Environments (SAGE II) and Genes-environments and Admixture in Latino Americans (GALA II) trials reporting discrimination experienced nearly 80% higher odds of having asthma and nearly 2 times the odds of poor asthma control compared to those who reported no discrimination, regardless of amount of African ancestry or socioeconomic status (249).

Healthcare access

Despite greater need for healthcare services, racial/ethnic minority populations and the poor are less likely to receive care compared to White populations and are less likely to have health insurance, attend fewer doctor visits, and receive greater fragmentation of care due to limited options in care providers and less continuity of care between visits (250, 251). Throughout the 1960s-1990s, America's inner cities saw a decline in the number of outpatient primary care providers as well as private hospitals, whose emergency rooms served as primary care sources for a large proportion of the mostly minority and poor population (252–255). Remaining private hospitals sometimes limit the number of Medicaid and Medicare patients they will serve (253). On the other hand, public hospitals are often overcrowded, under-resourced and require long wait times to see a healthcare provider, leading some patients to forgo preventive care services. Black children and children from non-English-speaking Hispanic families with a diagnosis of asthma were less likely to have a “usual source of care” and to identify a specific provider compared to White children (256). Evidence suggests that some communities with high proportions of racial and ethnic minorities are “pharmacy deserts”, which reduces access to prescription medications (257). Black people were more likely to receive care for asthma in the ED (250) and were less likely to receive guidelines-based asthma care including prescriptions for controller medications (258–260). Rural minority individuals experienced similar barriers to accessing healthcare and in some cases had less access to care than urban minority and rural White individuals (261–263).

Diet and Food security

Poor diet quality is an important contributor to chronic disease and to obesity. Diet quality in the U.S. varies by demographic and socioeconomic factors, with White adults estimated to consume more vegetables, whole grains and milk than Black adults and low-income adults having generally lower diet quality than higher-income adults (264–266). Nutrition during early life may have immunomodulating effects that influence development of allergic

diseases like asthma and food allergy (267). Maternal diets high in fresh fruits, vegetables and fish (Mediterranean diet) were consistently associated with reduced risk of allergic disease in children, while maternal intake of vegetable oils and margarines were associated with higher risk. Maternal intake of fat soluble vitamins like A and E were associated with reduced risk of AR (268). Eating a diverse diet in the first year of life was associated with lower prevalence of childhood asthma, food allergy and food sensitization up to age 6 years (269).

Access to nutritious food is affected by the local food environment, including geographic distance to stores, access to transportation to get to a store, the variety of food choices available and the price of food. In 2019, food insecurity affected 34.9% of U.S. families living below the poverty line, 15.6% of Hispanic/Latinx and 19.1% of Black households, compared to 7.9% of White households (270). Increasing poverty and higher neighborhood racial segregation were associated with fewer supermarkets, with census tracts with a high percentage of Black residents having the fewest supermarkets and White census tracts having the most (271). Low income neighborhoods tend to have more convenience stores, corner markets and fast food options that tend to offer fewer healthy food choices like fresh produce. Food prices are another major determinant of diet since healthier fresh foods tend to be more expensive than processed foods. In addition to the loss of health benefits associated with a healthy diet, food insecurity also impacts health in other ways. Households are sometimes forced to make the choice between buying food and buying medication to treat chronic illnesses. An analysis of data from the National Health Interview Survey showed that food insecurity and chronic illness in adults was associated with underuse of medications because of cost constraints, and these individuals were more likely to identify as Hispanic or non-Hispanic Black (272).

Food insecurity is also associated with increased risk of obesity (273). In addition to an increased risk for food insecurity, a larger proportion of non-Hispanic Black and Hispanic children and adults in the U.S. are obese compared to non-Hispanic White individuals (274). Differences in the sources and types of food available as well as targeting African Americans for marketing of high-calorie and poorly nutritious foods are both potential contributors to the racial disparities in obesity rates (275, 276). There is a large literature linking obesity with increased risk of asthma (277–279), increased asthma severity (280–282) and AD (279, 283).

Green Space

The health benefits of public green spaces such as parks, sporting fields, greenways and community gardens are well-described and have become particularly important during the COVID-19 pandemic with the need for physical distancing when socializing (284–286). Urban green spaces provide benefits like access to space for physical activity, which may influence obesity development (287) and psychological well-being (288). Some have hypothesized that reduced exposure to environmental microbiota due to urbanization leads to immune dysfunction and impaired tolerance due to loss of biodiversity (289, 290). Green space was inversely associated with risk of allergic sensitization (291), AD, AR (292), and asthma (293), though it is unclear whether greenness has a direct effect on allergic disease

risk and morbidity or whether it mediates other factors that contribute to allergic disease (294). For example, green spaces may partially mitigate the negative effects of TRAP on risk of asthma in children (295). Living near green spaces in early life was associated with higher lung function up to 24 years of age (296).

Access to urban green spaces and their health benefits are not equally distributed among populations, and low income and racial/ethnic minority groups tend to have the least access to these spaces (297–299). The average size of parks in predominantly White neighborhoods in New York City was over 3 times the size of parks in predominantly Black neighborhoods (297). Race, ethnicity and income are important modifiers of the health protective effects of greenspace. In a study of pregnant women in South Carolina, women from low-income Black communities with the lowest amount of greenspace experienced excess risk for poor pregnancy outcomes compared to White women from economically privileged White communities with similarly low greenspace (300). In addition to the amount of green space available, other factors lead to inequities in use of green spaces including lack of transportation, quality of parks, and safety concerns.

ENVIRONMENTAL JUSTICE

Environmental injustice is the disproportionate exposure of vulnerable groups to environmental hazards and their resultant negative health effects as well as unequal protection against these hazards by laws, policies and regulatory agencies (67). Beyond the more obvious hazards like pollution and toxic waste, environmental hazards also include poverty, psychosocial stress, exposure to violence and lack of access to resources and services like quality healthcare, healthy food choices and green space. The cumulative effects of these hazards may have profound implications for health, life opportunity, life expectancy and quality of life (301, 302). The goal of the environmental justice (EJ) movement is to provide equal protection from environmental health hazards to all people regardless of race, ethnicity, or income level and to ensure that all people have the opportunity to be involved in decision-making processes impacting their communities (303).

Community engagement is a key component of successful environmental intervention programs for asthma (126), but accomplishing this goal will require investment in EJ beyond the community level, with state and federal government resources needed to identify at-risk communities and implement solutions. State and community-based urban asthma programs have demonstrated improved asthma outcomes through home environmental interventions and education (304–306) but require stable funding to continue their work and expand to other at-risk communities. With respect to hazards like air pollution, land use planning and proposed zoning changes should be carefully and thoughtfully evaluated with involvement of public health professionals and input from community members (307). Unincorporated communities, which typically do not have local elected officials, should be recognized as particularly vulnerable due to their limited ability to participate in decision making about their communities (33). Governments should consider policies that limit idling of school and city buses and replace public transportation vehicles with newer, lower emission vehicles. State and federal environmental regulations on industry should be strictly enforced. Strategies for mitigation of indoor environmental hazards include education campaigns

focused on reducing asthma triggers in homes and schools, housing mobility programs, and improvements in quality inspection of public housing (308). Integrated pest management interventions and education programs in homes (147) and schools (309) have shown benefit in children with asthma in well-designed studies. Poverty is a key driver of exposure to environmental hazards like violence and psychosocial stress, food insecurity, and lack of access to healthcare services. There is a large need for community development through investment in affordable housing, schools, and health clinics as well as small businesses that create jobs and help residents to build wealth (226). Government programs similar to the Obama administration's "Let's Move!" program can help communities improve access to healthy food and prevent obesity by incentivizing the placement of supermarkets, farmers markets, food banks and pantries in underserved areas, establishing nutrition standards for school lunches and improving access to outdoor spaces for physical activity (310). Increased rates of psychosocial stress should be met with better screening for patient and caregiver stress and improved access to mental health care, including school-based resources for children, with social workers and school nurses assisting with identification of mental healthcare needs. Better collaborations between healthcare providers and community-based organizations will assist children and families with significant needs.

In this report, we have emphasized that the neighborhood is a key determinant of the overall health of its inhabitants, including asthma and other allergic diseases. Racial *and* economic integration (avoiding gentrification) of neighborhoods can create more equitable access to quality affordable housing, better education and employment opportunities, nutrition and green spaces and should be prioritized by public policy-makers and housing industry leaders.

REMAINING KNOWLEDGE GAPS AND DIRECTIONS FOR FUTURE RESEARCH

There is abundant evidence that exposure to environmental hazards is associated with poorer health, but more studies are needed to understand the extent to which exposure to hazards (or groups of hazards) explain health disparities. Future research should account for contextual factors as determinants of the distribution of allergic disease burden in a population and should be cognizant that the impact of environmental characteristics on allergic disease may vary by race and ethnicity (and plan their study's analysis strategy to account for this) (28). There is a need to identify community-level interventions that are most impactful for improving the health of community members. Interventions capable of effectively reducing exposure to hazards should be studied to determine whether these interventions also reduce allergic disease burden among populations and whether they reduce health disparities among groups within a population. Finally, most studies of environmental exposures have focused more on short-term health outcomes, like acute asthma exacerbations, for example. But there is a major gap in knowledge regarding the impact of long-term exposure on health later in life, such as the effect of early life exposures on adult lung health disparities in racial and ethnic minority populations. In conclusion, the scientific literature is clear that when it comes to health, where you live matters. Everyone has the right to live in a community that promotes health and opportunity.

Funding Sources:

A.J. Burbank received funding support from NIH NCATS UL1TR002489 and KL2TR002490. E.C. Matsui received funding support from NIH NHLBI R34HL159126, NIEHS R01ES026170, and NIAID K24AI114769. M.L. Hernandez received funding support from NIH NHLBI R01HL135235 and NCATS UL1TR002489 and KL2TR002490. J.A. Poole received funding support from Department of Defense #PR200793 and NIH NIOSH U54OH010162 and R01OH012045.

Abbreviations:

COU	Committee on the Underserved
DEI	now the Diversity, Equity and Inclusion Committee
EERH	Environmental Exposures and Respiratory Health
AR	allergic rhinitis
FHA	Federal Housing Administration
SO₂	sulfur dioxide
NO₂	nitrogen dioxide
PM	particulate matter
CAFO	concentrated animal feeding operations
PM_{2.5}	particulate matter less than 2.5 μm
NAACP	National Association for the Advancement of Colored People
TRAP	traffic related air pollution
NO_x	nitrogen oxides
CO	carbon monoxide
CO₂	carbon dioxide
DEP	diesel exhaust particles
ECHO	Environmental Influences on Child Health Outcomes
ED	emergency department
FeNO	fractional exhaled nitric oxide
VOC	volatile organic compounds
SHS	secondhand smoke
NHANES	National Health and Nutrition Examination Survey
SAGE II	Study of African Americans, Asthma, Genes, and Environments
GALA II	Genes-environments and Admixture in Latino American

References

1. Grove MO, L; Pickett S; Boone C; Buckley G; Locke DH; Lord C; Hall B The Legacy Effect: Understanding How Segregation and Environmental Injustice Unfold over Time in Baltimore. *Annals of the American Association of Geographers*. 2018;108(2):524–37.
2. Clark LP, Millet DB, Marshall JD. National patterns in environmental injustice and inequality: outdoor NO₂ air pollution in the United States. *PLoS One*. 2014;9(4):e94431. [PubMed: 24736569]
3. Dey T, Dominici F. COVID-19, Air Pollution, and Racial Inequity: Connecting the Dots. *Chem Res Toxicol*. 2021;34(3):669–71. [PubMed: 33377385]
4. Polcari I, Becker L, Stein SL, Smith MS, Paller AS. Filaggrin gene mutations in African Americans with both ichthyosis vulgaris and atopic dermatitis. *Pediatr Dermatol*. 2014;31(4):489–92. [PubMed: 24920311]
5. Davis CM, Apter AJ, Casillas A, Foggs MB, Louisias M, Morris EC, et al. Health disparities in allergic and immunologic conditions in racial and ethnic underserved populations: A Work Group Report of the AAAAI Committee on the Underserved. *J Allergy Clin Immunol*. 2021;147(5):1579–93. [PubMed: 33713767]
6. Pate CA, Zahran HS, Qin X, Johnson C, Hummelman E, Malilay J. Asthma Surveillance - United States, 2006–2018. *MMWR Surveill Summ*. 2021;70(5):1–32.
7. Akinbami LJ, Moorman JE, Bailey C, Zahran HS, King M, Johnson CA, et al. Trends in asthma prevalence, health care use, and mortality in the United States, 2001–2010. *NCHS Data Brief*. 2012(94):1–8.
8. Allergies and Hay Fever: Summary Health Statistics Tables. National Center for Health Statistics; 2018.
9. Everhart RS, Kopel SJ, Esteban CA, McQuaid EL, Klein R, McCue CE, et al. Allergic rhinitis quality of life in urban children with asthma. *Ann Allergy Asthma Immunol*. 2014;112(4):365–70 e1. [PubMed: 24589166]
10. Esteban CA, Klein RB, Kopel SJ, McQuaid EL, Fritz GK, Seifer R, et al. Underdiagnosed and Undertreated Allergic Rhinitis in Urban School-Aged Children with Asthma. *Pediatr Allergy Immunol Pulmonol*. 2014;27(2):75–81. [PubMed: 24963455]
11. Shaw TE, Currie GP, Koudelka CW, Simpson EL. Eczema prevalence in the United States: data from the 2003 National Survey of Children’s Health. *J Invest Dermatol*. 2011;131(1):67–73. [PubMed: 20739951]
12. National Research Council. *Critical Perspectives on Racial and Ethnic Differences in Health in Late Life*. Anderson N, Bulatao RA, Cohen B, editor. Washington D.C.: National Academies Press (US); 2004.
13. Rosenberg NA, Pritchard JK, Weber JL, Cann HM, Kidd KK, Zhivotovsky LA, et al. Genetic structure of human populations. *Science*. 2002;298(5602):2381–5. [PubMed: 12493913]
14. Chaturvedi N. Ethnicity as an epidemiological determinant—crudely racist or crucially important? *Int J Epidemiol*. 2001;30(5):925–7. [PubMed: 11689494]
15. Rumpel JA, Ahmedani BK, Peterson EL, Wells KE, Yang M, Levin AM, et al. Genetic ancestry and its association with asthma exacerbations among African American subjects with asthma. *J Allergy Clin Immunol*. 2012;130(6):1302–6. [PubMed: 23069492]
16. Levin AM, Wang Y, Wells KE, Padhukasahasram B, Yang JJ, Burchard EG, et al. Nocturnal asthma and the importance of race/ethnicity and genetic ancestry. *Am J Respir Crit Care Med*. 2014;190(3):266–73. [PubMed: 24937318]
17. Flores C, Ma SF, Pino-Yanes M, Wade MS, Perez-Mendez L, Kittles RA, et al. African ancestry is associated with asthma risk in African Americans. *PLoS One*. 2012;7(1):e26807. [PubMed: 22235241]
18. Vergara C, Caraballo L, Mercado D, Jimenez S, Rojas W, Rafaels N, et al. African ancestry is associated with risk of asthma and high total serum IgE in a population from the Caribbean Coast of Colombia. *Hum Genet*. 2009;125(5–6):565–79. [PubMed: 19290544]

19. Wells KE, Cajjal S, Peterson EL, Ahmedani BK, Kumar R, Lanfear DE, et al. Assessing differences in inhaled corticosteroid response by self-reported race-ethnicity and genetic ancestry among asthmatic subjects. *J Allergy Clin Immunol*. 2016;137(5):1364–9 e2. [PubMed: 27016472]
20. Gould W, Peterson EL, Karungi G, Zoratti A, Gaggin J, Toma G, et al. Factors predicting inhaled corticosteroid responsiveness in African American patients with asthma. *J Allergy Clin Immunol*. 2010;126(6):1131–8. [PubMed: 20864153]
21. Pino-Yanes M, Thakur N, Gignoux CR, Galanter JM, Roth LA, Eng C, et al. Genetic ancestry influences asthma susceptibility and lung function among Latinos. *J Allergy Clin Immunol*. 2015;135(1):228–35. [PubMed: 25301036]
22. Gamble C, Talbott E, Youk A, Holguin F, Pitt B, Silveira L, et al. Racial differences in biologic predictors of severe asthma: Data from the Severe Asthma Research Program. *J Allergy Clin Immunol*. 2010;126(6):1149–56 e1. [PubMed: 21051082]
23. Massoud AH, Charbonnier LM, Lopez D, Pellegrini M, Phipatanakul W, Chatila TA. An asthma-associated IL4R variant exacerbates airway inflammation by promoting conversion of regulatory T cells to TH17-like cells. *Nat Med*. 2016;22(9):1013–22. [PubMed: 27479084]
24. Lai PS, Massoud AH, Xia M, Petty CR, Cunningham A, Chatila TA, et al. Gene-environment interaction between an IL4R variant and school endotoxin exposure contributes to asthma symptoms in inner-city children. *J Allergy Clin Immunol*. 2018;141(2):794–6 e3. [PubMed: 28943468]
25. Sullivan K, Thakur N. Structural and Social Determinants of Health in Asthma in Developed Economies: a Scoping Review of Literature Published Between 2014 and 2019. *Curr Allergy Asthma Rep*. 2020;20(2):5. [PubMed: 32030507]
26. Diez Roux AV. Investigating neighborhood and area effects on health. *Am J Public Health*. 2001;91(11):1783–9. [PubMed: 11684601]
27. Tackett KJ, Jenkins F, Morrell DS, McShane DB, Burkhart CN. Structural racism and its influence on the severity of atopic dermatitis in African American children. *Pediatr Dermatol*. 2020;37(1):142–6. [PubMed: 31854003]
28. Zarate RA, Zigler C, Cubbin C, Matsui EC. Neighborhood-level variability in asthma-related emergency department visits in Central Texas. *J Allergy Clin Immunol*. 2021;148(5):1262–9 e6. [PubMed: 34506851]
29. The Great Migration: A&E Television Networks; [updated June 28, 2021. Available from: <https://www.history.com/topics/black-history/great-migration>.
30. Gross T. A “Forgotten History” of How the U.S. Government Segregated America: National Public Radio; 2017 [Available from: www.npr.org/2017/05/03/526655831/a-forgotten-history-of-how-the-u-s-government-segregated-america.
31. Rothstein R. *The Color of Law: A forgotten history of how our government segregated America*: Liveright; 2018.
32. Peterson R, Krivo LJ. Racial segregation and black urban homicide. *Social Forces*. 1993;71:1001–26.
33. Gomez-Vidal C, Gomez AM. Invisible and unequal: Unincorporated community status as a structural determinant of health. *Soc Sci Med*. 2021;285:114292. [PubMed: 34411968]
34. Loh T, Coes C, Buthe B. *The Great Real Estate Reset*: Brookings Institute; 2020 [Available from: <https://www.brookings.edu/essay/trend-1-separate-and-unequal-neighborhoods-are-sustaining-racial-and-economic-injustice-in-the-us/>.
35. Center on Budget and Policy Priorities. *Policy Basics: Public Housing 2021* [updated June 16, 2021. Available from: <https://www.cbpp.org/research/public-housing>.
36. Galster GGE. By Words and Deeds: Racial Steering by Real Estate Agents in the U.S. in 2000. *Journal of the American Planning Association*. 2005;71(3):251–68.
37. Verstraete JVP. Ethnic discrimination upon request? Real estate agents’ strategies for discriminatory questions of clients. *Journal of Housing and the Built Environment*. 2019;35:703–21.
38. Glantz A, Martinez E. Modern-day redlining: How banks block people of color from homeownership. *Chicago Tribune*. 2018 February 17, 2018.

39. Pager D, Shepherd H. The Sociology of Discrimination: Racial Discrimination in Employment, Housing, Credit, and Consumer Markets. *Annu Rev Sociol.* 2008;34:181–209. [PubMed: 20689680]
40. Kids Count Data Center: Children in poverty by race and ethnicity in the United States: Annie E. Casey Foundation; [Available from: <https://datacenter.kidscount.org/data/tables/44-children-in-poverty-by-race-and-ethnicity#detailed/1/any/false/1729,37,871,870,573,869,36,868,867,133/10,11,9,12,1,185,13/324,323>].
41. Gray SC, Edwards SE, Miranda ML. Race, socioeconomic status, and air pollution exposure in North Carolina. *Environ Res.* 2013;126:152–8. [PubMed: 23850144]
42. Brochu PJ, Yanosky JD, Paciorek CJ, Schwartz J, Chen JT, Herrick RF, et al. Particulate air pollution and socioeconomic position in rural and urban areas of the Northeastern United States. *Am J Public Health.* 2011;101 Suppl 1:S224–30. [PubMed: 21836114]
43. Miranda ML, Edwards SE, Keating MH, Paul CJ. Making the environmental justice grade: the relative burden of air pollution exposure in the United States. *Int J Environ Res Public Health.* 2011;8(6):1755–71. [PubMed: 21776200]
44. Bazyar J, Pourvakhshoori N, Khankeh H, Farrokhi M, Delshad V, Rajabi E. A comprehensive evaluation of the association between ambient air pollution and adverse health outcomes of major organ systems: a systematic review with a worldwide approach. *Environ Sci Pollut Res Int.* 2019;26(13):12648–61. [PubMed: 30903465]
45. Orellano P, Reynoso J, Quaranta N. Short-term exposure to sulphur dioxide (SO₂) and all-cause and respiratory mortality: A systematic review and meta-analysis. *Environ Int.* 2021;150:106434. [PubMed: 33601225]
46. Kravchenko J, Rhew SH, Akushevich I, Agarwal P, Lyerly HK. Mortality and Health Outcomes in North Carolina Communities Located in Close Proximity to Hog Concentrated Animal Feeding Operations. *N C Med J.* 2018;79(5):278–88. [PubMed: 30228132]
47. Amster E, Lew Levy C. Impact of Coal-fired Power Plant Emissions on Children’s Health: A Systematic Review of the Epidemiological Literature. *Int J Environ Res Public Health.* 2019;16(11).
48. Barbone F, Catelan D, Pistelli R, Accetta G, Grechi D, Rusconi F, et al. A Panel Study on Lung Function and Bronchial Inflammation among Children Exposed to Ambient SO₂ from an Oil Refinery. *Int J Environ Res Public Health.* 2019;16(6).
49. Prioleau T. Environmental Impact of the Petroleum Industry. Atlanta, GA: Hazardous Substance Research Centers/South & Southwest Outreach Program; 2003.
50. Bai S, Zhao X, Liu Y, Lin S, Liu Y, Wang Z, et al. The effect window for sulfur dioxide exposure in pregnancy on childhood asthma and wheezing: A case-control study. *Environ Res.* 2022;204(Pt C):112286. [PubMed: 34743895]
51. Oftedal B, Nafstad P, Magnus P, Bjorkly S, Skrondal A. Traffic related air pollution and acute hospital admission for respiratory diseases in Drammen, Norway 1995–2000. *Eur J Epidemiol.* 2003;18(7):671–5. [PubMed: 12952141]
52. Galan I, Tobias A, Banegas JR, Aranguiz E. Short-term effects of air pollution on daily asthma emergency room admissions. *Eur Respir J.* 2003;22(5):802–8. [PubMed: 14621088]
53. Smargiassi A, Kosatsky T, Hicks J, Plante C, Armstrong B, Villeneuve PJ, et al. Risk of asthmatic episodes in children exposed to sulfur dioxide stack emissions from a refinery point source in Montreal, Canada. *Environ Health Perspect.* 2009;117(4):653–9. [PubMed: 19440507]
54. Zheng XY, Orellano P, Lin HL, Jiang M, Guan WJ. Short-term exposure to ozone, nitrogen dioxide, and sulphur dioxide and emergency department visits and hospital admissions due to asthma: A systematic review and meta-analysis. *Environ Int.* 2021;150:106435. [PubMed: 33601224]
55. Zheng XY, Ding H, Jiang LN, Chen SW, Zheng JP, Qiu M, et al. Association between Air Pollutants and Asthma Emergency Room Visits and Hospital Admissions in Time Series Studies: A Systematic Review and Meta-Analysis. *PLoS One.* 2015;10(9):e0138146. [PubMed: 26382947]
56. Yu S, Park S, Park CS, Kim S. Association between the Ratio of FEV₁ to FVC and the Exposure Level to Air Pollution in Never-smoking Adult Refractory Asthmatics Using Data Clustered

- by Patient in the Soonchunhyang Asthma Cohort Database. *Int J Environ Res Public Health*. 2018;15(11).
57. Gehring U, Gruzjeva O, Agius RM, Beelen R, Custovic A, Cyrus J, et al. Air pollution exposure and lung function in children: the ESCAPE project. *Environ Health Perspect*. 2013;121(11–12):1357–64. [PubMed: 24076757]
 58. Gauderman WJ, Avol E, Gilliland F, Vora H, Thomas D, Berhane K, et al. The effect of air pollution on lung development from 10 to 18 years of age. *N Engl J Med*. 2004;351(11):1057–67. [PubMed: 15356303]
 59. Borlee F, Yzermans CJ, Aalders B, Rooijackers J, Krop E, Maassen CBM, et al. Air Pollution from Livestock Farms Is Associated with Airway Obstruction in Neighboring Residents. *Am J Respir Crit Care Med*. 2017;196(9):1152–61. [PubMed: 28489427]
 60. Wing S, Wolf S. Intensive livestock operations, health, and quality of life among eastern North Carolina residents. *Environ Health Perspect*. 2000;108(3):233–8. [PubMed: 10706529]
 61. Mirabelli MC, Wing S, Marshall SW, Wilcosky TC. Race, poverty, and potential exposure of middle-school students to air emissions from confined swine feeding operations. *Environ Health Perspect*. 2006;114(4):591–6. [PubMed: 16581551]
 62. Merchant JA, Naleway AL, Svendsen ER, Kelly KM, Burmeister LF, Stromquist AM, et al. Asthma and farm exposures in a cohort of rural Iowa children. *Environ Health Perspect*. 2005;113(3):350–6. [PubMed: 15743727]
 63. Schultz AA, Peppard P, Gangnon RE, Malecki KMC. Residential proximity to concentrated animal feeding operations and allergic and respiratory disease. *Environ Int*. 2019;130:104911. [PubMed: 31238264]
 64. Johnston J, Cushing L. Chemical Exposures, Health, and Environmental Justice in Communities Living on the Fenceline of Industry. *Curr Environ Health Rep*. 2020;7(1):48–57. [PubMed: 31970715]
 65. Fleischman L, Franklin M. Fumes Across the Fence-Line: The Health Impacts of Air Pollution from Oil & Gas Facilities on African American Communities. 2017.
 66. Bullard RD. Solid waste sites and the black Houston community. *Sociol Inq*. 1983;53(2–3):273–88. [PubMed: 11635985]
 67. Maantay J. Asthma and air pollution in the Bronx: methodological and data considerations in using GIS for environmental justice and health research. *Health Place*. 2007;13(1):32–56. [PubMed: 16311064]
 68. Perlin SA, Sexton K, Wong DW. An examination of race and poverty for populations living near industrial sources of air pollution. *J Expo Anal Environ Epidemiol*. 1999;9(1):29–48. [PubMed: 10189625]
 69. Martenies SE, Milano CW, Williams GO, Batterman SA. Disease and Health Inequalities Attributable to Air Pollutant Exposure in Detroit, Michigan. *Int J Environ Res Public Health*. 2017;14(10).
 70. Bullard RM, P; Saha R; Wright B Toxic Wastes and Race at Twenty: 1987–2007. United Church of Christ Justice & Witness Ministries; 2007.
 71. U.S. Department of Agriculture Economic Research Service. Data show U.S. Poverty rates in 2019 higher in rural areas than in urban for racial/ethnic groups; [updated August 23, 2021. Available from: <https://www.ers.usda.gov/data-products/chart-gallery/gallery/chart-detail/?chartId=101903>.
 72. Probst JS, ME; Jespersen KP; Willert K; Swann RS; McDuffie JA. Minorities in Rural America: An Overview of Population Characteristics. Columbia, SC: University of South Carolina, Norman J. Arnold School of Public Health, Department of Health Administration; 2002.
 73. Wing SJJ. Industrial Hog Operations in North Carolina Disproportionately Impact African-Americans, Hispanics and American Indians: NC Policy Watch; 2014 [Available from: ncpolicywatch.com.
 74. Stringfellow WT, Camarillo MK, Domen JK, Sandelin WL, Varadharajan C, Jordan PD, et al. Identifying chemicals of concern in hydraulic fracturing fluids used for oil production. *Environ Pollut*. 2017;220(Pt A):413–20. [PubMed: 27743793]

75. Maloney KO, Baruch-Mordo S, Patterson LA, Nicot JP, Entrekin SA, Fargione JE, et al. Unconventional oil and gas spills: Materials, volumes, and risks to surface waters in four states of the U.S. *Sci Total Environ*. 2017;581–582:369–77.
76. Czolowski ED, Santoro RL, Srebotnjak T, Shonkoff SBC. Toward Consistent Methodology to Quantify Populations in Proximity to Oil and Gas Development: A National Spatial Analysis and Review. *Environ Health Perspect*. 2017;125(8):086004. [PubMed: 28858829]
77. Health Effects Institute. Traffic-related air pollution: a critical review of the literature on emissions, exposure, and health effects. Boston, MA; 2010.
78. Manners S, Alam R, Schwartz DA, Gorska MM. A mouse model links asthma susceptibility to prenatal exposure to diesel exhaust. *J Allergy Clin Immunol*. 2014;134(1):63–72. [PubMed: 24365139]
79. Gruzieva O, Bellander T, Eneroth K, Kull I, Melen E, Nordling E, et al. Traffic-related air pollution and development of allergic sensitization in children during the first 8 years of life. *J Allergy Clin Immunol*. 2012;129(1):240–6. [PubMed: 22104609]
80. Codispoti CD, LeMasters GK, Levin L, Reponen T, Ryan PH, Biagini Myers JM, et al. Traffic pollution is associated with early childhood aeroallergen sensitization. *Ann Allergy Asthma Immunol*. 2015;114(2):126–33. [PubMed: 25499550]
81. Jung DY, Leem JH, Kim HC, Kim JH, Hwang SS, Lee JY, et al. Effect of Traffic-Related Air Pollution on Allergic Disease: Results of the Children’s Health and Environmental Research. *Allergy Asthma Immunol Res*. 2015;7(4):359–66. [PubMed: 25936911]
82. Dijkhoff IM, Drasler B, Karakocak BB, Petri-Fink A, Valacchi G, Eeman M, et al. Impact of airborne particulate matter on skin: a systematic review from epidemiology to in vitro studies. *Part Fibre Toxicol*. 2020;17(1):35. [PubMed: 32711561]
83. Lee YL, Su HJ, Sheu HM, Yu HS, Guo YL. Traffic-related air pollution, climate, and prevalence of eczema in Taiwanese school children. *J Invest Dermatol*. 2008;128(10):2412–20. [PubMed: 18449213]
84. Brunst KJ, Ryan PH, Brokamp C, Bernstein D, Reponen T, Lockey J, et al. Timing and Duration of Traffic-related Air Pollution Exposure and the Risk for Childhood Wheeze and Asthma. *Am J Respir Crit Care Med*. 2015;192(4):421–7. [PubMed: 26106807]
85. Carlsten C, Dybuncio A, Becker A, Chan-Yeung M, Brauer M. Traffic-related air pollution and incident asthma in a high-risk birth cohort. *Occup Environ Med*. 2011;68(4):291–5. [PubMed: 20952556]
86. Clark NA, Demers PA, Karr CJ, Koehoorn M, Lencar C, Tamburic L, et al. Effect of early life exposure to air pollution on development of childhood asthma. *Environ Health Perspect*. 2010;118(2):284–90. [PubMed: 20123607]
87. Gehring U, Wijga AH, Brauer M, Fischer P, de Jongste JC, Kerkhof M, et al. Traffic-related air pollution and the development of asthma and allergies during the first 8 years of life. *Am J Respir Crit Care Med*. 2010;181(6):596–603. [PubMed: 19965811]
88. Gehring U, Wijga AH, Hoek G, Bellander T, Berdel D, Bruske I, et al. Exposure to air pollution and development of asthma and rhinoconjunctivitis throughout childhood and adolescence: a population-based birth cohort study. *Lancet Respir Med*. 2015;3(12):933–42. [PubMed: 27057569]
89. Gruzieva O, Bergstrom A, Hulchiy O, Kull I, Lind T, Melen E, et al. Exposure to air pollution from traffic and childhood asthma until 12 years of age. *Epidemiology*. 2013;24(1):54–61. [PubMed: 23222555]
90. Jerrett M, Shankardass K, Berhane K, Gauderman WJ, Kunzli N, Avol E, et al. Traffic-related air pollution and asthma onset in children: a prospective cohort study with individual exposure measurement. *Environ Health Perspect*. 2008;116(10):1433–8. [PubMed: 18941591]
91. McConnell R, Islam T, Shankardass K, Jerrett M, Lurmann F, Gilliland F, et al. Childhood incident asthma and traffic-related air pollution at home and school. *Environ Health Perspect*. 2010;118(7):1021–6. [PubMed: 20371422]
92. Morgenstern V, Zutavern A, Cyrys J, Brockow I, Gehring U, Koletzko S, et al. Respiratory health and individual estimated exposure to traffic-related air pollutants in a cohort of young children. *Occup Environ Med*. 2007;64(1):8–16. [PubMed: 16912084]

93. Morgenstern V, Zutavern A, Cyrus J, Brockow I, Koletzko S, Kramer U, et al. Atopic diseases, allergic sensitization, and exposure to traffic-related air pollution in children. *Am J Respir Crit Care Med.* 2008;177(12):1331–7. [PubMed: 18337595]
94. Wu TJ, Wu CF, Chen BY, Lee YL, Guo YL. Age of asthma onset and vulnerability to ambient air pollution: an observational population-based study of adults from Southern Taiwan. *BMC Pulm Med.* 2016;16(1):54. [PubMed: 27094099]
95. Lindgren A, Strohm E, Montnemery P, Nihlen U, Jakobsson K, Axmon A. Traffic-related air pollution associated with prevalence of asthma and COPD/chronic bronchitis. A cross-sectional study in Southern Sweden. *Int J Health Geogr.* 2009;8:2. [PubMed: 19154599]
96. Lindgren A, Strohm E, Nihlen U, Montnemery P, Axmon A, Jakobsson K. Traffic exposure associated with allergic asthma and allergic rhinitis in adults. A cross-sectional study in southern Sweden. *Int J Health Geogr.* 2009;8:25. [PubMed: 19419561]
97. Commodore S, Ferguson PL, Neelon B, Newman R, Grobman W, Tita A, et al. Reported Neighborhood Traffic and the Odds of Asthma/Asthma-Like Symptoms: A Cross-Sectional Analysis of a Multi-Racial Cohort of Children. *Int J Environ Res Public Health.* 2020;18(1).
98. Delfino RJ, Wu J, Tjoa T, Gullesterian SK, Nickerson B, Gillen DL. Asthma morbidity and ambient air pollution: effect modification by residential traffic-related air pollution. *Epidemiology.* 2014;25(1):48–57. [PubMed: 24240657]
99. Chang J, Delfino RJ, Gillen D, Tjoa T, Nickerson B, Cooper D. Repeated respiratory hospital encounters among children with asthma and residential proximity to traffic. *Occup Environ Med.* 2009;66(2):90–8. [PubMed: 19151227]
100. Erbas B, Kelly AM, Physick B, Code C, Edwards M. Air pollution and childhood asthma emergency hospital admissions: estimating intra-city regional variations. *Int J Environ Health Res.* 2005;15(1):11–20. [PubMed: 16025811]
101. Esposito S, Galeone C, Lelii M, Longhi B, Ascolese B, Senatore L, et al. Impact of air pollution on respiratory diseases in children with recurrent wheezing or asthma. *BMC Pulm Med.* 2014;14:130. [PubMed: 25098250]
102. Jung CR, Chen WT, Tang YH, Hwang BF. Fine particulate matter exposure during pregnancy and infancy and incident asthma. *J Allergy Clin Immunol.* 2019;143(6):2254–62 e5. [PubMed: 30959062]
103. Castro-Giner F, Kunzli N, Jacquemin B, Forsberg B, de Cid R, Sunyer J, et al. Traffic-related air pollution, oxidative stress genes, and asthma (ECHRS). *Environ Health Perspect.* 2009;117(12):1919–24. [PubMed: 20049212]
104. Oftedal B, Nystad W, Brunekreef B, Nafstad P. Long-term traffic-related exposures and asthma onset in schoolchildren in oslo, norway. *Environ Health Perspect.* 2009;117(5):839–44. [PubMed: 19478970]
105. Molter A, Agius R, de Vocht F, Lindley S, Gerrard W, Custovic A, et al. Effects of long-term exposure to PM10 and NO2 on asthma and wheeze in a prospective birth cohort. *J Epidemiol Community Health.* 2014;68(1):21–8. [PubMed: 23999376]
106. Molter A, Simpson A, Berdel D, Brunekreef B, Custovic A, Cyrus J, et al. A multicentre study of air pollution exposure and childhood asthma prevalence: the ESCAPE project. *Eur Respir J.* 2015;45(3):610–24. [PubMed: 25323237]
107. Lindgren A, Strohm E, Bjork J, Jakobsson K. Asthma incidence in children growing up close to traffic: a registry-based birth cohort. *Environ Health.* 2013;12:91. [PubMed: 24160449]
108. Kramer U, Sugiri D, Ranft U, Krutmann J, von Berg A, Berdel D, et al. Eczema, respiratory allergies, and traffic-related air pollution in birth cohorts from small-town areas. *J Dermatol Sci.* 2009;56(2):99–105. [PubMed: 19713084]
109. Fuertes E, Standl M, Cyrus J, Berdel D, von Berg A, Bauer CP, et al. A longitudinal analysis of associations between traffic-related air pollution with asthma, allergies and sensitization in the GINIplus and LISAPLUS birth cohorts. *PeerJ.* 2013;1:e193. [PubMed: 24255809]
110. Schultz ES, Gruzieva O, Bellander T, Bottai M, Hallberg J, Kull I, et al. Traffic-related air pollution and lung function in children at 8 years of age: a birth cohort study. *Am J Respir Crit Care Med.* 2012;186(12):1286–91. [PubMed: 23103735]

111. Citerne A, Roda C, Viola M, Ranciere F, Momas I. Early postnatal exposure to traffic-related air pollution and asthma in adolescents: vulnerability factors in the PARIS birth cohort. *Environ Res.* 2021;201:111473. [PubMed: 34116015]
112. Lau N, Norman A, Smith MJ, Sarkar A, Gao Z. Association between Traffic Related Air Pollution and the Development of Asthma Phenotypes in Children: A Systematic Review. *Int J Chronic Dis.* 2018;2018:4047386. [PubMed: 30631772]
113. Bettiol A, Gelain E, Milanesio E, Asta F, Rusconi F. The first 1000 days of life: traffic-related air pollution and development of wheezing and asthma in childhood. A systematic review of birth cohort studies. *Environ Health.* 2021;20(1):46. [PubMed: 33865406]
114. Houston D, Li W, Wu J. Disparities in exposure to automobile and truck traffic and vehicle emissions near the Los Angeles-Long Beach port complex. *Am J Public Health.* 2014;104(1):156–64. [PubMed: 23678919]
115. Weaver GM, Gauderman WJ. Traffic-Related Pollutants: Exposure and Health Effects Among Hispanic Children. *Am J Epidemiol.* 2018;187(1):45–52. [PubMed: 28605396]
116. Hauptman M, Gaffin JM, Petty CR, Sheehan WJ, Lai PS, Coull B, et al. Proximity to major roadways and asthma symptoms in the School Inner-City Asthma Study. *J Allergy Clin Immunol.* 2020;145(1):119–26 e4. [PubMed: 31557500]
117. Kinney PL, Aggarwal M, Northridge ME, Janssen NA, Shepard P. Airborne concentrations of PM(2.5) and diesel exhaust particles on Harlem sidewalks: a community-based pilot study. *Environ Health Perspect.* 2000;108(3):213–8. [PubMed: 10706526]
118. O’Neill MS, Zanobetti A, Schwartz J. Disparities by race in heat-related mortality in four US cities: the role of air conditioning prevalence. *J Urban Health.* 2005;82(2):191–7. [PubMed: 15888640]
119. O’Lenick C, Chang HH, Kramer MR, Winquist A, Mulholland JA, Friberg MD, et al. Ozone and childhood respiratory disease in three US cities: evaluation of effect measure modification by neighborhood socioeconomic status using a Bayesian hierarchical approach. *Environ Health.* 2017;16(1):36. [PubMed: 28381221]
120. O’Lenick CR, Winquist A, Mulholland JA, Friberg MD, Chang HH, Kramer MR, et al. Assessment of neighbourhood-level socioeconomic status as a modifier of air pollution-asthma associations among children in Atlanta. *J Epidemiol Community Health.* 2017;71(2):129–36. [PubMed: 27422981]
121. Humphrey JL, Lindstrom M, Barton KE, Shrestha PM, Carlton EJ, Adgate JL, et al. Social and Environmental Neighborhood Typologies and Lung Function in a Low-Income, Urban Population. *Int J Environ Res Public Health.* 2019;16(7).
122. Kravitz-Wirtz N, Teixeira S, Hajat A, Woo B, Crowder K, Takeuchi D. Early-Life Air Pollution Exposure, Neighborhood Poverty, and Childhood Asthma in the United States, 1990(–)2014. *Int J Environ Res Public Health.* 2018;15(6).
123. Hughes HK, Matsui EC, Tschudy MM, Pollack CE, Keet CA. Pediatric Asthma Health Disparities: Race, Hardship, Housing, and Asthma in a National Survey. *Acad Pediatr.* 2017;17(2):127–34. [PubMed: 27876585]
124. Corburn J, Osleeb J, Porter M. Urban asthma and the neighbourhood environment in New York City. *Health Place.* 2006;12(2):167–79. [PubMed: 16338632]
125. Beck AF, Huang B, Chundur R, Kahn RS. Housing code violation density associated with emergency department and hospital use by children with asthma. *Health Aff (Millwood).* 2014;33(11):1993–2002. [PubMed: 25367995]
126. Bryant-Stephens TC, Strane D, Robinson EK, Bhambhani S, Kenyon CC. Housing and asthma disparities. *J Allergy Clin Immunol.* 2021;148(5):1121–9. [PubMed: 34599980]
127. Call RS, Smith TF, Morris E, Chapman MD, Platts-Mills TA. Risk factors for asthma in inner city children. *J Pediatr.* 1992;121(6):862–6. [PubMed: 1447646]
128. Sarpong SB, Hamilton RG, Eggleston PA, Adkinson NF Jr. Socioeconomic status and race as risk factors for cockroach allergen exposure and sensitization in children with asthma. *J Allergy Clin Immunol.* 1996;97(6):1393–401. [PubMed: 8648037]

129. Eggleston PA, Rosenstreich D, Lynn H, Gergen P, Baker D, Kattan M, et al. Relationship of indoor allergen exposure to skin test sensitivity in inner-city children with asthma. *J Allergy Clin Immunol.* 1998;102(4 Pt 1):563–70. [PubMed: 9802363]
130. Rosenstreich DL, Eggleston P, Kattan M, Baker D, Slavin RG, Gergen P, et al. The role of cockroach allergy and exposure to cockroach allergen in causing morbidity among inner-city children with asthma. *N Engl J Med.* 1997;336(19):1356–63. [PubMed: 9134876]
131. Cohn RD, Arbes SJ Jr., Yin M, Jaramillo R, Zeldin DC. National prevalence and exposure risk for mouse allergen in US households. *J Allergy Clin Immunol.* 2004;113(6):1167–71. [PubMed: 15208600]
132. Litonjua AA, Carey VJ, Burge HA, Weiss ST, Gold DR. Exposure to cockroach allergen in the home is associated with incident doctor-diagnosed asthma and recurrent wheezing. *J Allergy Clin Immunol.* 2001;107(1):41–7. [PubMed: 11149989]
133. Sarinho E, Schor D, Veloso MA, Rizzo JA. There are more asthmatics in homes with high cockroach infestation. *Braz J Med Biol Res.* 2004;37(4):503–10. [PubMed: 15064813]
134. Silva JM, Camara AA, Tobias KR, Macedo IS, Cardoso MR, Arruda E, et al. A prospective study of wheezing in young children: the independent effects of cockroach exposure, breast-feeding and allergic sensitization. *Pediatr Allergy Immunol.* 2005;16(5):393–401. [PubMed: 16101931]
135. Donohue KM, Al-alem U, Perzanowski MS, Chew GL, Johnson A, Divjan A, et al. Anti-cockroach and anti-mouse IgE are associated with early wheeze and atopy in an inner-city birth cohort. *J Allergy Clin Immunol.* 2008;122(5):914–20. [PubMed: 19000580]
136. Rabito FA, Carlson J, Holt EW, Iqbal S, James MA. Cockroach exposure independent of sensitization status and association with hospitalizations for asthma in inner-city children. *Ann Allergy Asthma Immunol.* 2011;106(2):103–9. [PubMed: 21277511]
137. Gruchalla RS, Pongracic J, Plaut M, Evans R 3rd, Visness CM, Walter M, et al. Inner City Asthma Study: relationships among sensitivity, allergen exposure, and asthma morbidity. *J Allergy Clin Immunol.* 2005;115(3):478–85. [PubMed: 15753892]
138. Phipatanakul W, Litonjua AA, Platts-Mills TA, Naccara LM, Celedon JC, Abdulkerim H, et al. Sensitization to mouse allergen and asthma and asthma morbidity among women in Boston. *J Allergy Clin Immunol.* 2007;120(4):954–6. [PubMed: 17590423]
139. Grant T, Aloe C, Perzanowski M, Phipatanakul W, Bollinger ME, Miller R, et al. Mouse Sensitization and Exposure Are Associated with Asthma Severity in Urban Children. *J Allergy Clin Immunol Pract.* 2017;5(4):1008–14 e1. [PubMed: 27923647]
140. Sheehan WJ, Permaul P, Petty CR, Coull BA, Baxi SN, Gaffin JM, et al. Association Between Allergen Exposure in Inner-City Schools and Asthma Morbidity Among Students. *JAMA Pediatr.* 2017;171(1):31–8. [PubMed: 27893060]
141. Matsui EC, Eggleston PA, Buckley TJ, Krishnan JA, Breyse PN, Rand CS, et al. Household mouse allergen exposure and asthma morbidity in inner-city preschool children. *Ann Allergy Asthma Immunol.* 2006;97(4):514–20. [PubMed: 17069107]
142. Pongracic JA, Visness CM, Gruchalla RS, Evans R 3rd, Mitchell HE. Effect of mouse allergen and rodent environmental intervention on asthma in inner-city children. *Ann Allergy Asthma Immunol.* 2008;101(1):35–41. [PubMed: 18681082]
143. Ahluwalia SK, Peng RD, Breyse PN, Diette GB, Curtin-Brosnan J, Aloe C, et al. Mouse allergen is the major allergen of public health relevance in Baltimore City. *J Allergy Clin Immunol.* 2013;132(4):830–5 e1–2. [PubMed: 23810154]
144. Torjusen EN, Diette GB, Breyse PN, Curtin-Brosnan J, Aloe C, Matsui EC. Dose-response relationships between mouse allergen exposure and asthma morbidity among urban children and adolescents. *Indoor Air.* 2013;23(4):268–74. [PubMed: 23067271]
145. Matsui EC, Sampson HA, Bahnson HT, Gruchalla RS, Pongracic JA, Teach SJ, et al. Allergen-specific IgE as a biomarker of exposure plus sensitization in inner-city adolescents with asthma. *Allergy.* 2010;65(11):1414–22. [PubMed: 20560910]
146. Sedaghat AR, Matsui EC, Baxi SN, Bollinger ME, Miller R, Perzanowski M, et al. Mouse Sensitivity is an Independent Risk Factor for Rhinitis in Children with Asthma. *J Allergy Clin Immunol Pract.* 2016;4(1):82–8 e1. [PubMed: 26441149]

147. Matsui EC, Perzanowski M, Peng RD, Wise RA, Balcer-Whaley S, Newman M, et al. Effect of an Integrated Pest Management Intervention on Asthma Symptoms Among Mouse-Sensitized Children and Adolescents With Asthma: A Randomized Clinical Trial. *JAMA*. 2017;317(10):1027–36. [PubMed: 28264080]
148. Grant T, Phipatanakul W, Perzanowski M, Balcer-Whaley S, Peng RD, Curtin-Brosnan J, et al. Reduction in mouse allergen exposure is associated with greater lung function growth. *J Allergy Clin Immunol*. 2020;145(2):646–53 e1. [PubMed: 31866099]
149. Sharpe R, Thornton CR, Osborne NJ. Modifiable factors governing indoor fungal diversity and risk of asthma. *Clin Exp Allergy*. 2014;44(5):631–41. [PubMed: 24471926]
150. Peat JK, Dickerson J, Li J. Effects of damp and mould in the home on respiratory health: a review of the literature. *Allergy*. 1998;53(2):120–8. [PubMed: 9534909]
151. Bjornsson E, Norback D, Janson C, Widstrom J, Palmgren U, Strom G, et al. Asthmatic symptoms and indoor levels of micro-organisms and house dust mites. *Clin Exp Allergy*. 1995;25(5):423–31. [PubMed: 7553245]
152. Gent JF, Ren P, Belanger K, Triche E, Bracken MB, Holford TR, et al. Levels of household mold associated with respiratory symptoms in the first year of life in a cohort at risk for asthma. *Environ Health Perspect*. 2002;110(12):A781–6. [PubMed: 12460818]
153. Belanger K, Beckett W, Triche E, Bracken MB, Holford T, Ren P, et al. Symptoms of wheeze and persistent cough in the first year of life: associations with indoor allergens, air contaminants, and maternal history of asthma. *Am J Epidemiol*. 2003;158(3):195–202. [PubMed: 12882940]
154. Bornehag CG, Blomquist G, Gyntelberg F, Jarvholm B, Malmberg P, Nordvall L, et al. Dampness in buildings and health. Nordic interdisciplinary review of the scientific evidence on associations between exposure to “dampness” in buildings and health effects (NORDDAMP). *Indoor Air*. 2001;11(2):72–86. [PubMed: 11394014]
155. Su HJ, Rotnitzky A, Burge HA, Spengler JD. Examination of fungi in domestic interiors by using factor analysis: correlations and associations with home factors. *Appl Environ Microbiol*. 1992;58(1):181–6. [PubMed: 1539973]
156. Park JH, Kreiss K, Cox-Ganser JM. Rhinosinusitis and mold as risk factors for asthma symptoms in occupants of a water-damaged building. *Indoor Air*. 2012;22(5):396–404. [PubMed: 22385263]
157. Borrás-Santos A, Jacobs JH, Taubel M, Haverinen-Shaughnessy U, Krop EJ, Huttunen K, et al. Dampness and mould in schools and respiratory symptoms in children: the HITEA study. *Occup Environ Med*. 2013;70(10):681–7. [PubMed: 23775866]
158. Jacobs J, Borrás-Santos A, Krop E, Taubel M, Leppanen H, Haverinen-Shaughnessy U, et al. Dampness, bacterial and fungal components in dust in primary schools and respiratory health in schoolchildren across Europe. *Occup Environ Med*. 2014;71(10):704–12. [PubMed: 25035116]
159. Salo PM, Arbes SJ Jr., Sever M, Jaramillo R, Cohn RD, London SJ, et al. Exposure to *Alternaria alternata* in US homes is associated with asthma symptoms. *J Allergy Clin Immunol*. 2006;118(4):892–8. [PubMed: 17030243]
160. Baxi SN, Sheehan WJ, Sordillo JE, Muilenberg ML, Rogers CA, Gaffin JM, et al. Association between fungal spore exposure in inner-city schools and asthma morbidity. *Ann Allergy Asthma Immunol*. 2019;122(6):610–5 e1. [PubMed: 30904580]
161. Verhoeff AP, Burge HA. Health risk assessment of fungi in home environments. *Ann Allergy Asthma Immunol*. 1997;78(6):544–54; quiz 55–6. [PubMed: 9207717]
162. Krieger J, Higgins DL. Housing and health: time again for public health action. *Am J Public Health*. 2002;92(5):758–68. [PubMed: 11988443]
163. Jacobs DE. Environmental health disparities in housing. *Am J Public Health*. 2011;101 Suppl 1:S115–22. [PubMed: 21551378]
164. Housing Assistance Council. Taking Stock: Rural People, Poverty, and Housing in the 21st Century. 2012.
165. Cheryan S, Ziegler SA, Plaut VC, Meltzoff AN. Designing Classrooms to Maximize Student Achievement. *Policy Insights from the Behavioral and Brain Sciences*. 2014;1(1):4–12.

166. Lewis SA, Weiss ST, Platts-Mills TA, Syring M, Gold DR. Association of specific allergen sensitization with socioeconomic factors and allergic disease in a population of Boston women. *J Allergy Clin Immunol.* 2001;107(4):615–22. [PubMed: 11295648]
167. Camacho-Rivera M, Kawachi I, Bennett GG, Subramanian SV. Associations of neighborhood concentrated poverty, neighborhood racial/ethnic composition, and indoor allergen exposures: a cross-sectional analysis of los angeles households, 2006–2008. *J Urban Health.* 2014;91(4):661–76. [PubMed: 24771244]
168. Matsui EC. Management of rodent exposure and allergy in the pediatric population. *Curr Allergy Asthma Rep.* 2013;13(6):681–6. [PubMed: 23912589]
169. Cho SJ, Ramachandran G, Grengs J, Ryan AD, Eberly LE, Adgate JL. Longitudinal evaluation of allergen and culturable fungal concentrations in inner-city households. *J Occup Environ Hyg.* 2008;5(2):107–18. [PubMed: 18085481]
170. Sarpong SB, Wood RA, Karrison T, Eggleston PA. Cockroach allergen (Bla g 1) in school dust. *J Allergy Clin Immunol.* 1997;99(4):486–92. [PubMed: 9111492]
171. Chew GL, Correa JC, Perzanowski MS. Mouse and cockroach allergens in the dust and air in northeastern United States inner-city public high schools. *Indoor Air.* 2005;15(4):228–34. [PubMed: 15982269]
172. Permaul P, Hoffman E, Fu C, Sheehan W, Baxi S, Gaffin J, et al. Allergens in urban schools and homes of children with asthma. *Pediatr Allergy Immunol.* 2012;23(6):543–9. [PubMed: 22672325]
173. Sheehan WJ, Rangsihienchai PA, Muilenberg ML, Rogers CA, Lane JP, Ghaemghami J, et al. Mouse allergens in urban elementary schools and homes of children with asthma. *Ann Allergy Asthma Immunol.* 2009;102(2):125–30. [PubMed: 19230463]
174. Baxi SN, Muilenberg ML, Rogers CA, Sheehan WJ, Gaffin J, Permaul P, et al. Exposures to molds in school classrooms of children with asthma. *Pediatr Allergy Immunol.* 2013;24(7):697–703. [PubMed: 24112429]
175. Pongracic JA, O'Connor GT, Muilenberg ML, Vaughn B, Gold DR, Kattan M, et al. Differential effects of outdoor versus indoor fungal spores on asthma morbidity in inner-city children. *J Allergy Clin Immunol.* 2010;125(3):593–9. [PubMed: 20132971]
176. Stevenson LA, Gergen PJ, Hoover DR, Rosenstreich D, Mannino DM, Matte TD. Sociodemographic correlates of indoor allergen sensitivity among United States children. *J Allergy Clin Immunol.* 2001;108(5):747–52. [PubMed: 11692099]
177. Franklin JM, Grunwell JR, Bruce AC, Smith RC, Fitzpatrick AM. Predictors of emergency department use in children with persistent asthma in metropolitan Atlanta, Georgia. *Ann Allergy Asthma Immunol.* 2017;119(2):129–36. [PubMed: 28479192]
178. Gaffin JM, Hauptman M, Petty CR, Sheehan WJ, Lai PS, Wolfson JM, et al. Nitrogen dioxide exposure in school classrooms of inner-city children with asthma. *J Allergy Clin Immunol.* 2018;141(6):2249–55 e2. [PubMed: 28988796]
179. Institute of Medicine. *Clearing the Air: Asthma and Indoor Air Exposures.* Washington DC: The National Academies Press; 2000.
180. U.S. Environmental Protection Agency. Report to Congress on indoor air quality. Washington, DC: 1989. Contract No.: EPA/400/1–89/001C.
181. Zota A, Adamkiewicz G, Levy JI, Spengler JD. Ventilation in public housing: implications for indoor nitrogen dioxide concentrations. *Indoor Air.* 2005;15(6):393–401. [PubMed: 16268829]
182. Dekker C, Dales R, Bartlett S, Brunekreef B, Zwanenburg H. Childhood asthma and the indoor environment. *Chest.* 1991;100(4):922–6. [PubMed: 1914606]
183. Volkmer RE, Ruffin RE, Wigg NR, Davies N. The prevalence of respiratory symptoms in South Australian preschool children. II. Factors associated with indoor air quality. *J Paediatr Child Health.* 1995;31(2):116–20. [PubMed: 7794611]
184. Belanger K, Holford TR, Gent JF, Hill ME, Kezik JM, Leaderer BP. Household levels of nitrogen dioxide and pediatric asthma severity. *Epidemiology.* 2013;24(2):320–30. [PubMed: 23337243]
185. Kattan M, Gergen PJ, Eggleston P, Visness CM, Mitchell HE. Health effects of indoor nitrogen dioxide and passive smoking on urban asthmatic children. *J Allergy Clin Immunol.* 2007;120(3):618–24. [PubMed: 17582483]

186. Permaul P, Gaffin JM, Petty CR, Baxi SN, Lai PS, Sheehan WJ, et al. Obesity may enhance the adverse effects of NO₂ exposure in urban schools on asthma symptoms in children. *J Allergy Clin Immunol*. 2020;146(4):813–20 e2. [PubMed: 32197971]
187. Belanger K, Gent JF, Triche EW, Bracken MB, Leaderer BP. Association of indoor nitrogen dioxide exposure with respiratory symptoms in children with asthma. *Am J Respir Crit Care Med*. 2006;173(3):297–303. [PubMed: 16254270]
188. Mohsenin V. Airway responses to nitrogen dioxide in asthmatic subjects. *J Toxicol Environ Health*. 1987;22(4):371–80. [PubMed: 3320381]
189. Salome CM, Brown NJ, Marks GB, Woolcock AJ, Johnson GM, Nancarrow PC, et al. Effect of nitrogen dioxide and other combustion products on asthmatic subjects in a home-like environment. *Eur Respir J*. 1996;9(5):910–8. [PubMed: 8793451]
190. Kanchongkittiphon W, Mendell MJ, Gaffin JM, Wang G, Phipatanakul W. Indoor environmental exposures and exacerbation of asthma: an update to the 2000 review by the Institute of Medicine. *Environ Health Perspect*. 2015;123(1):6–20. [PubMed: 25303775]
191. McGwin G, Lienert J, Kennedy JI. Formaldehyde exposure and asthma in children: a systematic review. *Environ Health Perspect*. 2010;118(3):313–7. [PubMed: 20064771]
192. Centers for Disease Control and Prevention. Current Cigarette Smoking Among Adults in the United States: 2020 [updated December 10, 2020. Available from: https://www.cdc.gov/tobacco/data_statistics/fact_sheets/adult_data/cig_smoking/index.htm.
193. Singh GK, Siahpush M, Kogan MD. Disparities in children's exposure to environmental tobacco smoke in the United States, 2007. *Pediatrics*. 2010;126(1):4–13. [PubMed: 20587673]
194. Abell TD, Baker LC, Ramsey CN Jr. The effects of maternal smoking on infant birth weight. *Fam Med*. 1991;23(2):103–7. [PubMed: 2037208]
195. Milner AD, Marsh MJ, Ingram DM, Fox GF, Susiva C. Effects of smoking in pregnancy on neonatal lung function. *Arch Dis Child Fetal Neonatal Ed*. 1999;80(1):F8–14. [PubMed: 10325804]
196. Sherwood RA, Keating J, Kavvadia V, Greenough A, Peters TJ. Substance misuse in early pregnancy and relationship to fetal outcome. *Eur J Pediatr*. 1999;158(6):488–92. [PubMed: 10378398]
197. Wang X, Tager IB, Van Vunakis H, Speizer FE, Hanrahan JP. Maternal smoking during pregnancy, urine cotinine concentrations, and birth outcomes. A prospective cohort study. *Int J Epidemiol*. 1997;26(5):978–88. [PubMed: 9363518]
198. Ko TJ, Tsai LY, Chu LC, Yeh SJ, Leung C, Chen CY, et al. Parental smoking during pregnancy and its association with low birth weight, small for gestational age, and preterm birth offspring: a birth cohort study. *Pediatr Neonatol*. 2014;55(1):20–7. [PubMed: 23850094]
199. Windham GC, Hopkins B, Fenster L, Swan SH. Prenatal active or passive tobacco smoke exposure and the risk of preterm delivery or low birth weight. *Epidemiology*. 2000;11(4):427–33. [PubMed: 10874550]
200. Lodrup Carlsen KC, Jaakkola JJ, Nafstad P, Carlsen KH. In utero exposure to cigarette smoking influences lung function at birth. *Eur Respir J*. 1997;10(8):1774–9. [PubMed: 9272918]
201. Castro-Rodriguez JA, Forno E, Rodriguez-Martinez CE, Celedon JC. Risk and Protective Factors for Childhood Asthma: What Is the Evidence? *J Allergy Clin Immunol Pract*. 2016;4(6):1111–22. [PubMed: 27286779]
202. Perzanowski MS, Divjan A, Mellins RB, Canfield SM, Rosa MJ, Chew GL, et al. Exhaled NO among inner-city children in New York City. *J Asthma*. 2010;47(9):1015–21. [PubMed: 20936992]
203. Tariq SM, Matthews SM, Hakim EA, Stevens M, Arshad SH, Hide DW. The prevalence of and risk factors for atopy in early childhood: a whole population birth cohort study. *J Allergy Clin Immunol*. 1998;101(5):587–93. [PubMed: 9600493]
204. Vork KL, Broadwin RL, Blaisdell RJ. Developing asthma in childhood from exposure to secondhand tobacco smoke: insights from a meta-regression. *Environ Health Perspect*. 2007;115(10):1394–400. [PubMed: 17938726]

205. Sturm JJ, Yeatts K, Loomis D. Effects of tobacco smoke exposure on asthma prevalence and medical care use in North Carolina middle school children. *Am J Public Health*. 2004;94(2):308–13. [PubMed: 14759947]
206. Cook DG, Strachan DP. Health effects of passive smoking-10: Summary of effects of parental smoking on the respiratory health of children and implications for research. *Thorax*. 1999;54(4):357–66. [PubMed: 10092699]
207. Chapman RS, Hadden WC, Perlin SA. Influences of asthma and household environment on lung function in children and adolescents: the third national health and nutrition examination survey. *Am J Epidemiol*. 2003;158(2):175–89. [PubMed: 12851231]
208. Soussan D, Liard R, Zureik M, Tournon D, Rogeaux Y, Neukirch F. Treatment compliance, passive smoking, and asthma control: a three year cohort study. *Arch Dis Child*. 2003;88(3):229–33. [PubMed: 12598385]
209. Wang HC, McGeady SJ, Yousef E. Patient, home residence, and neighborhood characteristics in pediatric emergency department visits for asthma. *J Asthma*. 2007;44(2):95–8. [PubMed: 17454322]
210. Lannero E, Wickman M, van Hage M, Bergstrom A, Pershagen G, Nordvall L. Exposure to environmental tobacco smoke and sensitisation in children. *Thorax*. 2008;63(2):172–6. [PubMed: 18089631]
211. Saulyte J, Regueira C, Montes-Martinez A, Khudyakov P, Takkouche B. Active or passive exposure to tobacco smoking and allergic rhinitis, allergic dermatitis, and food allergy in adults and children: a systematic review and meta-analysis. *PLoS Med*. 2014;11(3):e1001611. [PubMed: 24618794]
212. Eriksson J, Ekerljung L, Sundblad BM, Lotvall J, Toren K, Ronmark E, et al. Cigarette smoking is associated with high prevalence of chronic rhinitis and low prevalence of allergic rhinitis in men. *Allergy*. 2013;68(3):347–54. [PubMed: 23346908]
213. Topp R, Thefeld W, Wichmann HE, Heinrich J. The effect of environmental tobacco smoke exposure on allergic sensitization and allergic rhinitis in adults. *Indoor Air*. 2005;15(4):222–7. [PubMed: 15982268]
214. Lee CH, Chuang HY, Hong CH, Huang SK, Chang YC, Ko YC, et al. Lifetime exposure to cigarette smoking and the development of adult-onset atopic dermatitis. *Br J Dermatol*. 2011;164(3):483–9. [PubMed: 21054333]
215. Lewis J, Hernandez D, & Geronimus AT. Energy efficiency as energy justice: addressing racial inequities through investments in people and places. *Energy Efficiency*. 2020;13:419–32.
216. Adua LSJ. Explaining residential energy consumption: A focus on location and race differences in natural gas use. *Journal of Rural Social Sciences*. 2011;26(1).
217. Goldstein B, Reames TG, Newell JP. Racial inequity in household energy efficiency and carbon emissions in the United States: An emissions paradox. *Energy Research & Social Science*. 2022;84.
218. U.S. Census Bureau. Quarterly Residential Vacancies and Homeownership, Fourth Quarter 2021 (Release number: CB22-10). 2022.
219. D'Souza J, Jia C, Mukherjee B, Batterman S. Ethnicity, housing and personal factors as determinants of VOC exposures. *Atmospheric Environment*. 2009;43:2884–92.
220. Yang TC, Shoff C, Noah AJ, Black N, Sparks CS. Racial segregation and maternal smoking during pregnancy: a multilevel analysis using the racial segregation interaction index. *Soc Sci Med*. 2014;107:26–36. [PubMed: 24602968]
221. Hunt BR, Whitman S. Maternal smoking in Chicago: a community-level analysis. *J Health Care Poor Underserved*. 2011;22(1):194–210. [PubMed: 21317515]
222. Martin LT, McNamara M, Milot A, Bloch M, Hair EC, Halle T. Correlates of smoking before, during, and after pregnancy. *Am J Health Behav*. 2008;32(3):272–82. [PubMed: 18067467]
223. Merianos AL, Jandarov RA, Choi K, Mahabee-Gittens EM. Tobacco smoke exposure disparities persist in U.S. children: NHANES 1999–2014. *Prev Med*. 2019;123:138–42. [PubMed: 30902698]
224. Wilson KM, Klein JD, Blumkin AK, Gottlieb M, Winickoff JP. Tobacco-smoke exposure in children who live in multiunit housing. *Pediatrics*. 2011;127(1):85–92. [PubMed: 21149434]

225. Jaffee KD, Liu GC, Canty-Mitchell J, Qi RA, Austin J, Swigonski N. Race, urban community stressors, and behavioral and emotional problems of children with special health care needs. *Psychiatr Serv*. 2005;56(1):63–9. [PubMed: 15637194]
226. Jutte DP, Miller JL, Erickson DJ. Neighborhood adversity, child health, and the role for community development. *Pediatrics*. 2015;135 Suppl 2:S48–57. [PubMed: 25733725]
227. Rodriguez EM, Pollack CE, Keet C, Peng RD, Balcer-Whaley S, Custer J, et al. Neighborhoods, Caregiver Stress, and Children’s Asthma Symptoms. *J Allergy Clin Immunol Pract*. 2021.
228. van de Loo KF, van Gelder MM, Roukema J, Roeleveld N, Merkus PJ, Verhaak CM. Prenatal maternal psychological stress and childhood asthma and wheezing: a meta-analysis. *Eur Respir J*. 2016;47(1):133–46. [PubMed: 26541526]
229. Kopel LS, Gaffin JM, Ozonoff A, Rao DR, Sheehan WJ, Friedlander JL, et al. Perceived neighborhood safety and asthma morbidity in the school inner-city asthma study. *Pediatr Pulmonol*. 2015;50(1):17–24. [PubMed: 24421055]
230. Kopel LS, Petty CR, Gaffin JM, Sheehan WJ, Baxi SN, Kanchongkittiphon W, et al. Caregiver stress among inner-city school children with asthma. *J Allergy Clin Immunol Pract*. 2017;5(4):1132–4 e3. [PubMed: 28433340]
231. Brehm JM, Ramratnam SK, Tse SM, Croteau-Chonka DC, Pino-Yanes M, Rosas-Salazar C, et al. Stress and Bronchodilator Response in Children with Asthma. *Am J Respir Crit Care Med*. 2015;192(1):47–56. [PubMed: 25918834]
232. Bartlett SJ, Krishnan JA, Riekert KA, Butz AM, Malveaux FJ, Rand CS. Maternal depressive symptoms and adherence to therapy in inner-city children with asthma. *Pediatrics*. 2004;113(2):229–37. [PubMed: 14754931]
233. Oren E, Gerald L, Stern DA, Martinez FD, Wright AL. Self-Reported Stressful Life Events During Adolescence and Subsequent Asthma: A Longitudinal Study. *J Allergy Clin Immunol Pract*. 2017;5(2):427–34 e2. [PubMed: 27815066]
234. Wing R, Gjelsvik A, Nocera M, McQuaid EL. Association between adverse childhood experiences in the home and pediatric asthma. *Ann Allergy Asthma Immunol*. 2015;114(5):379–84. [PubMed: 25843164]
235. Cohen RT, Canino GJ, Bird HR, Celedon JC. Violence, abuse, and asthma in Puerto Rican children. *Am J Respir Crit Care Med*. 2008;178(5):453–9. [PubMed: 18556622]
236. Sternthal MJ, Jun HJ, Earls F, Wright RJ. Community violence and urban childhood asthma: a multilevel analysis. *Eur Respir J*. 2010;36(6):1400–9. [PubMed: 20413538]
237. Wright RJ, Mitchell H, Visness CM, Cohen S, Stout J, Evans R, et al. Community violence and asthma morbidity: the Inner-City Asthma Study. *Am J Public Health*. 2004;94(4):625–32. [PubMed: 15054016]
238. Tonorezos ES, Breyse PN, Matsui EC, McCormack MC, Curtin-Brosnan J, Williams D, et al. Does neighborhood violence lead to depression among caregivers of children with asthma? *Soc Sci Med*. 2008;67(1):31–7. [PubMed: 18406503]
239. Bartlett SJ, Kolodner K, Butz AM, Eggleston P, Malveaux FJ, Rand CS. Maternal depressive symptoms and emergency department use among inner-city children with asthma. *Arch Pediatr Adolesc Med*. 2001;155(3):347–53. [PubMed: 11231800]
240. Williams D. Race, stress, and mental health: Findings from the Commonwealth Minority Health Survey. Baltimore, MD; 2000.
241. Carliner H, Gary D, McLaughlin KA, Keyes KM. Trauma Exposure and Externalizing Disorders in Adolescents: Results From the National Comorbidity Survey Adolescent Supplement. *J Am Acad Child Adolesc Psychiatry*. 2017;56(9):755–64 e3. [PubMed: 28838580]
242. Sheats KJ, Irving SM, Mercy JA, Simon TR, Crosby AE, Ford DC, et al. Violence-Related Disparities Experienced by Black Youth and Young Adults: Opportunities for Prevention. *Am J Prev Med*. 2018;55(4):462–9. [PubMed: 30139709]
243. Redelings M, Lieb L, Sorvillo F. Years off your life? The effects of homicide on life expectancy by neighborhood and race/ethnicity in Los Angeles county. *J Urban Health*. 2010;87(4):670–6. [PubMed: 20556528]
244. Schubiner H, Scott R, Tzelepis A. Exposure to violence among inner-city youth. *J Adolesc Health*. 1993;14(3):214–9. [PubMed: 8323933]

245. Zarei K, Kahle L, Buckman DW, Choi K, Williams F. Parent-child nativity, race, ethnicity, and adverse childhood experiences among United States children. *J Pediatr*. 2022.
246. Kurani S, Webb L, Cadet K, Ma M, Gibson M, Jallah N, et al. Area-level deprivation and adverse childhood experiences among high school students in Maryland. *BMC Public Health*. 2022;22(1):811. [PubMed: 35459200]
247. Torres-Harding S, Turner T. Assessing Racial Microaggression Distress in a Diverse Sample. *Eval Health Prof*. 2015;38(4):464–90. [PubMed: 25237154]
248. Coogan PF, Yu J, O'Connor GT, Brown TA, Cozier YC, Palmer JR, et al. Experiences of racism and the incidence of adult-onset asthma in the Black Women's Health Study. *Chest*. 2014;145(3):480–5. [PubMed: 23887828]
249. Thakur N, Barcelo NE, Borrell LN, Singh S, Eng C, Davis A, et al. Perceived Discrimination Associated With Asthma and Related Outcomes in Minority Youth: The GALA II and SAGE II Studies. *Chest*. 2017;151(4):804–12. [PubMed: 27916618]
250. Institute of Medicine. *Unequal Treatment: Confronting Racial and Ethnic Disparities in Health Care*. Washington DC: The National Academies Press; 2003.
251. Lieu TA, Newacheck PW, McManus MA. Race, ethnicity, and access to ambulatory care among US adolescents. *Am J Public Health*. 1993;83(7):960–5. [PubMed: 8328617]
252. Kindig DA, Movassaghi H, Dunham NC, Zwick DI, Taylor CM. Trends in physician availability in 10 urban areas from 1963 to 1980. *Inquiry*. 1987;24(2):136–46. [PubMed: 2954910]
253. Watson S. *Health Care in the Inner City: Asking the Right Question*. NC L Rev 1647. 1993;71(5).
254. Collins KSHA, Neuhaus C. *U.S. Minority Health: A Chartbook*. New York: The Commonwealth Fund; 1999.
255. Lillie-Blanton MMR, Salganicoff A. Site of medical care: Do racial and ethnic differences persist? *Yale Journal of Health Policy, Law, and Ethics*. 2001;1(1):1–17. [PubMed: 12669318]
256. Greek AA, Kieckhefer GM, Kim H, Joesch JM, Baydar N. Family perceptions of the usual source of care among children with asthma by race/ethnicity, language, and family income. *J Asthma*. 2006;43(1):61–9. [PubMed: 16448968]
257. Qato DMDM, Wilder J, Lee T, Qato D, Lambert B. 'Pharmacy Deserts' are prevalent in Chicago's predominantly minority communities, raising medication access concerns. *Health Affairs*. 2014;33(11):1958–65. [PubMed: 25367990]
258. Riekert KA, Butz AM, Eggleston PA, Huss K, Winkelstein M, Rand CS. Caregiver-physician medication concordance and undertreatment of asthma among inner-city children. *Pediatrics*. 2003;111(3):e214–20. [PubMed: 12612274]
259. van Ryn M, Burke J. The effect of patient race and socio-economic status on physicians' perceptions of patients. *Soc Sci Med*. 2000;50(6):813–28. [PubMed: 10695979]
260. Okelo SO, Wu AW, Merriman B, Krishnan JA, Diette GB. Are physician estimates of asthma severity less accurate in black than in white patients? *J Gen Intern Med*. 2007;22(7):976–81. [PubMed: 17453263]
261. Cyr ME, Etchin AG, Guthrie BJ, Benneyan JC. Access to specialty healthcare in urban versus rural US populations: a systematic literature review. *BMC Health Serv Res*. 2019;19(1):974. [PubMed: 31852493]
262. Mueller KJ, Ortega ST, Parker K, Patil K, Askenazi A. Health status and access to care among rural minorities. *J Health Care Poor Underserved*. 1999;10(2):230–49. [PubMed: 10224828]
263. Probst JC, Moore CG, Glover SH, Samuels ME. Person and place: the compounding effects of race/ethnicity and rurality on health. *Am J Public Health*. 2004;94(10):1695–703. [PubMed: 15451735]
264. Hiza HA, Casavale KO, Guenther PM, Davis CA. Diet quality of Americans differs by age, sex, race/ethnicity, income, and education level. *J Acad Nutr Diet*. 2013;113(2):297–306. [PubMed: 23168270]
265. McCabe-Sellers BJ, Bowman S, Stuff JE, Champagne CM, Simpson PM, Bogle ML. Assessment of the diet quality of US adults in the Lower Mississippi Delta. *Am J Clin Nutr*. 2007;86(3):697–706. [PubMed: 17823435]

266. Wang Y, Chen X. How much of racial/ethnic disparities in dietary intakes, exercise, and weight status can be explained by nutrition- and health-related psychosocial factors and socioeconomic status among US adults? *J Am Diet Assoc.* 2011;111(12):1904–11. [PubMed: 22117667]
267. Netting MJ, Middleton PF, Makrides M. Does maternal diet during pregnancy and lactation affect outcomes in offspring? A systematic review of food-based approaches. *Nutrition.* 2014;30(11–12):1225–41. [PubMed: 25280403]
268. Maslova E, Hansen S, Strom M, Halldorsson TI, Olsen SF. Maternal intake of vitamins A, E and K in pregnancy and child allergic disease: a longitudinal study from the Danish National Birth Cohort. *Br J Nutr.* 2014;111(6):1096–108. [PubMed: 24229579]
269. Roduit C, Frei R, Depner M, Schaub B, Loss G, Genuneit J, et al. Increased food diversity in the first year of life is inversely associated with allergic diseases. *J Allergy Clin Immunol.* 2014;133(4):1056–64. [PubMed: 24508301]
270. U.S. Department of Agriculture Economic Research Service. Food Security and Nutrition Assistance 2020 [updated November 8, 2021. Available from: <https://www.ers.usda.gov/data-products/ag-and-food-statistics-charting-the-essentials/food-security-and-nutrition-assistance/>.
271. Bower KM, Thorpe RJ Jr., Rohde C, Gaskin DJ. The intersection of neighborhood racial segregation, poverty, and urbanicity and its impact on food store availability in the United States. *Prev Med.* 2014;58:33–9. [PubMed: 24161713]
272. Berkowitz SA, Seligman HK, Choudhry NK. Treat or eat: food insecurity, cost-related medication underuse, and unmet needs. *Am J Med.* 2014;127(4):303–10 e3. [PubMed: 24440543]
273. Kaur J, Lamb MM, Ogden CL. The Association between Food Insecurity and Obesity in Children-The National Health and Nutrition Examination Survey. *J Acad Nutr Diet.* 2015;115(5):751–8. [PubMed: 25737437]
274. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of childhood and adult obesity in the United States, 2011–2012. *JAMA.* 2014;311(8):806–14. [PubMed: 24570244]
275. Williams JD, Crockett D, Harrison RL, Thomas KD. The role of food culture and marketing activity in health disparities. *Prev Med.* 2012;55(5):382–6. [PubMed: 22227280]
276. Grier SA, Kumanyika SK. The context for choice: health implications of targeted food and beverage marketing to African Americans. *Am J Public Health.* 2008;98(9):1616–29. [PubMed: 18633097]
277. Beuther DA, Sutherland ER. Overweight, obesity, and incident asthma: a meta-analysis of prospective epidemiologic studies. *Am J Respir Crit Care Med.* 2007;175(7):661–6. [PubMed: 17234901]
278. Beckett WS, Jacobs DR Jr., Yu X, Iribarren C, Williams OD. Asthma is associated with weight gain in females but not males, independent of physical activity. *Am J Respir Crit Care Med.* 2001;164(11):2045–50. [PubMed: 11739133]
279. Kusunoki T, Morimoto T, Nishikomori R, Heike T, Ito M, Hosoi S, et al. Obesity and the prevalence of allergic diseases in schoolchildren. *Pediatr Allergy Immunol.* 2008;19(6):527–34. [PubMed: 18221473]
280. Dixon AE, Pratley RE, Forgione PM, Kaminsky DA, Whittaker-Leclair LA, Griffes LA, et al. Effects of obesity and bariatric surgery on airway hyperresponsiveness, asthma control, and inflammation. *J Allergy Clin Immunol.* 2011;128(3):508–15 e1–2. [PubMed: 21782230]
281. Scott HA, Gibson PG, Garg ML, Wood LG. Airway inflammation is augmented by obesity and fatty acids in asthma. *Eur Respir J.* 2011;38(3):594–602. [PubMed: 21310876]
282. Telenga ED, Tideman SW, Kerstjens HA, Hacken NH, Timens W, Postma DS, et al. Obesity in asthma: more neutrophilic inflammation as a possible explanation for a reduced treatment response. *Allergy.* 2012;67(8):1060–8. [PubMed: 22686834]
283. Silverberg JI, Silverberg NB, Lee-Wong M. Association between atopic dermatitis and obesity in adulthood. *Br J Dermatol.* 2012;166(3):498–504. [PubMed: 21999468]
284. Wolch J, Byrne J, Newell JP. Urban green space, public health, and environmental justice: The challenge of making cities ‘just green enough’. *Landscape and Urban Planning.* 2014;125:234–44.

285. Maller C, Townsend M, Pryor A, Brown P, St Leger L. Healthy nature healthy people: 'contact with nature' as an upstream health promotion intervention for populations. *Health Promot Int*. 2006;21(1):45–54. [PubMed: 16373379]
286. Kleinschroth F, Kowarik I. COVID-19 crisis demonstrates the urgent need for urban greenspaces. *Front Ecol Environ*. 2020;18(6):318–9. [PubMed: 32834788]
287. Wolch J, Jerrett M, Reynolds K, McConnell R, Chang R, Dahmann N, et al. Childhood obesity and proximity to urban parks and recreational resources: a longitudinal cohort study. *Health Place*. 2011;17(1):207–14. [PubMed: 21075670]
288. Woo J, Tang N, Suen E, Leung J, Wong M. Green space, psychological restoration, and telomere length. *Lancet*. 2009;373(9660):299–300.
289. Haahtela T, Holgate S, Pawankar R, Akdis CA, Benjaponpitak S, Carballo L, et al. The biodiversity hypothesis and allergic disease: world allergy organization position statement. *World Allergy Organ J*. 2013;6(1):3. [PubMed: 23663440]
290. Selway CA, Mills JG, Weinstein P, Skelly C, Yadav S, Lowe A, et al. Transfer of environmental microbes to the skin and respiratory tract of humans after urban green space exposure. *Environ Int*. 2020;145:106084. [PubMed: 32977191]
291. Paciencia I, Moreira A, Moreira C, Cavaleiro Rufo J, Sokhatska O, Rama T, et al. Neighbourhood green and blue spaces and allergic sensitization in children: A longitudinal study based on repeated measures from the Generation XXI cohort. *Sci Total Environ*. 2021;772:145394. [PubMed: 33561602]
292. Kim HJ, Min JY, Kim HJ, Min KB. Association between green areas and allergic disease in Korean adults: a cross-sectional study. *Ann Occup Environ Med*. 2020;32:e5. [PubMed: 32082587]
293. Sbihi H, Tamburic L, Koehoorn M, Brauer M. Greenness and Incident Childhood Asthma: A 10-Year Follow-up in a Population-based Birth Cohort. *Am J Respir Crit Care Med*. 2015;192(9):1131–3. [PubMed: 26517419]
294. Hartley K, Ryan P, Brokamp C, Gillespie GL. Effect of greenness on asthma in children: A systematic review. *Public Health Nurs*. 2020;37(3):453–60. [PubMed: 31899558]
295. Feng X, Astell-Burt T. Is Neighborhood Green Space Protective against Associations between Child Asthma, Neighborhood Traffic Volume and Perceived Lack of Area Safety? Multilevel Analysis of 4447 Australian Children. *Int J Environ Res Public Health*. 2017;14(5).
296. Fuertes E, Markevych I, Thomas R, Boyd A, Granell R, Mahmoud O, et al. Residential greenspace and lung function up to 24 years of age: The ALSPAC birth cohort. *Environ Int*. 2020;140:105749. [PubMed: 32380303]
297. Dai D. Racial/ethnic and socioeconomic disparities in urban green space accessibility: Where to intervene? *Landscape and Urban Planning*. 2011;102(4):234–44.
298. Estabrooks PA, Lee RE, Gyurcsik NC. Resources for physical activity participation: does availability and accessibility differ by neighborhood socioeconomic status? *Ann Behav Med*. 2003;25(2):100–4. [PubMed: 12704011]
299. Nesbitt L, Meitner M, Girling C, Sheppard S, Lu Y Who has access to urban vegetation? A spatial analysis of distributional green equity in 10 US cities. *Landscape and Urban Planning*. 2018;181:51–79.
300. Runkle JD, Matthews JL, Sparks L, McNicholas L, Sugg MM. Racial and ethnic disparities in pregnancy complications and the protective role of greenspace: A retrospective birth cohort study. *Sci Total Environ*. 2022;808:152145. [PubMed: 34871679]
301. Ruffin J. A renewed commitment to environmental justice in health disparities research. *Am J Public Health*. 2011;101 Suppl 1:S12–4. [PubMed: 22039043]
302. Bloom DE, Bowser DM. The population health and income nexus in the Mississippi River Delta Region and beyond. *J Health Hum Serv Adm*. 2008;31(1):105–23. [PubMed: 18575150]
303. U.S. Environmental Protection Agency. Environmental Justice [Available from: <https://www.epa.gov/environmentaljustice>].
304. Kearney GD, Johnson LC, Xu X, Balanay JA, Lamm KM, Allen DL. Eastern Carolina Asthma Prevention Program (ECAPP): An Environmental Intervention Study Among Rural

- and Underserved Children with Asthma in Eastern North Carolina. *Environ Health Insights*. 2014;8:27–37. [PubMed: 25057240]
305. U.S. Department of Housing and Urban Development, Office of Lead Hazard Control and Healthy Homes. Guide to Sustaining Effective Asthma Home Intervention Programs. 2018.
306. Crocker DD, Kinyota S, Dumitru GG, Ligon CB, Herman EJ, Ferdinands JM, et al. Effectiveness of home-based, multi-trigger, multicomponent interventions with an environmental focus for reducing asthma morbidity: a community guide systematic review. *Am J Prev Med*. 2011;41(2 Suppl 1):S5–32. [PubMed: 21767736]
307. Perdue W, Stone, LA, Gostin, LO. The Built Environment and Its Relationship to the Public's Health: The Legal Framework. *American Journal of Public Health*. 2003;93(9):1390–4. [PubMed: 12948949]
308. Scally C. Urban Wire: Health and Health Care [Internet]: Urban Institute. 2017. [cited 2022]. Available from: <https://www.urban.org/urban-wire/exposing-and-addressing-child-asthma-hud-assisted-renter-households>.
309. Phipatanakul W, Koutrakis P, Coull BA, Petty CR, Gaffin JM, Sheehan WJ, et al. Effect of School Integrated Pest Management or Classroom Air Filter Purifiers on Asthma Symptoms in Students With Active Asthma: A Randomized Clinical Trial. *JAMA*. 2021;326(9):839–50. [PubMed: 34547084]
310. Let's Move: America's Move to Raise a Healthier Generation of Kids [Available from: <https://letsmove.obamawhitehouse.archives.gov/become-lets-move-city-or-town>].




Figure 1.
 Conceptual Framework for Race/Ethnicity, Environmental Exposure Disparities, and Allergic Disease Burden

Table 1.

Summary of disparities in exposure to environmental hazards by race, ethnicity and income level

Environmental Hazards	Groups with Disproportionate Exposure
Exposure to violence and psychological stress	<ul style="list-style-type: none"> • Higher psychological stress and exposure to community violence among populations of color (240, 241) • Increased risk of aggravated assault and homicide among Black youth compared to White youth (242, 243) • Higher reported exposure to adverse childhood experiences among Black and Hispanic/Latinx children (242, 245, 246) • Black youth more likely to miss school due to safety concerns (242) • Experience of racial discrimination as a source of psychological stress linked to poor health (247–249)
Air pollution	<ul style="list-style-type: none"> • Poorest air quality in U.S. counties with high percentages of non-Hispanic Black residents (43) • African Americans and families living in poverty live closer to industrial emission sources (68) • Proportion of residents from racial/ethnic minority groups increases with decreasing distance from hazardous waste facilities in the U.S. (70) • Larger proportion of Non-Hispanic Black and Hispanic/Latinx children exposed to neighborhood traffic (97) and live in close proximity to major roadways (116) • Bus depots commonly located in low income neighborhoods (117) • Neighborhood disadvantage strengthens associations between pollution and respiratory disease (119–122) • Black and Hispanic/Latinx populations exposed to higher levels of volatile organic compounds compared to White populations (219) • Low educational attainment and lower income associated with higher rates of current smoking and maternal smoking in pregnancy (220–222) • Exposure to secondhand smoke higher in Black and high poverty populations (193, 223)
Substandard housing and Indoor allergens	<ul style="list-style-type: none"> • People of color and low income families more likely to live in poor housing compared to White and higher income families (162–164) • Racial/ethnic minority and low income families more likely to experience pests in the home (131, 166, 167) and are exposed to higher levels of indoor allergens in home and school environments (128, 160, 168–176) • High rates of sensitization to pest allergens in Black and Hispanic/Latinx populations (143, 168)
Access to healthcare	<ul style="list-style-type: none"> • Racial/ethnic minority populations less likely to have health insurance (250) • People of color attend fewer doctor visits, experience greater fragmentation of care and less continuity of care (250, 251) • Black children and children from non-English speaking families less likely to have a usual source of care (256) • Minority populations less likely to receive guidelines based asthma care (258–260) • Pharmacy deserts in predominantly minority and low income neighborhoods limit access to medications (257) • African American and Hispanic/Latinx patients reported having fewer choices in where they receive medical care (250) • Minority populations more likely to receive care from safety net hospitals (254) including emergency departments as opposed to office-based care, even after controlling for insurance coverage (255)
Diet and Food security	<ul style="list-style-type: none"> • Fewer supermarkets in predominantly Black neighborhoods (271) • Targeting of African Americans for marketing of high calorie, poorly nutritious foods (275, 276)
Green space access	<ul style="list-style-type: none"> • Less access to green space in low income and minority neighborhoods (297–299) • Smaller average size of parks in Black neighborhoods compared to White neighborhoods (297)