



Influenza (Flu) (</flu/index.htm>)

# Weekly U.S. Influenza Surveillance Report

Updated May 26, 2023



**Note:** CDC is also tracking the impact of other respiratory viruses, including COVID-19. Data comparing the impact of these viruses can be found in two new dashboards: RESP-NET (<https://www.cdc.gov/surveillance/resp-net/dashboard.html>) and NSSP's Emergency Department Visits for COVID-19, Influenza and Respiratory Syncytial Virus (<https://www.cdc.gov/ncird/surveillance/respiratory-illnesses/index.html>).

## Key Updates for Week 20, ending May 20, 2023

Seasonal influenza activity remains low nationally.

This is the last full FluView report of the 2022-2023 season. An abbreviated report will be published during the summer, and the first full report of the 2023-2024 season will be published on October 13, 2023.

## Viruses

### Clinical Lab

**1.1%** (Trend →)  
positive for influenza  
this week

[\(/flu/weekly/index.htm#ClinicalLaboratories\)](/flu/weekly/index.htm#ClinicalLaboratories)

### Public Health Lab

Influenza B viruses were the most  
frequently reported this week.

[\(/flu/weekly/index.htm#PublicHealthLaboratories\)](/flu/weekly/index.htm#PublicHealthLaboratories)

### Virus Characterization

Genetic and antigenic  
characterization and antiviral  
susceptibility are summarized in  
this report.

[\(/flu/weekly/index.htm#VirusCharacterization\)](/flu/weekly/index.htm#VirusCharacterization)

## Illness

### Outpatient Respiratory Illness

**2.0%** (Trend →)  
of visits to a health care provider this week were for respiratory illness  
(*below baseline*).

[\(/flu/weekly/index.htm#ILINet\)](/flu/weekly/index.htm#ILINet)

## Outpatient Respiratory Illness: Activity Map

This week 1 jurisdiction experienced moderate activity and 2 jurisdictions experienced high activity.

[\(/flu/weekly/index.htm#ORIAM\)](/flu/weekly/index.htm#ORIAM)

## Long-term Care Facilities

**0.3%** (Trend →)

of facilities reported  
≥ 1 influenza-positive test  
among residents this week.

[\(/flu/weekly/index.htm#LTCF\)](/flu/weekly/index.htm#LTCF)

## FluSurv-NET

**62.6 per 100,000**

cumulative hospitalization rate

[\(/flu/weekly/index.htm#FluSurvNet\)](/flu/weekly/index.htm#FluSurvNet)

## HHS Protect Hospitalizations

**939** (Trend →)

patients admitted to hospitals with influenza  
this week.

[\(/flu/weekly/index.htm#HHSProtect\)](/flu/weekly/index.htm#HHSProtect)

## NCHS Mortality

**6.5%** (Trend →)

of deaths attributed to pneumonia, influenza, or COVID-19 this week (**above threshold**).

[\(/flu/weekly/index.htm#NCHSMortality\)](/flu/weekly/index.htm#NCHSMortality)

## Pediatric Deaths

**2**

deaths were reported this week for a total of  
154 so far this season

[\(/flu/weekly/index.htm#PedMortality\)](/flu/weekly/index.htm#PedMortality)

*All data are preliminary and may change as more reports are received.*

*Directional arrows indicate changes between the current week and the previous week. Additional information on the arrows can be found at the bottom of this page.*

*A description of the CDC influenza surveillance system, including methodology and detailed descriptions of each data component is available on the [surveillance methods \(/flu/weekly/overview.htm\)](/flu/weekly/overview.htm) page.*

*Additional information on the current and previous influenza seasons for each surveillance component are available on [FluView Interactive \(/flu/weekly/fluviewinteractive.htm\)](/flu/weekly/fluviewinteractive.htm).*

### Key Points

- Seasonal influenza activity remains low nationally.
- This week is the last full version of FluView for the 2022-2023 influenza season. Starting with week 21, an abbreviated summer version of FluView will be published. The full version is expected to resume for week 40 of 2023, which is the start of the 2023-2024 influenza season.
- Nationally, outpatient respiratory illness is below baseline, and eight of 10 HHS regions are below their respective baselines.
- The number of flu hospital admissions remains low.
- During week 20, 36.1% of the 36 viruses reported by public health laboratories were influenza A and 63.9% were influenza B. All six influenza A viruses detected and subtyped during week 20 were influenza A(H1N1).
- Two influenza-associated pediatric death that occurred during the 2022-2023 season were reported this week, for a total of 154 pediatric flu deaths reported so far this season.
- CDC estimates that, so far this season, there have been at least 27 million illnesses, 300,000 hospitalizations, and 19,000 deaths from flu.

- The majority of influenza viruses tested are in the same genetic subclade as and antigenically similar to the influenza viruses included in this season's influenza vaccine.
- All viruses collected and evaluated this season have been susceptible to the influenza antivirals peramivir, zanamivir, and baloxavir, and all viruses except for one (> 99.9%) have been susceptible to the influenza antiviral oseltamivir.
- CDC continues to recommend that everyone ages 6 months and older get an annual flu vaccine as long as flu activity continues.

## U.S. Virologic Surveillance

(<https://www.cdc.gov/flu/weekly/overview.htm#LabSurveillance>)


Nationally, the percentage of specimens testing positive for influenza in clinical laboratories remained stable (change of <0.5 percentage points). For regional and state level data and age group distribution, please visit [FluView Interactive](https://gis.cdc.gov/grasp/fluview/fluportaldashboard.html) (<https://gis.cdc.gov/grasp/fluview/fluportaldashboard.html>). Viruses known to be associated with recent live attenuated influenza vaccine (LAIV) receipt or found upon further testing to be a vaccine virus are not included, as they are not circulating influenza viruses.

## Clinical Laboratories

The results of tests performed by clinical laboratories nationwide are summarized below. Data from clinical laboratories (the percentage of specimens tested that are positive for influenza) are used to monitor whether influenza activity is increasing or decreasing.

	Week 20	Data Cumulative since October 2, 2022 (Week 40)
No. of specimens tested	31,894	3,395,600
No. of positive specimens (%)	355 (1.1%)	352,572 (10.4%)
<i>Positive specimens by type</i>		
Influenza A	108 (30.4%)	345,544 (98.0%)
Influenza B	247 (69.6%)	7,028 (2.0%)

## Influenza Positive Tests Reported to CDC by U.S. Clinical Laboratories, National Summary, October 2, 2022 – May 20, 2023



<http://gis.cdc.gov/grasp/fluview/fluportaldashboard.html>

[View Chart Data \(/flu/weekly/weeklyarchives2022-2023/data/whoAllregt\\_cl20.html\)](/flu/weekly/weeklyarchives2022-2023/data/whoAllregt_cl20.html) | [View Full Screen \(/flu/weekly/WeeklyArchives2022-2023/WHONPHL20.html\)](/flu/weekly/WeeklyArchives2022-2023/WHONPHL20.html)


### Public Health Laboratories

The results of tests performed by public health laboratories nationwide are summarized below. Data from public health laboratories are used to monitor the proportion of circulating viruses that belong to each influenza subtype/lineage.

	Week 20	Data Cumulative since October 2, 2022 (Week 40)
No. of specimens tested	2,780	248,451
No. of positive specimens	36	29,364
<i>Positive specimens by type/subtype</i>		
Influenza A	13 (36.1%)	28,582 (97.3%)
(H1N1)pdm09	6 (100%)	6,545 (27.2%)
H3N2	0 (0%)	17,500 (72.8%)
H3N2v	0	1 (<0.1%)
Subtyping not performed	7	4,536

	Week 20	Data Cumulative since October 2, 2022 (Week 40)
Influenza B	23 (63.9%)	782 (2.7%)
Yamagata lineage	0	0
Victoria lineage	17 (100%)	593 (100%)
Lineage not performed	6	189

### Influenza Positive Tests Reported to CDC by U.S. Public Health Laboratories, National Summary, October 2, 2022 – May 20, 2023



(<http://gis.cdc.gov/grasp/fluview/fluportaldashboard.html>)

[View Chart Data \(/flu/weekly/weeklyarchives2022-2023/data/whoAllregt\\_phl20.html\)](/flu/weekly/weeklyarchives2022-2023/data/whoAllregt_phl20.html) | [View Full Screen \(/flu/weekly/weeklyarchives2022-2023/WhoPHL20.html\)](/flu/weekly/weeklyarchives2022-2023/WhoPHL20.html)

#### Additional virologic surveillance information for current and past seasons:

[Surveillance Methods \(/flu/weekly/overview.htm#LabSurveillance\)](/flu/weekly/overview.htm#LabSurveillance) | [FluView Interactive: National, Regional, and State Data \(http://gis.cdc.gov/grasp/fluview/fluportaldashboard.html\)](#) or [Age Data \(https://gis.cdc.gov/grasp/fluview/flu\\_by\\_age\\_virus.html\)](https://gis.cdc.gov/grasp/fluview/flu_by_age_virus.html)

## Influenza Virus Characterization

[\(/flu/weekly/overview.htm#VirusCharacterization\)](/flu/weekly/overview.htm#VirusCharacterization)

CDC performs [genetic \(https://www.cdc.gov/flu/about/professionals/genetic-characterization.htm\)](https://www.cdc.gov/flu/about/professionals/genetic-characterization.htm) and [antigenic \(https://www.cdc.gov/flu/about/professionals/antigenic.htm\)](https://www.cdc.gov/flu/about/professionals/antigenic.htm) characterization of U.S. viruses submitted from state and local public health laboratories according to the Right Size Roadmap submission guidance. These data are used to compare how similar the currently circulating influenza viruses are to the reference viruses representing viruses contained in the current influenza vaccines. The data are also used to monitor evolutionary changes that continually occur in influenza viruses circulating in humans.

CDC genetically characterized 2,912 influenza viruses collected since October 2, 2022. To reflect the genetic diversity of the HA genes in recent A(H1N1)pdm09 and A(H3N2) viruses, new subclades have been designated. The vast majority of A(H1N1)pdm09 viruses collected in the U.S. this season express HA genes belonging to 6B.1A.5a.2, but this clade has been split into the 2a and 2a.1 subclades. All A(H3N2) viruses collected thus far in the U.S. this season express HA genes belonging to clade 3C.2a1b.2a.2, and this clade has been split into multiple subclades. The new subclade designations improve the ability to track the evolution and co-circulation of multiple groups of influenza viruses.

Virus Subtype or Lineage	Genetic Characterization				
	Total No. of Subtype/Lineage Tested	HA Clade	Number (% of subtype/lineage tested)	HA Subclade	Number (% of subtype/lineage tested)
<b>A/H1</b>	1,062				
		6B.1A.5a	1,062 (100%)	1	6 (0.6%)
				2a	166 (15.6%)
				2a.1	890 (83.8%)
<b>A/H3</b>	1,634				
		3C.2a1b.2a	1,634 (100%)	2a	25 (1.5%)
				2a.1	185 (11.3%)
				2a.1b	137 (8.4%)
				2a.3	45 (2.8%)
				2a.3a	3 (0.2%)
				2a.3a.1	71 (4.3%)
				2a.3b	11 (0.7%)
				2b	1,157 (70.8%)
				2c	0 (0%)
<b>B/Victoria</b>	216				
		V1A	216 (100%)	3	5 (2.3%)
				3a.2	211 (97.7%)

Virus Subtype or Lineage	Genetic Characterization				
	Total No. of Subtype/Lineage Tested	HA Clade	Number (% of subtype/lineage tested)	HA Subclade	Number (% of subtype/lineage tested)
B/Yamagata	0				
		Y3	0	Y3	0 (0%)

CDC antigenically characterizes (<https://www.cdc.gov/flu/about/professionals/antigenic.htm>) influenza viruses by hemagglutination inhibition (HI) (<http://www.cdc.gov/flu/professionals/laboratory/antigenic.htm>) (H1N1pdm09, B/Victoria, and B/Yamagata viruses) or neutralization-based HINT (<https://www.cdc.gov/flu/spotlights/2018-2019/new-lab-method-test-flu.html>) (H3N2 viruses) using antisera that ferrets make after being infected with reference viruses representing the 2022-2023 Northern Hemisphere recommended cell or recombinant-based vaccine viruses. Antigenic differences between viruses are determined by comparing how well the antibodies made against the vaccine reference viruses recognize the circulating viruses that have been grown in cell culture. Ferret antisera are useful because antibodies raised against a particular virus can often recognize small changes in the surface proteins of other viruses. In HI assays, viruses with similar antigenic properties have antibody titer differences of less than or equal to 4-fold when compared to the reference (vaccine) virus. In HINT, viruses with similar antigenic properties have antibody neutralization titer differences of less than 8-fold. Viruses selected for antigenic characterization are a subset representing the genetic changes in the surface proteins seen in genetically characterized viruses.

#### Influenza A Viruses

- **A (H1N1)pdm09:** Three hundred and fourteen A(H1N1)pdm09 viruses were antigenically characterized by HI, and 310 (99%) were well-recognized (reacting at titers that were within 4-fold of the homologous virus titer) by ferret antisera to cell-grown A/Wisconsin/588/2019-like reference viruses representing the A(H1N1)pdm09 component for the cell- and recombinant-based influenza vaccines.
- **A (H3N2):** Two hundred and twenty-five A(H3N2) viruses were antigenically characterized by HINT, and 212 (94%) were well-recognized (reacting at titers that were within 8-fold of the homologous virus titer) by ferret antisera to cell-grown A/Darwin/6/2021-like reference viruses representing the A(H3N2) component for the cell- and recombinant-based influenza vaccines.

#### Influenza B Viruses

- **B/Victoria:** Fifty-eight influenza B/Victoria-lineage virus were antigenically characterized by HI, and all were well-recognized (reacting at titers that were within 4-fold of the homologous virus titer) by ferret antisera to cell-grown B/Austria/1359417/2021-like reference viruses representing the B/Victoria component for the cell- and recombinant-based influenza vaccines.
- **B/Yamagata:** No influenza B/Yamagata-lineage viruses were available for antigenic characterization.

#### Assessment of Virus Susceptibility to Antiviral Medications

CDC assesses susceptibility of influenza viruses to antiviral medications including the neuraminidase inhibitors (oseltamivir, zanamivir, and peramivir) and the PA endonuclease inhibitor baloxavir using next generation sequence analysis supplemented by laboratory assays. Information about antiviral susceptibility test methods can be found at [U.S. Influenza Surveillance: Purpose and Methods | CDC](https://www.cdc.gov/flu/weekly/overview.htm) (<https://www.cdc.gov/flu/weekly/overview.htm>).

Viruses collected in the U.S. since October 2, 2022, were tested for antiviral susceptibility as follows:

Antiviral Medication	Total Viruses	A/H1	A/H3	B/Victoria	B/Yamagata

Antiviral Medication			Total Viruses	A/H1	A/H3	B/Victoria	B/Yamagata
Neuraminidase Inhibitors	Oseltamivir	Viruses Tested	2,904	1,059	1,630	215	0
		Reduced Inhibition	1 (<0.1%)	1 (0.1%)	0 (0%)	0 (0%)	0 (0%)
		Highly Reduced Inhibition	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
	Peramivir	Viruses Tested	2,904	1,059	1,630	215	0
		Reduced Inhibition	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
		Highly Reduced Inhibition	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
	Zanamivir	Viruses Tested	2,904	1,059	1,630	215	0
		Reduced Inhibition	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
		Highly Reduced Inhibition	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
PA Cap-Dependent Endonuclease Inhibitor	Baloxavir	Viruses Tested	2,820	1,016	1,593	211	0
		Reduced Susceptibility	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)

One A(H1N1)pdm09 virus had NA-S247G amino acid substitution and showed reduced inhibition by oseltamivir.


## Outpatient Respiratory Illness Surveillance

(<https://www.cdc.gov/flu/weekly/overview.htm#ILINet>)

The U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet) monitors outpatient visits for respiratory illness referred to as influenza-like illness [ILI (fever plus cough or sore throat)], not laboratory-confirmed influenza, and will therefore capture respiratory illness visits due to infection with any pathogen that can present with similar symptoms, including influenza, SARS-CoV-2, and RSV. Due to the COVID-19 pandemic, health care-seeking behaviors have changed, and people may be accessing the health care system in alternative settings not captured as a part of ILINet or at a different point in their illness than they might have before the pandemic. Therefore, it is important to evaluate syndromic surveillance data, including that from ILINet, in the context of other sources of surveillance data to obtain a complete and accurate picture of influenza, SARS-CoV-2, and other respiratory virus activity. CDC is tracking COVID-19 activity in a weekly publication called [COVID Data Tracker Weekly Review](https://www.cdc.gov/coronavirus/2019-ncov/covid-data/covidview/index.html) (<https://www.cdc.gov/coronavirus/2019-ncov/covid-data/covidview/index.html>). Information about other respiratory virus activity can be found on [CDC's National Respiratory and Enteric Virus Surveillance System \(NREVSS\) website](https://www.cdc.gov/surveillance/nrevss/index.html) (<https://www.cdc.gov/surveillance/nrevss/index.html>).

## Outpatient Respiratory Illness Visits

Nationwide during week 20, 2.0% of patient visits reported through ILINet were due to respiratory illness that included fever plus a cough or sore throat, also referred to as ILI. This has remained stable (change of  $\leq 0.1$  percentage points) compared to week 19 and is below the national baseline of 2.5%. Eight of 10 HHS regions are below their respective baselines; Region 2 is above its baseline, and Region 9 is at its baseline. Multiple respiratory viruses are co-circulating, and the relative contribution of influenza virus infection to ILI varies by location.



<http://gis.cdc.gov/grasp/fluview/fluportaldashboard.html>

\* Effective October 3, 2021 (week 40), the ILI definition (fever plus cough or sore throat) no longer includes “without a known cause other than influenza.”


[View Chart Data \(current season only\) \(/flu/weekly/weeklyarchives2022-2023/data/senAllregt20.html\)](/flu/weekly/weeklyarchives2022-2023/data/senAllregt20.html) | [View Full Screen \(/flu/weekly/weeklyarchives2022-2023/ILI20.html\)](/flu/weekly/weeklyarchives2022-2023/ILI20.html)

## Outpatient Respiratory Illness Visits by Age Group

More than 70% of ILINet participants provide both the number of patient visits for respiratory illness and the total number of patient visits for the week broken out by age group. Data from this subset of providers are used to calculate the percentages of patient visits for respiratory illness by age group.

The percentage of visits for respiratory illness reported in ILINet remained stable (change of  $\leq 0.1$  percentage point) for all age groups (0-4 years, 5-24 years, 25-49 years, 50-64 years, and 65+ years) in week 20 compared to week 19.

**Percentage of Outpatient Visits for Respiratory Illness by Age Group  
Reported by the U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet),  
Weekly National Summary, October 2, 2022-May 20, 2023**



(<http://gis.cdc.gov/grasp/fluview/fluportaldashboard.html>)

[View Chart Data \(/flu/weekly/weeklyarchives2022-2023/data/iliage20.html\)](/flu/weekly/weeklyarchives2022-2023/data/iliage20.html) | [View Full Screen \(/flu/weekly/weeklyarchives2022-2023/ILIAge20.html\)](/flu/weekly/weeklyarchives2022-2023/ILIAge20.html)

## Outpatient Respiratory Illness Activity Map

Data collected in ILINet are used to produce a measure of ILI activity\*

([https://www.cdc.gov/flu/weekly/overview.htm#anchor\\_1633697504110](https://www.cdc.gov/flu/weekly/overview.htm#anchor_1633697504110)) by state/jurisdiction and Core Based Statistical Areas (CBSA).

Activity Level	Number of Jurisdictions		Number of CBSAs	
	Week 20 (Week ending May 20, 2023)	Week 19 (Week ending May 13, 2023)	Week 20 (Week ending May 20, 2023)	Week 19 (Week ending May 13, 2023)
Very High	0	0	1	1
High	2	4	6	6
Moderate	1	0	12	15
Low	3	3	58	58
Minimal	49	48	585	606
Insufficient Data	0	0	267	243

# A Weekly Influenza Surveillance Report Prepared by the Influenza Division

## Outpatient Respiratory Illness Activity Map Determined by Data Reported to ILINet

This system monitors visits for respiratory illness that includes fever plus a cough or sore throat, also referred to as ILI, not laboratory confirmed influenza and may capture patient visits due to other respiratory pathogens that cause similar symptoms.



prev Play Pause next

40 50 1 10 20  
weeks

State CBSA

2022-23 Influenza Season Week 20 ending May 20, 2023

ILI Activity Level



Season: 2022-23 ▲

Download Image

Download Data

(<https://www.cdc.gov/flu/weekly/fluereport.xml>)View Full Screen (<http://gis.cdc.gov/grasp/fluview/main.html>)

\*Data collected in ILINet may disproportionately represent certain populations within a jurisdiction or CBSA, and therefore, may not accurately depict the full picture of influenza activity for the entire jurisdiction or CBSA. Differences in the data presented here by CDC and independently by some health departments likely represent differing levels of data completeness with data presented by the health department likely being the more complete.

### Additional information about medically attended visits for ILI for current and past seasons:


[Surveillance Methods \(/flu/weekly/overview.htm#ILINet\)](#) | [FluView Interactive: National, Regional, and State Data \(http://gis.cdc.gov/grasp/fluview/fluportaldashboard.html\)](#) or [ILI Activity Map \(https://gis.cdc.gov/grasp/fluview/main.html\)](#)

## Long-term Care Facility (LTCF) Surveillance

(<https://www.cdc.gov/flu/weekly/overview.htm#LongTermCare>)

LTCFs (e.g., nursing homes/skilled nursing, long-term care for the developmentally disabled, and assisted living facilities) from all 50 states and U.S. territories report data on influenza virus infections among residents through the [National Healthcare Safety Network \(NHSN\) Long-term Care Facility Component \(https://www.cdc.gov/nhsn/lc/index.html\)](#). During 20, 43 (0.3%) of 14,074 facilities reported at least one influenza positive test among their residents. This percentage remains stable compared to week 19.

## Percent of Long-term Care Facilities (LTCF) with at Least One Confirmed Influenza Positive Test among Residents, Reported to CDC National Healthcare Safety Network (NHSN), National Summary, October 3, 2022 – May 21, 2023



</flu/weekly/weeklyarchives2022-2023/LTCF20.html> View Chart Data  </flu/weekly/weeklyarchives2022-2023/data/LTCFData20.csv> | View Full Screen </flu/weekly/weeklyarchives2022-2023/LTCF20.html>

### Additional information about long-term care facility surveillance:

[Surveillance Methods \(/flu/weekly/overview.htm#LongTermCare\)](/flu/weekly/overview.htm#LongTermCare) | [Additional Data !\[\]\(65669ef2a9341eca7c5ba6092e766555\_img.jpg\) \(https://data.cms.gov/covid-19/covid-19-nursing-home-data\)](https://data.cms.gov/covid-19/covid-19-nursing-home-data)

## Hospitalization Surveillance

<http://www.cdc.gov/flu/weekly/overview.htm#HospitalizationSurv>

### FluSurv-NET

The Influenza Hospitalization Surveillance Network (FluSurv-NET) conducts population-based surveillance for laboratory-confirmed influenza-related hospitalizations in select counties in 13 states and represents approximately 9% of the U.S. population. FluSurv-NET hospitalization data are preliminary. As data are received each week, prior case counts and rates are updated accordingly.

A total of 18,306 laboratory-confirmed influenza-associated hospitalizations were reported by FluSurv-NET sites between October 1, 2022, and April 30, 2023. The weekly hospitalization rate observed in week 17 the last week of FluSurv-NET enrollment for the 2023-2024 season, was 0.2 per 100,000 population. The weekly rate observed during week 48 (week ending December 3, 2022) is the third highest peak weekly rate observed during all seasons going back to 2010-2011; this follows the 2017-2018 season, which peaked during week 1 (week ending January 6, 2018) and the 2014-2015 season, which peaked during week 52 (week ending December 27, 2014).


The overall cumulative hospitalization rate was 62.6 per 100,000 population. This cumulative hospitalization rate is similar to the hospitalization rates for 4 seasons (2014-2015, 2016-2017, 2018-2019, and 2019-2020 seasons) and lower than the hospitalization rate for the 2017-2018 season, going back to 2010-2011.

When examining rates by age, the highest rate of hospitalization per 100,000 population was among adults aged 65 and older (186.8). Among adults aged 65 and older, rates were highest among adults aged 85 and older (102.1). Among persons aged <65 years, hospitalization rates per 100,000 population were highest among children aged 0-4 years (80.4), followed by adults aged 50-64 years (68.0). When examining rates by race and ethnicity, the highest rate of hospitalization per 100,000 population was among non-Hispanic Black persons (88.5), followed by non-Hispanic American Indian or Alaska Native persons (82.7), Hispanic/Latino persons (55.6), non-Hispanic White persons (54.1), and non-Hispanic Asian/Pacific Islander persons (27.7).

Among 18,306 hospitalizations, 17,487 (95.5%) were associated with influenza A virus, 649 (3.5%) with influenza B virus, 30 (0.2%) with influenza A virus and influenza B virus co-infection, and 140 (0.8%) with influenza virus for which the type was not determined. Among 4,564 hospitalizations with influenza A subtype information, 3,412 (74.8%) were A(H3N2), and 1,152 (25.3%) were A(H1N1)pdm09. Based on preliminary data, of the 5,689 laboratory-confirmed influenza-associated hospitalizations with more complete data, 3.7% (95% CI: 3.1%-4.4%) also tested positive for SARS-CoV-2.

Among 3,926 hospitalized adults with information on underlying medical conditions, 97.0% had at least one reported underlying medical condition, the most commonly reported were hypertension, cardiovascular disease, metabolic disorder, and obesity. Among 1,422 hospitalized women of childbearing age (15-49 years) with information on pregnancy status, 36.8% were pregnant. Among 1,385 hospitalized children with information on underlying medical conditions, 66.4% had at least one reported underlying medical condition; the most commonly reported was asthma, followed by neurologic disease, and obesity.


While patients admitted after April 30, 2023, will not be included, data on patients admitted through April 30, 2023, will continue to be updated as additional information is received.



<https://gis.cdc.gov/grasp/fluview/FluHospRates.html>

[View Full Screen \(/flu/weekly/weeklyarchives2022-2023/EIPRates20.html\)](/flu/weekly/weeklyarchives2022-2023/EIPRates20.html)

Cumulative Rate of Laboratory-Confirmed Influenza Hospitalizations among cases of all ages, 2014-15 to 2022-23, MMWR Week 17



\*\*In this figure, weekly rates for all seasons reflect end-of-season rates. As hospitalization data are received each week, prior case counts and rates are updated accordingly.

(<https://gis.cdc.gov/grasp/fluview/FluHospChars.html>)

[View Full Screen \(/flu/weekly/weeklyarchives2022-2023/EIPConditions20.html\)](/flu/weekly/weeklyarchives2022-2023/EIPConditions20.html)


### Additional FluSurv-NET hospitalization surveillance information for current and past seasons and additional age groups:

Surveillance Methods (<https://www.cdc.gov/flu/weekly/overview.htm#FluSurvNet>) | FluView Interactive: Rates by Age, Sex, and Race/Ethnicity (<http://gis.cdc.gov/GRASP/Fluview/FluHospRates.html>) or Data on Patient Characteristics (<http://gis.cdc.gov/grasp/fluview/FluHospChars.html>) | RESP-NET Interactive (<https://www.cdc.gov/surveillance/resp-net/dashboard.html>)

## HHS Protect Hospitalization Surveillance

Hospitals report to HHS Protect the number of patients admitted with laboratory-confirmed influenza. During week 20, 939 patients with laboratory-confirmed influenza were admitted to a hospital. The number of patients admitted to a hospital with laboratory-confirmed influenza remained stable compared to week 19.

## New Influenza Hospital Admissions Reported to HHS Protect, National Summary, October 2, 2022 – May 20, 2023



[View Chart Data](/flu/weekly/weeklyarchives2022-2023/Protect20.html) [View Full Screen](/flu/weekly/weeklyarchives2022-2023/data/ProtectData20.csv)

### Additional HHS Protect hospitalization surveillance information:

[Surveillance Methods](https://www.cdc.gov/flu/weekly/overview.htm#HHSProtect) | [Additional Data](https://healthdata.gov/Hospital/COVID-19-Reported-Patient-Impact-and-Hospital-Capa/anag-cw7u)

## Mortality Surveillance


<https://www.cdc.gov/flu/weekly/overview.htm#MortalitySurveillance>

### National Center for Health Statistics (NCHS) Mortality Surveillance

Based on NCHS mortality surveillance data available on May 25, 2023, 6.5% of the deaths that occurred during the week ending May 20, 2023 (week 20), were due to pneumonia, influenza, and/or COVID-19 (PIC). This percentage remained stable ( $\leq 0.3$  percentage point change) compared to week 19 and is above the epidemic threshold of 6.4% for this week. Among the 1,483 PIC deaths reported for this week, 304 had COVID-19 listed as an underlying or contributing cause of death on the death certificate, and 15 listed influenza. The data presented are preliminary and may change as more data are received and processed.

# Pneumonia, Influenza, and COVID-19 Mortality from the National Center for Health Statistics Mortality Surveillance System

Data as of May 25, 2023



(<http://gis.cdc.gov/GRASP/Fluview/mortality.html>)

[View Chart Data](#) (x) (</flu/weekly/weeklyarchives2022-2023/data/NCHSData20.csv>) | [View Full Screen](#) (</flu/weekly/weeklyarchives2022-2023/NCHS20.html>)

**Additional pneumonia, influenza and COVID-19 mortality surveillance information for current and past seasons:**

[Surveillance Methods](https://www.cdc.gov/flu/weekly/overview.htm#NCHSMortality) (<https://www.cdc.gov/flu/weekly/overview.htm#NCHSMortality>) | [FluView Interactive](#)


(<https://gis.cdc.gov/grasp/fluview/mortality.html>)

## Influenza-Associated Pediatric Mortality

Two influenza-associated pediatric deaths occurring during the 2022-2023 season were reported to CDC during week 20. One death was associated with an influenza A(H1N1) virus and the other death was associated with an influenza B virus with no lineage determined. Both deaths occurred during week 19 (the week ending May 13, 2023).

A total of 154 influenza-associated pediatric deaths occurring during the 2022-2023 season have been reported to CDC.

# Influenza-Associated Pediatric Deaths by Week of Death, 2019-2020 season to 2022-2023 season



<http://gis.cdc.gov/GRASP/Fluview/PedFluDeath.html>

[View Full Screen \(/flu/weekly/weeklyarchives2022-2023/PedFlu20.html\)](/flu/weekly/weeklyarchives2022-2023/PedFlu20.html)

## Additional pediatric mortality surveillance information for current and past seasons:

[Surveillance Methods \(https://www.cdc.gov/flu/weekly/overview.htm#PediatricMortality\)](https://www.cdc.gov/flu/weekly/overview.htm#PediatricMortality) | [FluView Interactive](#)

<https://gis.cdc.gov/GRASP/Fluview/PedFluDeath.html>

## Trend Indicators

Increasing: ↑

Decreasing: ↓

Stable: →

## Indicators Status by System

**Clinical Labs:** Up or down arrows indicate a change of greater than or equal to 0.5 percentage points in the percent of specimens positive for influenza compared to the previous week.

**Outpatient Respiratory Illness (ILINet):** Up or down arrows indicate a change of greater than 0.1 percentage points in the percent of visits due to respiratory illness (ILI) compared to the previous week.

**Long-term Care Facilities:** Up or down arrows indicate change of greater than or equal to 5% of the percent of facilities reporting at least one influenza positive test among their residents compared to the previous week.

**HHS Protect Hospitalizations:** Up or down arrows indicate change of greater than or equal to 5% of the number of patients admitted with laboratory-confirmed influenza compared to the previous week.

**NCHS Mortality:** Up or down arrows indicate change of greater than 0.3 percentage points of the percent of deaths due to PIC compared to the previous week.

# Additional National and International Influenza Surveillance Information

**FluView Interactive:** FluView includes enhanced web-based interactive applications that can provide dynamic visuals of the influenza data collected and analyzed by CDC. These [FluView Interactive applications](http://www.cdc.gov/flu/weekly/fluviewinteractive.htm) (<http://www.cdc.gov/flu/weekly/fluviewinteractive.htm>) allow people to create customized, visual interpretations of influenza data, as well as make comparisons across flu seasons, regions, age groups and a variety of other demographics.

**National Institute for Occupational Safety and Health:** Monthly surveillance data on the prevalence of health-related workplace absenteeism among full-time workers in the United States are [available from NIOSH](https://www.cdc.gov/niosh/topics/absences/default.html) (<https://www.cdc.gov/niosh/topics/absences/default.html>).

**U.S. State and local influenza surveillance:** Select a jurisdiction below to access the latest local influenza information.

<a href="http://adph.org/influenza/">Alabama</a> ( <a href="http://adph.org/influenza/">http://adph.org/influenza/</a> )	<a href="http://dhss.alaska.gov/dph/Epi/id/Pages/influenza/fluintfo.aspx">Alaska</a> ( <a href="http://dhss.alaska.gov/dph/Epi/id/Pages/influenza/fluintfo.aspx">http://dhss.alaska.gov/dph/Epi/id/Pages/influenza/fluintfo.aspx</a> )
<a href="https://www.colorado.gov/pacific/cdphe/influenza">Colorado</a> ( <a href="https://www.colorado.gov/pacific/cdphe/influenza">https://www.colorado.gov/pacific/cdphe/influenza</a> )	<a href="https://portal.ct.gov/DPH/Epidemiology-and-Emerging-Infections/Influenza-Surveillance-and-Statistics">Connecticut</a> ( <a href="https://portal.ct.gov/DPH/Epidemiology-and-Emerging-Infections/Influenza-Surveillance-and-Statistics">https://portal.ct.gov/DPH/Epidemiology-and-Emerging-Infections/Influenza-Surveillance-and-Statistics</a> )
<a href="https://dph.georgia.gov/flu-activity-georgia">Georgia</a> ( <a href="https://dph.georgia.gov/flu-activity-georgia">https://dph.georgia.gov/flu-activity-georgia</a> )	<a href="http://health.hawaii.gov/docd/resources/reports/influenza-reports/">Hawaii</a> ( <a href="http://health.hawaii.gov/docd/resources/reports/influenza-reports/">http://health.hawaii.gov/docd/resources/reports/influenza-reports/</a> )
<a href="https://idph.iowa.gov/influenza/reports">Iowa</a> ( <a href="https://idph.iowa.gov/influenza/reports">https://idph.iowa.gov/influenza/reports</a> )	<a href="http://www.kdheks.gov/flu/surveillance.htm">Kansas</a> ( <a href="http://www.kdheks.gov/flu/surveillance.htm">http://www.kdheks.gov/flu/surveillance.htm</a> )
<a href="https://phpa.health.maryland.gov/influenza/fluwatch/">Maryland</a> ( <a href="https://phpa.health.maryland.gov/influenza/fluwatch/">https://phpa.health.maryland.gov/influenza/fluwatch/</a> )	<a href="https://www.mass.gov/influenza">Massachusetts</a> ( <a href="https://www.mass.gov/influenza">https://www.mass.gov/influenza</a> )
<a href="http://health.mo.gov/living/healthcondiseases/communicable/influenza/reports.php">Missouri</a> ( <a href="http://health.mo.gov/living/healthcondiseases/communicable/influenza/reports.php">http://health.mo.gov/living/healthcondiseases/communicable/influenza/reports.php</a> )	<a href="https://dphhs.mt.gov/publichealth/cdepi/diseases/influenza/in">Montana</a> ( <a href="https://dphhs.mt.gov/publichealth/cdepi/diseases/influenza/in">https://dphhs.mt.gov/publichealth/cdepi/diseases/influenza/in</a> )
<a href="http://www.nj.gov/health/cd/topics/flu.shtml">New Jersey</a> ( <a href="http://www.nj.gov/health/cd/topics/flu.shtml">http://www.nj.gov/health/cd/topics/flu.shtml</a> )	<a href="https://nmhealth.org/about/erd/ideb/isp/">New Mexico</a> ( <a href="https://nmhealth.org/about/erd/ideb/isp/">https://nmhealth.org/about/erd/ideb/isp/</a> )
<a href="http://www.flu.ohio.gov">Ohio</a> ( <a href="http://www.flu.ohio.gov">http://www.flu.ohio.gov</a> )	<a href="https://oklahoma.gov/health/health-education/acute-disease-service/disease-information/influenza-home-page.html">Oklahoma</a> ( <a href="https://oklahoma.gov/health/health-education/acute-disease-service/disease-information/influenza-home-page.html">https://oklahoma.gov/health/health-education/acute-disease-service/disease-information/influenza-home-page.html</a> )
<a href="http://www.scdhec.gov/Health/DiseasesandConditions/InfectiousDiseases/Flu/FluData/">South Carolina</a> ( <a href="http://www.scdhec.gov/Health/DiseasesandConditions/InfectiousDiseases/Flu/FluData/">http://www.scdhec.gov/Health/DiseasesandConditions/InfectiousDiseases/Flu/FluData/</a> )	<a href="https://doh.sd.gov/diseases/infectious/flu/surveillance.aspx">South Dakota</a> ( <a href="https://doh.sd.gov/diseases/infectious/flu/surveillance.aspx">https://doh.sd.gov/diseases/infectious/flu/surveillance.aspx</a> )
<a href="http://www.healthvermont.gov/immunizations-infectious-disease/influenza/flu-activity-and-surveillance">Vermont</a> ( <a href="http://www.healthvermont.gov/immunizations-infectious-disease/influenza/flu-activity-and-surveillance">http://www.healthvermont.gov/immunizations-infectious-disease/influenza/flu-activity-and-surveillance</a> )	<a href="http://www.vdh.virginia.gov/epidemiology/influenza-in-virginia/influenza-surveillance/">Virginia</a> ( <a href="http://www.vdh.virginia.gov/epidemiology/influenza-in-virginia/influenza-surveillance/">http://www.vdh.virginia.gov/epidemiology/influenza-in-virginia/influenza-surveillance/</a> )
<a href="https://health.wyo.gov/publichealth/infectious-disease-epidemiology-unit/disease/influenza/">Wyoming</a> ( <a href="https://health.wyo.gov/publichealth/infectious-disease-epidemiology-unit/disease/influenza/">https://health.wyo.gov/publichealth/infectious-disease-epidemiology-unit/disease/influenza/</a> )	<a href="http://www1.nyc.gov/site/doh/providers/health-topics/flu-alerts.page">New York City</a> ( <a href="http://www1.nyc.gov/site/doh/providers/health-topics/flu-alerts.page">http://www1.nyc.gov/site/doh/providers/health-topics/flu-alerts.page</a> )

## World Health Organization:

Additional influenza surveillance information from participating WHO member nations is available through [FluNet](https://www.who.int/tools/flunet) (<https://www.who.int/tools/flunet>) and the [Global Epidemiology Reports](https://www.who.int/teams/global-influenza-programme/surveillance-and-monitoring/influenza-surveillance-outputs). (<https://www.who.int/teams/global-influenza-programme/surveillance-and-monitoring/influenza-surveillance-outputs>)

## WHO Collaborating Centers for Influenza:

[Australia](http://www.influenzacentre.org/Surveillance_Samples_Received.html) ([http://www.influenzacentre.org/Surveillance\\_Samples\\_Received.html](http://www.influenzacentre.org/Surveillance_Samples_Received.html)), [China](http://www.chinaivdc.cn/cnic/) (<http://www.chinaivdc.cn/cnic/>), [Japan](http://idsc.nih.go.jp/index.html) (<http://idsc.nih.go.jp/index.html>), the [United Kingdom](https://www.crick.ac.uk/research/worldwide-influenza-centre) (<https://www.crick.ac.uk/research/worldwide-influenza-centre>), and the [United States](http://www.cdc.gov/flu/) (CDC in Atlanta, Georgia) (<http://www.cdc.gov/flu/>)

**Europe:**

The most up-to-date influenza information from Europe is available from [WHO/Europe and the European Centre for Disease Prevention and Control](http://www.flunewseurope.org/)  (<http://www.flunewseurope.org/>).

**Public Health Agency of Canada:**

The most up-to-date influenza information from Canada is available in [Canada's weekly FluWatch report](http://www.phac-aspc.gc.ca/fluwatch/)  (<http://www.phac-aspc.gc.ca/fluwatch/>).

**Public Health England:**

The most up-to-date influenza information from the United Kingdom is available from [Public Health England](http://www.hpa.org.uk/Topics/InfectiousDiseases/InfectionsAZ/SeasonalInfluenza/)  (<http://www.hpa.org.uk/Topics/InfectiousDiseases/InfectionsAZ/SeasonalInfluenza/>).

**Any links provided to non-Federal organizations are provided solely as a service to our users. These links do not constitute an endorsement of these organizations or their programs by CDC or the Federal Government, and none should be inferred. CDC is not responsible for the content of the individual organization web pages found at these links.**

A description of the CDC influenza surveillance system, including methodology and detailed descriptions of each data component is available on the [surveillance methods](http://www.cdc.gov/flu/weekly/overview.htm) (<http://www.cdc.gov/flu/weekly/overview.htm>) page.

Last Reviewed: May 26, 2023, 11:00 AM