
Leishmania donovani Transmission Cycle Associated with Human Infection, *Phlebotomus alexandri* Sand Flies, and Hare Blood Meals, Israel¹

Liora Studentsky, Laor Orshan, Fouad Akad, Irina Ben Avi, Debora Diaz, Shirly Elbaz,
Orly Sagi, Gal Zagron, Lea Valinsky, Maya Davidovich-Cohen, Gad Baneth

Cutaneous leishmaniasis caused by *Leishmania major* or *L. tropica* and visceral leishmaniasis caused by *L. infantum* have been reported in Israel. We collected *Phlebotomus* spp. sand flies in the Negev desert of southern Israel to identify circulating *Leishmania* spp. Of 22,636 trapped sand flies, 80% were *P. alexandri*. We sequenced *Leishmania*-specific internal transcribed spacer 1 fragments and *K26* genes. Of 5,019 *Phlebotomus* female sand flies, 2.5% were *Leishmania* DNA-positive; 92% of infections were *L. donovani*. Phylogenetic analyses showed separate clustering of *L. donovani* and *L. infantum*. *P. alexandri* flies positive for *L. donovani* harbored blood meals from European hares. *Leishmania* DNA isolated from a patient with cutaneous leishmaniasis who lived in the survey area was identical to *L. donovani* from *P. alexandri* flies. We report circulation of *L. donovani*, a cause of visceral leishmaniasis, in southern Israel. Prompt diagnosis and *Leishmania* spp. identification are critical to prevent leishmaniasis progression.

Zoonotic leishmaniasis is endemic to Israel. *Leishmania tropica*, *L. major*, and *L. infantum* infect humans in different areas of Israel and circulate through distinct zoonotic transmission cycles (1). Cutaneous

leishmaniasis (CL) is caused by *L. major*, which is transmitted by *Phlebotomus papatasi* sand flies, and *L. tropica*, which is transmitted by *P. sergenti* and *P. arabicus* sand flies. Canine leishmaniasis and human visceral leishmaniasis (VL) are caused by *L. infantum* in Israel, and the putative vectors are *P. perfiliewi*, *P. syriacus*, and *P. tobbi* sand flies (1–9). Reservoirs for *L. major* are sand rats (*Psammomys obesus*), gerbils (*Gerbillus dasyurus*), jirds (*Meriones crassus* and *M. tristrami*), and possibly also voles (10–13), whereas rock hyraxes (*Procapra capensis*) are considered the animal reservoir for *L. tropica* in Israel (14). Domestic dogs (*Canis lupus familiaris*), jackals (*C. aureus*), foxes (*Vulpes vulpes*), and wolves (*C. lupus*) are recognized reservoir hosts for *L. infantum* (15).

A substantial increase in CL incidence has been recorded since 2002, and endemic transmission has occurred in areas of Israel where it was previously unknown (6,16,17). Although not life-threatening, CL is a considerable public health problem in Israel; CL is diagnosed in hundreds of new patients annually. During 2001–2018, CL incidence rates increased 7-fold, from 0.4 to 2.9/100,000 population; a peak was observed in 2012, when the mean annual incidence increased to 4.4/100,000 population (18,19). Our study combines results from sand fly surveys, *Phlebotomus* spp. blood meal analysis, and human patient clinical data from the mountainous area of central Negev in southern Israel during the summer months of 2018–2020. We found a fourth leishmaniasis transmission cycle associated with human illness.

Author affiliations: Ministry of Health Central Laboratories for Public Health, Jerusalem, Israel (L. Studentsky, L. Orshan, I. Ben Avi, D. Diaz, S. Elbaz, M. Davidovich-Cohen); The Hebrew University of Jerusalem, Rehovot, Israel (L. Studentsky, G. Baneth); Ministry of Health National Laboratory for Public Health, Tel Aviv, Israel (F. Akad); Soroka University Medical Center, Beer-Sheba, Israel (O. Sagi); Israeli Ministry of Environmental Protection, Jerusalem (G. Zagron); Maccabi Healthcare Services, Tel Aviv (L. Valinsky)

DOI: <https://doi.org/10.3201/eid2905.221657>

¹Data from this study were presented at the Israeli Society for Parasitology, Protozoology and Tropical Diseases Annual Meeting; March 21, 2022; Kfar Hamaccabiah, Ramat Gan, Israel.

Materials and Methods

Study Area and Sand Fly Trapping

We conducted our study in the mountainous desert area of central Negev in southern Israel (Figure 1). In this region, elevations range from 50 to 1,037 m above sea level, large differences occur between peak daytime and nighttime temperatures, and annual average precipitation is 30–150 mm (20,21). We collected sand flies outdoors in August 2018, September 2019, and August 2020 by using modified traps from the




Figure 1. Locations of *Phlebotomus* spp. sand fly collection sites within the central Negev region of Israel in study of *Leishmania donovani* transmission cycle associated with human infection, *Phlebotomus alexandri* sand flies, and hare blood meals. We collected sand flies outdoors in August 2018, September 2019, and August 2020 by using modified traps from the US Centers for Disease Control and Prevention. The traps operated without light and were powered by 2 AA (1.2V) rechargeable batteries and baited with ≈ 1 kg dry ice. Traps were placed in an updraft vertical position overnight; openings were ≈ 10 cm above the ground, and collection cups hung above the motor and fan. Different colored circles indicate sites where specific *Leishmania* spp. infections were identified in trapped *Phlebotomus* sand flies. Empty circles indicate sites where sand flies were negative for *Leishmania* spp. Inset shows location of the survey area in Israel (red box).

US Centers for Disease Control and Prevention. The traps operated without light and were powered by 2 AA (1.2V) rechargeable batteries and baited with ≈ 1 kg dry ice. We placed traps in an updraft vertical position overnight; openings were ≈ 10 cm above the ground, and collection cups hung above the motor and fan (22,23).

Identification and Sample Preparation

We transferred live sand fly catches to the laboratory, which we then chilled and processed. We counted dead sand flies and sorted by sex, identifying all male flies at the species level by using specific morphologic keys for genitalia (24,25). We kept all engorged females and ≤ 10 –15 unfed females from each trap individually. If the number of female sand flies in the trap was > 15 , we pooled those flies with others in groups of 20 specimens each. We noted the blood meal size and freshness for each engorged female (26). We stored all female fly specimens in collection microtubes at -80°C until DNA extraction.

Molecular Analysis by Real-Time PCR, HRM Assay, and Sequencing

We extracted total DNA from sand fly samples by using the QIAasympphony DSP DNA Mini Kit and QIAasympphony SP robot (QIAGEN, <https://www.qiagen.com>). We homogenized the samples for 5 min in 50 μL lysis buffer and stainless steel beads by using a TissueLyser II instrument (QIAGEN). The lysis buffer contained DNase- and proteinase-free RNaseA (ThermoFisher Scientific, <https://www.thermo-fisher.com>), proteinase K, and ATL tissue lysis buffer (QIAGEN). After homogenization, we added 200 μL lysis buffer to each samples and incubated at 56°C for 2 h. We performed centrifugation and transferred the samples directly to the robot. We extracted DNA in accordance with the manufacturer's instructions and eluted the DNA in 100 μL of elution buffer.

We performed all real-time PCR reactions by using a Roche LightCycler 96 (Roche, <https://www.roche.com>) and AccuMelt HRM SuperMix (Quantabio, <https://www.quantabio.com>). We analyzed all female sand flies for *Leishmania* spp. infection and single and engorged female flies to determine *Phlebotomus* sand fly species and blood meal source. We performed high-resolution melting (HRM) assays at the final step of each real-time PCR. We performed amplicon dissociation analysis by capturing fluorescence signals in $0.1^{\circ}\text{C}/\text{s}$ increments and holding for 60 s in each range of the melting curve (60°C – 85°C for sand fly species and blood meal detection assays or $\leq 95^{\circ}\text{C}$ for *Leishmania* PCR). Sanger sequencing was

performed at the Center for Genomic Technologies at Hebrew University of Jerusalem.

We screened all female sand flies for *Leishmania* DNA and identified parasite species by amplifying an internal transcribed spacer (ITS) 1 rRNA fragment with ITS1–219 PCR primers (Appendix Table 1, <https://wwwnc.cdc.gov/EID/article/29/5/22-1657-App1.pdf>) and by using the HRM assay (27). For PCR controls, we extracted DNA from parasite promastigote cultures of international reference strains: *L. major* (MHOM/PS/1967/JerichoII), *L. tropica* (MHOM/IL/1990/P283), *L. infantum* (MHOM/SD/62/2S), *L. donovani* (MHOM/SD/1962/1S-CLD2), and *L. aethiopicus* (MHOM/ET/1972/L102). High purity water for molecular biology (Bio-Lab, <http://www.biolab-chemicals.com>) was used as a negative control.

We included DNA isolated from skin lesions from 4 patients who had leishmaniasis diagnosed at the parasitology laboratory at Soroka Medical Center, Beer-Sheba, Israel; leishmaniasis was caused by *L. donovani*/*L. infantum* complex in those patients (Table 1). Leishmaniasis was diagnosed at the hospital by using multiplex real-time PCR with 5 probes for the ITS region of *Leishmania* sp. (28,29). We analyzed the samples further at the Ministry of Health by using real-time PCR–HRM amplification of the ITS1 fragment and ITS region and then sequencing.

We amplified the entire 1,020-bp ribosomal ITS region from *Leishmania*-positive field, clinical, and control samples by using PCR primers LITSR and LITSV (Appendix Table 1). If the entire ITS region was not successfully amplified with LITSR and LITSV primers, we used an internal pair of primers, L5.8S and L5.8SR, to amplify ITS1 and ITS2 separately (30). We performed ITS1 amplicon sequencing for 12 of the positive samples, and the entire ITS region was sequenced from 6 unfed and 3 engorged females, 3 pooled *Phlebotomus* spp. samples, all 4 human samples, and 4 *Leishmania*-positive controls. We amplified the repeat region of the *L. donovani* and *L. infantum* HASPB (known as K26) gene for additional

separation of *L. donovani* complex-positive samples by using primers K26F and K26R (31).

To identify *Phlebotomus* spp., we amplified a 368–393-bp fragment of the cytochrome b gene by using a universal primer set designed for this study (cytb-F and cytb-R; Appendix Table 1). The specificity of the designed primers was tested against DNA sequences from hematophagous arthropods, including sand flies, mosquitoes, and ticks. Male sand flies identified at the species level by using morphologic characteristics were used as positive controls and molecular biology grade water was used as a negative control. We analyzed all individual samples, and 1 third of samples from each melting curve pattern were sequenced.

We identified blood meal sources in *Phlebotomus* sand fly specimens by amplifying a 500-bp segment of host 12S and 16S mitochondrial rRNA genes by using modified vertebrate universal primers N12–16F and N12–16R (32). We included negative (water) and positive (100 ng of human DNA) controls in each PCR. We sequenced 50 samples that represented all HRM curve patterns and all female sand flies containing blood meals that had a melting curve of a rare host (<5 samples).

We used DNA from *Leishmania* reference strains, male sand fly specimens identified by morphologic characteristics, and human blood as templates for real-time PCR and HRM curve standardization. Each species produced a unique melting curve that was easily distinguishable from other species and consistent with observed nucleotide differences (Appendix Figures 1–3). We compared normalized HRM curves of field samples with the positive control included in each PCR, which enabled species determination (27). We validated species identification by sequencing 1 third of the samples; complete matches were observed for speciation by HRM curve analysis and DNA sequencing.

We aligned and corrected nucleotide sequences by using BioNumerics version 8.0 software (Applied Maths, <https://www.applied-maths.com>) and

Table 1. Human clinical samples from Soroka Medical Center in study of *Leishmania donovani* transmission cycle associated with human infection, *Phlebotomus alexandri* sand flies, and hare blood meals, Israel*

Patient no.	Diagnosis	Age, y/sex	Clinical description	Residence	Infecting <i>Leishmania</i> sp.†
1	VL	47/M	Splenomegaly	Northern Israel	<i>L. infantum</i> (100% identity with <i>L. infantum</i> , GenBank accession no. KU680954)
2	VL	4/F	Splenomegaly, hepatomegaly	Hebron	<i>L. infantum</i> (99.19% identity with <i>L. infantum</i> , GenBank accession no. MN503527)
3	CL	69/F	Skin ulcer	Negev	<i>L. infantum</i> (99.87% identity with <i>L. infantum</i> , GenBank accession no. KU680954)
4	CL	51/M	Skin ulcer	Arava	<i>L. donovani</i> (99.75% identity with <i>L. donovani</i> , GenBank accession no. LC459330)

*CL, cutaneous leishmaniasis; VL, visceral leishmaniasis.

†*Leishmania* spp. were identified by PCR high resolution melting curves and Sanger sequencing.




Figure 2. Phylogenetic analysis of *Leishmania* internal transcribed spacer 1 rRNA fragments in study of *Leishmania donovani* transmission cycle associated with human infection, *Phlebotomus alexandri* sand flies, and hare blood meals, Israel. *Leishmania*-specific internal transcribed spacer 1 rRNA fragments (201 bp) were amplified by PCR from *P. alexandri* sand flies, pooled female *Phlebotomus* spp. flies, and patient samples and then sequenced. Tree was constructed by using the maximum-likelihood method and Tamura 3-parameter model, estimated by using the Akaike information criterion (33). Dendrogram includes sequences from *L. donovani* and *L. infantum* isolated from sand flies and clinical samples in this study compared with *Leishmania* spp. reference controls and GenBank sequences from Israel and other countries. Tree shows substantial separate clustering of *L. infantum* (bootstrap 94%) and *L. donovani* (bootstrap 89%) sequences. Empty circles are *Leishmania* international reference strains, black triangles are the 10 sequences from our study deposited in GenBank, and black circles are additional *L. infantum*-positive sand flies samples from Israel. Available GenBank sequences for *L. major*, *L. tropica*, *L. infantum*, and *L. donovani* from Israel and other countries are also included. GenBank accession numbers, *Leishmania* spp., isolate source, and country are indicated. Only bootstrap values >70% are shown. Not to scale.

compared sequences against the GenBank database by using BLASTN (<http://blast.ncbi.nlm.nih.gov>). We identified *Leishmania* spp., blood meal sources, and sand fly species on the basis of >98% identity with sequences obtained during the BLAST search.

We submitted sequences of the ITS1 fragments and entire ITS and K26 regions obtained in this study to GenBank (Appendix Table 2).

We constructed phylogenetic trees on the basis of marker gene sequences in this study and relevant




Figure 3. Phylogenetic analysis of entire *Leishmania* internal transcribed spacer region in study of *Leishmania donovani* transmission cycle associated with human infection, *Phlebotomus alexandri* sand flies, and hare blood meals, Israel. *Leishmania*-specific internal transcribed spacer region (988 bp) was amplified by PCR from *P. alexandri* sand flies, pooled female *Phlebotomus* spp. flies, and patient samples and then sequenced. Tree was constructed by using the maximum-likelihood method and Tamura 3-parameter model of all relevant *Leishmania* spp. and *Trypanosoma cruzi* as an outgroup. Sand fly and clinical samples from this study (black triangles), *L. infantum* isolates from Israel (black circles), *Leishmania* international reference strains (empty circles), and available GenBank *Leishmania* sequences are shown. GenBank accession numbers, isolate source, and country of origin are shown for each sequence. Only bootstrap values >70% are shown next to branches. Not to scale.




Figure 4. Phylogenetic analysis of *Leishmania* K26 gene in study of *Leishmania donovani* transmission cycle associated with human infection, *Phlebotomus alexandri* sand flies, and hare blood meals, Israel. *Leishmania*-specific K26 gene fragment (348 bp) was amplified by PCR from *P. alexandri* flies, pooled female *Phlebotomus* spp. flies, and patient samples and then sequenced. Tree was constructed by using the maximum-likelihood method and Hasegawa-Kishino-Yano model. K26 phylogenetic analysis shows separation between *L. infantum* and *L. donovani*. Sand fly and clinical samples from this study (black triangles), *L. infantum* isolates from Israel (black circles), *Leishmania* international reference strains (empty circles), and available GenBank *Leishmania* sequences are shown. GenBank accession number, isolate source, and country of origin are shown for each sequence. Only bootstrap values >70% are shown next to branches. Not to scale.

sequences of other *Leishmania* spp. deposited in GenBank. We used MEGA X software (33) to infer phylogenetic trees after nucleotide sequence alignment was performed by using ClustalW software (<http://www.clustal.org>) and maximum-likelihood and neighbor-joining algorithms. We used 1,000 bootstrap replicates to determine percentages of replicate trees. We constructed a phylogenetic tree composed of 45 analyzed partial sequences of the ITS1 locus, including sequences of *Leishmania* spp. from Israel and other countries deposited in GenBank and *Trypanosoma*

cruzi as an outgroup (Figure 2). We constructed a second tree that included 30 nearly complete ITS sequences of all relevant *Leishmania* spp. and *T. cruzi* as an outgroup (Figure 3) and an additional phylogenetic tree that included 20 K26 gene sequences of *L. donovani* complex-positive samples (Figure 4).

Results

We collected 22,636 *Phlebotomus* spp. sand fly specimens (15,720 female and 6,916 male; sex ratio 2.3) during 7 trapping nights by using 118 traps placed

Table 2. Number of female and male *Phlebotomus* spp. collected during 2018–2020 in central Negev in study of *Leishmania donovani* transmission cycle associated with human infection, *Phlebotomus alexandri* sand flies, and hare blood meals, Israel

Phlebotomus spp.	2018		2019		2020		Total no. (%)
	F	M	F	M	F	M	
<i>P. alexandri</i>	129	1,452	70	126	431	4,039	6,247 (80.0)
<i>P. kazeruni</i>	38	837	22	11	78	142	1,128 (14.5)
<i>P. sergenti</i>	5	109	26	9	34	56	239 (3.1)
<i>P. papatasi</i>	25	97	8	8	8	27	173 (2.2)
<i>P. syriacus</i>	0	0	3	0	17	3	23 (0.3)
Not identified*	8,344	0	265	0	6,217	0	14,826
Total	8,541	2,495	394	154	6,785	4,267	22,636

*Sand flies were pooled before molecular testing in batches of 20 specimens each.

at 94 sites. After identifying all male and 894 (6%) female flies, we found the catches consisted of 5 species. The most abundant sand fly species were *P. alexandri* (80%), *P. kazeruni* (14.4%), *P. sergenti* (3.1%), *P. papatasi* (2.2%), and *P. syriacus* (0.3%) (Table 2).

Among the 4,140 unfed female sand flies tested in 210 pools, we found 41 pools were positive for *Leishmania* spp. In addition, 6/688 single female flies and 4/206 engorged female flies were positive for *Leishmania* spp. Of the 51 *Leishmania*-positive samples, the HRM curves of 47 (36 pools, 6 single females, and 3 engorged female flies) were similar to the HRM curve of the *L. donovani* control (Table 3; Appendix Figure 1). The HRM curves for 2 pooled fly samples from 2018 were identical to the HRM curve of the *L. tropica* control. One pooled fly sample and 1 engorged female fly collected in 2020 had an HRM curve identical to the *L. major* control. The ITS1-PCR sequences of 20 samples (11 pools and all 9 single and engorged females) that had HRM curves similar to the *L. donovani* control HRM curve were also 100% identical to the *L. donovani* control sequence.

Leishmaniasis was diagnosed in 4 human patients (Table 1). The ITS1 HRM curve and sequence from patient 4 with CL were similar to the *L. donovani* control and 47 *L. donovani*-positive *Phlebotomus* spp. samples. The ITS1 HRM and sequences from patients

1 and 2 with VL and patient 3 with CL were similar to the *L. infantum* control. Alignment of ITS1 sequences from *L. infantum* and *L. donovani* controls, 3 representative sand fly samples showing HRM identical to *L. donovani*, and the 4 patient samples showed clustering into 2 distinct groups. The first group comprised the *L. donovani* control, 3 sand fly samples, and patient 4. The second group comprised the *L. infantum* control and samples from patients 1, 2, and 3. The difference between the groups was at position 71–74 in ITS1; the *L. donovani* group had a 4-nt (ATAT) insertion that was missing in the *L. infantum*-positive samples. A comparison of ITS1 sequences with those in GenBank showed 100% query coverage and 99.65%–100% identity with GenBank sequences for *L. infantum* and *L. donovani* from various countries (data not shown).

We aligned DNA sequences from the entire ITS region obtained from *Leishmania*-positive *P. alexandri* samples from our study and the *L. infantum* and *L. donovani* controls (Appendix Figure 4). We found 2 additional regions containing polymorphic sites that distinguished between *L. infantum* and *L. donovani*: a 2-nt (GG) deletion at position 724–725 and 1-nt (G) insertion at position 817 in the *L. donovani* sequence.

We constructed a phylogenetic tree of ITS1 rRNA fragments of *Leishmania* sequences obtained from *P. alexandri* flies, pooled female *Phlebotomus* spp. flies, and

Table 3. Number of *Leishmania* spp. detected in sand fly samples by PCR during 2018–2020 in study of *Leishmania donovani* transmission cycle associated with human infection, *Phlebotomus alexandri* sand flies, and hare blood meals, Israel*

Year	Total no. tested	<i>L. donovani</i>	<i>L. tropica</i>	<i>L. major</i>
2018				
Females (no. pools)	2,938 (148)	24	2	0
Single unfed females	108	0	0	0
Single engorged females	89	0	0	0
2019				
Females (no. pools)	262 (15)	4	0	0
Single unfed females	121	3	0	0
Single engorged females	8	0	0	0
2020				
Females (no. pools)	940 (47)	10	0	1
Single unfed females	459	3	0	0
Single engorged females	109	3	0	1
Total no.	5,019	47	2	2

*All engorged females and ≤10–15 unfed females from each trap were maintained and analyzed individually. If the number of female sand flies in the trap was >15, they were pooled in batches of 20 specimens each.

Table 4. Number of female *Phlebotomus* spp. sand flies collected in the central Negev region engorged with different blood meals in study of *Leishmania donovani* transmission cycle associated with human infection, *Phlebotomus alexandri* sand flies, and hare blood meals, Israel

Blood meal source	<i>Phlebotomus</i> spp.				Total no.
	<i>P. alexandri</i>	<i>P. kazeruni</i>	<i>P. papatasi</i>	<i>P. sergenti</i>	
<i>Lepus europaeus</i> (hare)	107	8	5	6	126
<i>Equus hemionus</i> (onager)	31	1	0	1	33
<i>Gazella dorcas</i> (gazelle)	10	0	4	2	16
<i>Canis lupus familiaris</i> (dog)	1	0	3	0	4
<i>Homo sapiens</i> (human)	1	0	0	0	1
<i>Psammomys obesus</i> (fat sand rat)	0	0	1	0	1
<i>Vulpes vulpes</i> (fox)	1	0	0	0	1
Total no.	151	9	13	9	182

patient samples from this study. We compared those sequences with *Leishmania* spp. controls and GenBank sequences from Israel and other countries. The tree showed substantial separate clustering of *L. infantum* (bootstrap 94%) and *L. donovani* (bootstrap 89%) sequences (Figure 2). Phylogenetic analysis of the entire ITS region showed separate clustering of *L. infantum* (bootstrap 84%) and *L. donovani* (bootstrap 95%) sequences (Figure 3). K26 phylogenetic analysis also showed separation between *L. infantum* and *L. donovani* (Figure 4).

We identified the blood meal source for 182/206 (88%) engorged female sand flies that represented the 4 most abundant sand fly species within the study area. We observed 7 types of HRM curves. We compared blood meal sequences with GenBank sequences and determined similarities between HRM curves. We identified European brown hare (*Lepus europaeus*) blood in 126 (69.2%) flies, onager (*Equus hemionus*) blood in 33 (18.3%) flies, gazelle (*Gazella dorcas*) blood in 16 (8.8%) flies, and domestic dog (*C. lupus familiaris*) blood in 4 (2.2%) flies; 1 female sand fly each contained blood from either a fat sand rat (*Psammomys obesus*), fox (*V. vulpes*), or human (Table 4). Hare blood was the dominant blood meal found in all 4 *Phlebotomus* spp. flies: *P. papatasi*, 38%; *P. sergenti*, 67%; *P. alexandri*, 71%; and *P. kazeruni*, 89%. The 12S–16S hare blood meal sequences were 99.8% similar to *L. europaeus* hares and only 95.3% similar to *L. capensis* hares.

Of the 47 *Phlebotomus* spp. sand fly samples with HRM curve patterns and sequences similar to the *L. donovani* control, 9 were single *P. alexandri* female sand flies, 3 of which were engorged with hare blood. Of the 2 *Phlebotomus* spp. samples positive for *L. major*, 1 was in a single engorged *P. papatasi* female sand fly that had an unsuccessful blood meal identification. The 2 identified *L. tropica* samples were from pooled female sand flies.

Discussion

We found a fourth transmission cycle of leishmaniasis in the central Negev region of southern Israel.

On the basis of molecular analysis of the ITS region and K26 gene and phylogenetic analysis, we concluded that the parasite found in patient 4, who lives in the survey area, and in *P. alexandri* sand flies was *L. donovani* sensu stricto. We found that *L. infantum* was the cause of illness in the other 3 patients with leishmaniasis. A case report describing patient 4 was published in 2016; the authors concluded that the infecting parasite was likely *L. infantum* because of prevailing knowledge of endemic *Leishmania* transmission in Israel (29). An earlier study reported another patient from the Arava region of central Negev, close to where patient 4 lives, who had symptoms of both CL and VL (34). The cause of infection was identified as *L. donovani*; the authors noted that this infection was unusual because *L. donovani* was not known to circulate in Israel. The earlier study substantiates our findings of *L. donovani* in both sand flies and another human patient within the same geographic area. *L. infantum* was identified as the causative agent in the other 3 patients in our study and was also described in canines in Israel (1).

The high abundance of *P. alexandri* sand flies within the study area and the association with *L. donovani* infections suggest that the *P. alexandri* sand fly is the putative vector of *L. donovani* in Israel. *P. alexandri* flies have been associated with *L. donovani* sensu lato transmission in other parts of the Old World. Natural infection by *L. donovani* was found in field-collected *P. alexandri* sand fly specimens in China, and inoculation of hamsters with those parasites caused VL (35). Another study reported the susceptibility of *P. alexandri* to artificial infection with *L. donovani* isolated from human patients in China (36).

We found blood meals from European brown hares in ≈70% of engorged female *Phlebotomus* sand flies in our study. The high feeding rates on hares, presence of *L. donovani* in female *P. alexandri* sand flies engorged with hare blood and illnesses reported in humans infected with *L. donovani* suggest a zoonotic *L. donovani* transmission cycle in Israel. Those data

suggest that the hare could be a potential reservoir and *P. alexandri* flies could be the putative vector for *L. donovani*. The role of hares as a reservoir host for *L. donovani* requires further investigation; however, a related hare species, *Lepus granatensis*, was reported as a potential sylvatic reservoir for *L. infantum* in a leishmaniasis outbreak in Madrid, Spain (37,38). Furthermore, studies in Greece and Italy detected *L. donovani* complex infection in *L. europaeus* hares (39,40), providing support for hares as a potential reservoir for *L. donovani* in Israel. Dogs were identified as reservoirs for *L. donovani* in India, Sudan, and Ethiopia, and different rodent species have been identified as possible reservoirs of *Leishmania* spp. from the *L. donovani* complex (41–48). However, no *L. donovani* infections in canines and rodents have been reported in Israel; infections in sand fly blood meals found in our study do not implicate those hosts in the local life cycle of *L. donovani*.

In conclusion, we found circulation of *L. donovani* in the Negev region of southern Israel that was associated with cutaneous lesions in humans. We determined that *P. alexandri* was the putative sand fly vector and that hares were the main reservoir host of *L. donovani*. We found 2 distinct *Leishmania* spp. in the *L. donovani* complex in Israel. Previously, the few reported human cases of CL resulting from *L. donovani* infections were attributed to either *L. infantum* or non-autochthonous infections. Analysis of patient samples in our study indicates that, in addition to *L. major* and *L. tropica* (the known agents causing CL), *L. donovani* is also a cause of autochthonous CL in Israel. Our results suggest that CL in Israel can be caused by *L. donovani*, a primary cause of VL. Therefore, prompt diagnosis, identification of the *Leishmania* sp., and treatment with drugs intended for visceral leishmaniasis, such as pentavalent antimonials or liposomal amphotericin B (49), are critical to prevent disease progression and death among patients with leishmaniasis.

Acknowledgments

We thank Simha Shilo, Anna Sarner, and Arik Levy for assisting with sand fly sorting; Aviv Rashti, Anat Berkowitz, Karin Cohen, and Alina Bazarski for assisting with human sample analysis; Itay Naveh and Dan Ish-Shalom for supporting and participating in the fieldwork; Heather Schnur for editorial remarks; Yaarit Nachum-Biala for sharing professional knowledge of phylogenetic tree construction; and Guy Nizry for helping with map creation.

This study was supported by internal funding from the Ministry of Health, Jerusalem, Israel.

About the Author

Ms. Studentsky is a PhD student at the Koret School of Veterinary Medicine, The Hebrew University of Jerusalem, Israel, and works at the Medical Entomology Laboratory, Ministry of Health, Jerusalem. Her primary research interests focus on leishmaniasis, sand fly surveillance, and vector competence.

References

- Jaffe CL, Baneth G, Abdeen ZA, Schlein Y, Warburg A. Leishmaniasis in Israel and the Palestinian Authority. *Trends Parasitol.* 2004;20:328–32. <https://doi.org/10.1016/j.pt.2004.05.001>
- Schlein Y, Warburg A, Schnur LF, Gunders AE. Leishmaniasis in the Jordan Valley II. Sandflies and transmission in the central endemic area. *Trans R Soc Trop Med Hyg.* 1982;76:582–6. [https://doi.org/10.1016/0035-9203\(82\)90215-2](https://doi.org/10.1016/0035-9203(82)90215-2)
- Schlein Y, Warburg A, Schnur LF, Le Blancq SM, Gunders AE. Leishmaniasis in Israel: reservoir hosts, sandfly vectors and leishmanial strains in the Negev, Central Arava and along the Dead Sea. *Trans R Soc Trop Med Hyg.* 1984;78:480–4. [https://doi.org/10.1016/0035-9203\(84\)90067-1](https://doi.org/10.1016/0035-9203(84)90067-1)
- Anis E, Leventhal A, Elkana Y, Wilamowski A, Pener H. Cutaneous leishmaniasis in Israel in the era of changing environment. *Public Health Rev.* 2001;29:37–47.
- Jacobson RL, Eisenberger CL, Svobodova M, Baneth G, Szttern J, Carvalho J, et al. Outbreak of cutaneous leishmaniasis in northern Israel. *J Infect Dis.* 2003;188:1065–73. <https://doi.org/10.1086/378204>
- Schnur LF, Nasereddin A, Eisenberger CL, Jaffe CL, El Fari M, Azmi K, et al. Multifarious characterization of *Leishmania tropica* from a Judean desert focus, exposing intraspecific diversity and incriminating *Phlebotomus sergenti* as its vector. *Am J Trop Med Hyg.* 2004;70:364–72. <https://doi.org/10.4269/ajtmh.2004.70.364>
- Svobodova M, Votypka J, Peckova J, Dvorak V, Nasereddin A, Baneth G, et al. Distinct transmission cycles of *Leishmania tropica* in 2 adjacent foci, northern Israel. *Emerg Infect Dis.* 2006;12:1860–8. <https://doi.org/10.3201/eid1212.060497>
- Jacobson RL. Leishmaniasis in an era of conflict in the Middle East. *Vector Borne Zoonotic Dis.* 2011;11:247–58. <https://doi.org/10.1089/vbz.2010.0068>
- Ready PD. Biology of phlebotomine sand flies as vectors of disease agents. *Annu Rev Entomol.* 2013;58:227–50. <https://doi.org/10.1146/annurev-ento-120811-153557>
- Gunders AE, Foner A, Montilio B. Identification of *Leishmania* species isolated from rodents in Israel. *Nature.* 1968;219:85–6. <https://doi.org/10.1038/219085a0>
- Gunders AE, Lidror R, Montilo B, Amitai P. Isolation of *Leishmania* sp. from *Psammodmys obesus* in Judea. *Trans R Soc Trop Med Hyg.* 1968;62:465. [https://doi.org/10.1016/0035-9203\(68\)90129-6](https://doi.org/10.1016/0035-9203(68)90129-6)
- Wasserberg G, Abramsky Z, Anders G, El-Fari M, Schoenian G, Schnur L, et al. The ecology of cutaneous leishmaniasis in Nizzana, Israel: infection patterns in the reservoir host, and epidemiological implications. *Int J Parasitol.* 2002;32:133–43. [https://doi.org/10.1016/S0020-7519\(01\)00326-5](https://doi.org/10.1016/S0020-7519(01)00326-5)
- Faiman R, Abbasi I, Jaffe C, Motro Y, Nasereddin A, Schnur LF, et al. A newly emerged cutaneous leishmaniasis focus in northern Israel and two new reservoir hosts of

- Leishmania major*. PLoS Negl Trop Dis. 2013;7:e2058. <https://doi.org/10.1371/journal.pntd.0002058>
14. Talmi-Frank D, Jaffe CL, Nasereddin A, Warburg A, King R, Svobodova M, et al. *Leishmania tropica* in rock hyraxes (*Procapra capensis*) in a focus of human cutaneous leishmaniasis. Am J Trop Med Hyg. 2010;82:814–8. <https://doi.org/10.4269/ajtmh.2010.09-0513>
 15. Ya'ari A, Jaffe CL, Garty BZ. Visceral leishmaniasis in Israel, 1960–2000. Isr Med Assoc J. 2004;6:205–8.
 16. Singer SR, Abramson N, Shoob H, Zaken O, Zentner G, Stein-Zamir C. Ecoepidemiology of cutaneous leishmaniasis outbreak, Israel. Emerg Infect Dis. 2008;14:1424–6. <https://doi.org/10.3201/eid1409.071100>
 17. Azmi K, Krayter L, Nasereddin A, Ereqat S, Schnur LF, Al-Jawabreh A, et al. Increased prevalence of human cutaneous leishmaniasis in Israel and the Palestinian Authority caused by the recent emergence of a population of genetically similar strains of *Leishmania tropica*. Infect Genet Evol. 2017;50:102–9. <https://doi.org/10.1016/j.meegid.2016.07.035>
 18. Gandacu D, Glazer Y, Anis E, Karakis I, Warshavsky B, Slater P, et al. Resurgence of cutaneous leishmaniasis in Israel, 2001–2012. Emerg Infect Dis. 2014;20:1605–11. <https://doi.org/10.3201/eid2010.140182>
 19. Israel Ministry of Health. Annual report of central laboratories, 2019 (Hebrew) [cited 2021 Jan 5]. https://www.health.gov.il/PublicationsFiles/LAB_JER2019.pdf
 20. Nezer O, Bar-David S, Gueta T, Carmel Y. High-resolution species-distribution model based on systematic sampling and indirect observations. Biodivers Conserv. 2017;26:421–37. <https://doi.org/10.1007/s10531-016-1251-2>
 21. Stern E, Gardus Y, Meir A, Krakover S, Tzoar H. Atlas of the Negev. Jerusalem (Israel): Keter Publishing House; 1986.
 22. Orshan L, Szekely D, Khalfa Z, Bitton S. Distribution and seasonality of *Phlebotomus* sand flies in cutaneous leishmaniasis foci, Judean Desert, Israel. J Med Entomol. 2010;47:319–28. <https://doi.org/10.1093/jmedent/47.3.319>
 23. Orshan L, Elbaz S, Ben-Ari Y, Akad F, Afik O, Ben-Avi I, et al. Distribution and dispersal of *Phlebotomus papatasi* (Diptera: Psychodidae) in a zoonotic cutaneous leishmaniasis focus, the northern Negev, Israel. PLoS Negl Trop Dis. 2016;10:e0004819. <https://doi.org/10.1371/journal.pntd.0004819>
 24. Abonnenc E. Les Phlébotomes de la région éthiopienne (Diptera, Psychodidae). In: Memoires ORSTOM series. Paris: Office de la Recherche Scientifique et Technique; 1972
 25. Lewis DJ. A taxonomic review of the genus *Phlebotomus* (Diptera: Psychodidae). Bull Br Mus Nat Hist. 1982;45:121–209.
 26. Lukes J, Mauricio IL, Schönian G, Dujardin JC, Soteriadou K, Dedet JP, et al. Evolutionary and geographical history of the *Leishmania donovani* complex with a revision of current taxonomy. Proc Natl Acad Sci USA. 2007;104:9375–80. <https://doi.org/10.1073/pnas.0703678104>
 27. Talmi-Frank D, Nasereddin A, Schnur LF, Schönian G, Töz SO, Jaffe CL, et al. Detection and identification of old world *Leishmania* by high resolution melt analysis. PLoS Negl Trop Dis. 2010;4:e581. <https://doi.org/10.1371/journal.pntd.0000581>
 28. Sagi O, Berkowitz A, Codish S, Novack V, Rashti A, Akad F, et al. Sensitive molecular diagnostics for cutaneous leishmaniasis. Open Forum Infect Dis. 2017;4:ofx037. <https://doi.org/10.1093/ofid/ofx037>
 29. Ben-Shimol S, Sagi O, Horev A, Avni YS, Ziv M, Riesenber K. Cutaneous leishmaniasis caused by *Leishmania infantum* in southern Israel. Acta Parasitol. 2016;61:855–8. <https://doi.org/10.1515/ap-2016-0118>
 30. el Tai NO, Osman OF, el Fari M, Presber W, Schönian G. Genetic heterogeneity of ribosomal internal transcribed spacer in clinical samples of *Leishmania donovani* spotted on filter paper as revealed by single-strand conformation polymorphisms and sequencing. Trans R Soc Trop Med Hyg. 2000;94:575–9. [https://doi.org/10.1016/S0035-9203\(00\)90093-2](https://doi.org/10.1016/S0035-9203(00)90093-2)
 31. Haralambous C, Antoniou M, Pralong F, Dedet JP, Soteriadou K. Development of a molecular assay specific for the *Leishmania donovani* complex that discriminates *L. donovani*/*Leishmania infantum* zymodemes: a useful tool for typing MON-1. Diagn Microbiol Infect Dis. 2008;60:33–42. <https://doi.org/10.1016/j.diagmicrobio.2007.07.019>
 32. Valinsky L, Ettinger G, Bar-Gal GK, Orshan L. Molecular identification of bloodmeals from sand flies and mosquitoes collected in Israel. J Med Entomol. 2014;51:678–85. <https://doi.org/10.1603/ME13125>
 33. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: molecular evolutionary genetics analysis across computing platforms. Mol Biol Evol. 2018;35:1547–9. <https://doi.org/10.1093/molbev/msy096>
 34. Ben-Ami R, Schnur LF, Golan Y, Jaffe CL, Mardi T, Zeltser D. Cutaneous involvement in a rare case of adult visceral leishmaniasis acquired in Israel. J Infect. 2002;44:181–4. <https://doi.org/10.1053/jinf.2002.0953>
 35. Guan LR, Xu YX, Li BS, Dong J. The role of *Phlebotomus alexandri* Sinton, 1928 in the transmission of kala-azar. Bull World Health Organ. 1986;64:107–12.
 36. Guan LR, Zhou ZB, Jin CF, Fu Q, Chai JJ. Phlebotomine sand flies (Diptera: Psychodidae) transmitting visceral leishmaniasis and their geographical distribution in China: a review. Infect Dis Poverty. 2016;5:15. <https://doi.org/10.1186/s40249-016-0107-z>
 37. Molina R, Jiménez MI, Cruz I, Iriso A, Martín-Martín I, Sevillano O, et al. The hare (*Lepus granatensis*) as potential sylvatic reservoir of *Leishmania infantum* in Spain. Vet Parasitol. 2012;190:268–71. <https://doi.org/10.1016/j.vetpar.2012.05.006>
 38. Sevá ADP, Martcheva M, Tuncer N, Fontana I, Carrillo E, Moreno J, et al. Efficacies of prevention and control measures applied during an outbreak in southwest Madrid, Spain. PLoS One. 2017;12:e0186372. <https://doi.org/10.1371/journal.pone.0186372>
 39. Tsokana CN, Sokos C, Giannakopoulos A, Mamuris Z, Birtsas P, Papaspyropoulos K, et al. First evidence of *Leishmania* infection in European brown hare (*Lepus europaeus*) in Greece: GIS analysis and phylogenetic position within the *Leishmania* spp. Parasitol Res. 2016;115:313–21. <https://doi.org/10.1007/s00436-015-4749-8>
 40. Rocchigiani G, Ebani VV, Nardoni S, Bertelloni F, Bascherini A, Leoni A, et al. Molecular survey on the occurrence of arthropod-borne pathogens in wild brown hares (*Lepus europaeus*) from Central Italy. Infect Genet Evol. 2018;59:142–7. <https://doi.org/10.1016/j.meegid.2018.02.005>
 41. Jambulingam P, Pradeep Kumar N, Nandakumar S, Paily KP, Srinivasan R. Domestic dogs as reservoir hosts for *Leishmania donovani* in the southernmost Western Ghats in India. Acta Trop. 2017;171:64–7. <https://doi.org/10.1016/j.actatropica.2017.03.006>
 42. Hassan MM, Osman OF, El-Raba' FM, Schallig HD, Elnaiem DE. Role of the domestic dog as a reservoir host of *Leishmania donovani* in eastern Sudan. Parasit Vectors. 2009;2:26. <https://doi.org/10.1186/1756-3305-2-26>
 43. Dereure J, El-Safi SH, Bucheton B, Boni M, Kheir MM, Davoust B, et al. Visceral leishmaniasis in eastern Sudan: parasite identification in humans and dogs; host-parasite


- relationships. *Microbes Infect.* 2003;5:1103–8. <https://doi.org/10.1016/j.micinf.2003.07.003>
44. Bsrat A, Berhe M, Gadissa E, Taddele H, Tekle Y, Hagos Y, et al. Serological investigation of visceral *Leishmania* infection in human and its associated risk factors in Welkait District, Western Tigray, Ethiopia. *Parasite Epidemiol Control.* 2017;3:13–20. <https://doi.org/10.1016/j.parepi.2017.10.004>
 45. Bashaye S, Nombela N, Argaw D, Mulugeta A, Herrero M, Nieto J, et al. Risk factors for visceral leishmaniasis in a new epidemic site in Amhara Region, Ethiopia. *Am J Trop Med Hyg.* 2009;81:34–9. <https://doi.org/10.4269/ajtmh.2009.81.34>
 46. Kalayou S, Tadelles H, Bsrat A, Abebe N, Haileselassie M, Schallig HDFH. Serological evidence of *Leishmania donovani* infection in apparently healthy dogs using direct agglutination test (DAT) and rk39 dipstick tests in Kafta Humera, north-west Ethiopia. *Transbound Emerg Dis.* 2011;58:255–62. <https://doi.org/10.1111/j.1865-1682.2011.01209.x>
 47. Magri A, Galuppi R, Fioravanti M, Caffara M. Survey on the presence of *Leishmania* sp. in peridomestic rodents from the Emilia-Romagna Region (northeastern Italy). *Vet Res Commun.* 2023;47:291–6. <https://doi.org/10.1007/s11259-022-09925-4>
 48. Özbilgin A, Çavuş İ, Yıldırım A, Gündüz C. Do the rodents have a role in transmission of cutaneous leishmaniasis in Turkey? [in Turkish]. *Mikrobiyol Bul.* 2018;52:259–72. <https://doi.org/10.5578/mb.66828>
 49. Frézard F, Aguiar MMG, Ferreira LAM, Ramos GS, Santos TT, Borges GSM, et al. Liposomal amphotericin B for treatment of leishmaniasis: from the identification of critical physicochemical attributes to the design of effective topical and oral formulations. *Pharmaceutics.* 2022;15:99. <https://doi.org/10.3390/pharmaceutics15010099>

Address for correspondence: Liora Studentsky, Laboratory of Medical Entomology, Ministry of Health, Elaiv Yaakov 9, Jerusalem 9546208, Israel; email: liora.studenetsky@moh.gov.il

etymologia revisited

Dermatophilus congolensis

[dur"mə-tof'ī-ləs con-gō-len'sis]



Originally published
in August 2022

From the Greek *derma* (skin) + *philos* (loving), *Dermatophilus congolensis* is a Gram-positive, aerobic actinomycete, and facultatively anaerobic bacteria (Figure 1). *D. congolensis* infects the epidermis and produces exudative dermatitis termed dermatophilosis that was previously known as rain rot, rain scald, streptotrichosis, and mycotic dermatitis.

In 1915, René Van Saceghem (Figure 2), a Belgian military veterinarian stationed at a veterinary laboratory in the former Belgian Congo (thus, the species name *congolensis*), reported *D. congolensis* from exudative dermatitis in cattle. Local breeders and veterinarians had observed the disease since 1910, but the causal agent was not identified.

Dermatophilosis affects animals, mainly cattle, and more rarely humans. Outbreaks of *D. congolensis* infection have severe economic implications in the livestock and leather industries.

References:

1. Amor A, Enríquez A, Corcuera MT, Toro C, Herrero D, Baquero M. Is infection by *Dermatophilus congolensis* underdiagnosed? *J Clin Microbiol.* 2011;49:449–51.
2. Branford I, Johnson S, Chapwanya A, Zayas S, Boyen F, Mielcarska MB, et al. Comprehensive molecular dissection of *Dermatophilus congolensis* genome and first observation of tet(z) tetracycline resistance. *Int J Mol Sci.* 2021;22:7128.
3. Dorland's illustrated medical dictionary. 32nd ed. Philadelphia: Elsevier Saunders; 2012.
4. Van Saceghem R. Contagious skin disease (contagious impetigo) [in French]. *Bull Soc Pathol Exot.* 1915;8:354–9.

https://wwwnc.cdc.gov/eid/article/28/8/et-2808_article