



# HHS Public Access

Author manuscript

*Depress Anxiety*. Author manuscript; available in PMC 2023 June 01.

Published in final edited form as:

*Depress Anxiety*. 2022 June ; 39(6): 536–547. doi:10.1002/da.23273.

## Depressive and Anxious Symptoms Among Young Adults in the COVID-19 Pandemic: Results from Monitoring the Future

Noah T. Kreski, MPH<sup>1,\*</sup>, Katherine M. Keyes, PhD<sup>1</sup>, Michael J. Parks, PhD<sup>2</sup>, Megan E. Patrick, PhD<sup>3</sup>

<sup>1</sup>Columbia University, Department of Epidemiology, Mailman School of Public Health, New York, NY

<sup>2</sup>University of Minnesota, Center for Applied Research and Educational Improvement, Minneapolis, MN

<sup>3</sup>University of Michigan, Institute for Social Research, Ann Arbor, MI

### Abstract

**Purpose.**—The SARS-CoV-2 pandemic is associated with worsening mental health among young adults, but further research is necessary to quantify the associations with depression and anxiety.

**Methods.**—Using Monitoring the Future data (N=1,244 young adults, modal age: 19, Fall 2020 supplement), we examined internalizing symptoms (PHQ-8 and GAD-7 separately), dividing the sample into those without clinically significant scores, significant scores but minimal pandemic-attributed symptoms, and significant scores with substantial pandemic-attributed symptoms. Logistic regression analyses linked demographic factors, pandemic-related experiences, and coping methods to symptom groups.

**Results.**—Internalizing symptoms were highly prevalent, with many occurring among a majority at least several days over the past two weeks. Major changes in education, employment, and resource availability predicted elevated symptom risk (e.g., lacking a place to sleep or money for rent, gas, or food led to 4.43 [95% CI:2.59-7.55] times the risk of high depressive symptoms significantly attributed to the pandemic). High internalizing symptoms were linked to underutilization of healthy coping behaviors, substance use overutilization, and dietary changes. High depressive and anxious symptoms attributed to the pandemic were marked by high levels of taking breaks from the news/social media and contacting healthcare providers.

**Conclusions.**—The pandemic's associations with young adults' depressive and anxious symptoms warrants urgent attention through improved mental health treatment infrastructure and stronger structural support.

\*Corresponding Author: Noah T. Kreski, Columbia University, Mailman School of Public Health, Department of Epidemiology, ntk2109@cumc.columbia.edu.

Conflict of Interest: Authors have no conflicts of interest to disclose. Study sponsors had no role in the study design, data collection, analysis or interpretation, report writing, or the decision to submit this manuscript for publication.

Ethics: Monitoring the Future is approved by the IRB of University of Michigan.

## Keywords

COVID-19; Depression; Anxiety; Epidemiology; Coping

---

## Introduction:

The SARS-CoV-2 pandemic is associated with worsening mental health among young adults, but further research is necessary to quantify the associations with depression and anxiety. Available evidence suggests that depressive and anxious symptoms increased among adults during the course of the pandemic, with the highest increases among adults ages 18 to 29.<sup>1</sup> Depressive and anxious symptoms among young adults increased in the presence of educational, financial, and other pandemic-related stressors,<sup>2</sup> and meta-analytic estimates suggest that 34% and 32% of young adults have reported depressive and anxious symptoms, respectively, during the pandemic.<sup>3</sup> However, the specific domains of depressive or anxious symptoms most directly affected by the pandemic are a gap in the literature that can inform intervention efforts. Similarly, it remains unclear the extent to which highly prevalent depressive and anxious symptoms are primarily related to the pandemic, rather than symptoms that would arise regardless of the pandemic. Understanding the specific symptoms and individuals most directly affected by pandemic-related stressors provides important context for future mental health interventions.

The COVID-19 pandemic has disrupted young people's social, educational, and occupational practices, at times leading to isolation, remote learning and employment, and changes to work hours or compensation. Job insecurity in the pandemic predicts higher depression and anxiety symptoms among young adults in the US,<sup>4</sup> and pandemic-related academic disruptions and uncertainty can lead to psychological distress that results in young adults reducing or stopping their engagement with education.<sup>5</sup> These, along with other pandemic-related experiences, including economic hardship and resource scarcity, have worsened young adults' mental health, though further study is necessary to understand the full magnitude of their link with clinically significant levels of depressive or anxious symptoms. Additionally, while understanding the role of the COVID-19 pandemic for these symptoms is important, it is perhaps more crucial to identify and implement effective coping strategies against pandemic-related stressors. While some techniques, such as connecting socially with others<sup>6</sup> and maintaining a healthy diet,<sup>7</sup> have been reported by young adults for improved mental wellbeing, others strategies, such as substance use to cope with pandemic-related stress, are potentially damaging and can lead to future substance use problems.<sup>8</sup> Furthermore, less is known about the extent to which coping behaviors vary between those with pandemic-related depressive and anxious symptoms and those with similar symptoms not attributed to the pandemic.

In order to understand the role of COVID-19 pandemic-related experiences in young adults' depressive and anxious symptoms, this study had the following aims: (1) Estimate the distribution of depressive and anxious symptoms in this population and determine the extent to which these symptoms are attributed by young adults to the COVID-19 pandemic; (2) Disentangle the demographic risk profiles for depressive and anxious symptoms

that are attributed to pandemic-related experiences vs. not; (3) Understand the extent to which specific pandemic-related experiences have elevated risk for depressive and anxious symptoms; and (4) Link patterns of depressive and anxious symptoms to various coping behaviors to recognize potential areas for intervention.

## Materials and Methods

Monitoring the Future (MTF), an ongoing study of adolescent beliefs and behaviors,<sup>9</sup> included a unique supplemental study in Fall 2020 that featured items relating to the COVID-19 pandemic.<sup>8</sup> The MTF Vaping Supplement collected data between September and November 2020. Participants were recruited from grade 12 students participating in the Spring 2019 MTF survey. Of 13,713 MTF participants in grade 12, there were 7,850 eligible for the Vaping Supplement. Ineligibility criteria included missing contact information or random selection into MTF's longitudinal study. Of eligible participants, 4,358 were selected at random, oversampling those who reported vaping or other substance use. Of these participants, 1,244 (28.5%) participated in a 1-year follow-up survey, which provides the data for these analyses. These participants ranged in age from 18 to 24, though the vast majority (93%) were 19 or 20. The Institutional Review Board of University of Michigan approved this study. This supplement's baseline survey captured a broad array of internalizing symptom items, as well as information on coping behaviors, educational trajectories, and economic outcomes related to the COVID-19 pandemic. This sample oversampled 12<sup>th</sup> grade students who vaped, but survey weights accounted for selection probabilities.

### Measures:

**Internalizing Symptoms**—Depressive symptoms were assessed with the Patient Health Questionnaire (PHQ-8)<sup>10</sup> and anxious symptoms were assessed with the Generalized Anxiety Disorder Scale (GAD-7).<sup>11</sup> These scales have been used to assess depression and anxiety in a wide variety of contexts, providing meaningful screening tools that capture core affective and behavioral symptoms in psychiatric settings<sup>12,13</sup> and general population surveys alike.<sup>10,14,15</sup> Full items for each of these scales can be seen in Supplemental Table A1. These scales were structured similarly, with each asking “Over the past two weeks, how often have you been bothered by the following problems?” before reporting each symptom. Responses ranged from “not at all” (score of 0) to “several days” (1), “more than half the days” (2), and “nearly every day” (3). Each scale also generated a total score, adding the scores for every item together. The PHQ-8 scale exhibits strong sensitivity and specificity for major depression (88% for both metrics)<sup>10</sup> with a validated cutpoint of a total score 10 (out of 24). The GAD-7 had a sensitivity of 89% for diagnosing generalized anxiety disorder in the general population, and a specificity of 82% with a validated cutpoint of a total score 10 (out of 21).<sup>15</sup> We queried the PHQ-8 (which excludes an item on suicidality) rather than the PHQ-9, as we were unable to conduct follow-up phone calls with people who indicate suicidality. This constituted a human subjects and liability concern, gathering that information without the ability to react to it in real time. Prior research has demonstrated the strong equivalency of the PHQ-8 and PHQ-9 in terms of scores, sensitivity, and specificity, suggesting minimal effect on our results.<sup>16</sup>

Groups were divided further based on the extent to which their symptoms were attributed to the pandemic. For each PHQ-8 and GAD-7 item, respondents were asked “How much has COVID-19 influenced the following problems?” with response options of “No influence” (score of 0), “Somewhat” (1), “Very much (2), and “Completely” (3). These scores were summed for each scale to generate an index of symptom attribution to the COVID-19 pandemic. We used this index divide those with significantly high symptoms into two groups: those whose symptoms were more than somewhat attributed to the pandemic, and those whose symptoms were somewhat or not related attributed to the pandemic. This threshold for overall symptoms being “more than somewhat” attributed to the COVID pandemic was a COVID attribution index >8 for PHQ-8 and >7 for GAD-7. Ultimately, this resulted in three groups each for analyses of depressive and anxious symptoms: those without high symptoms, those with high symptoms more than somewhat attributed to the pandemic, and those with high symptoms *not* attributed to the pandemic more than somewhat. While some prior studies have implemented items examining attribution of life changes and mental health to the pandemic among young adults,<sup>17,18</sup> these have typically used a single item rather than a more comprehensive index relating to the attribution of specific symptoms to the pandemic.

**Demographic Variables**—Demographic variables were included to assess heterogeneity of depressive and anxious symptom group membership by sex, race/ethnicity, parental education and urbanicity. These variables were also included as covariates in adjusted models. Responses were coded as male and female for sex. Race and ethnicity were pulled from two items: for ethnicity, people answered “Are you Hispanic or Latino/a?” with a yes or no. Then, for race, people selected all that applied of this list: Asian/South Asian, Native Hawaiian or Other Pacific Islander, Black or African American, American Indian or Alaskan Native, white, Arab/Middle Eastern/North African, or Other. These responses were then coded into the following groups: white, Hispanic/Latino, Black, Asian, and Other. Responses were coded as “college or more” and “some college or less” for highest level of parental education. For urbanicity, responses were coded as rural, a small or medium city, a suburb of a small or medium city, a large or very large city, a suburb of a very large city, or “can’t say”/mixed.

**Pandemic-Related Experiences**—Respondents described the effect that the COVID-19 pandemic had on their education, employment, and other areas of their lives. We examined the educational effect of the pandemic with two items. First, respondents who were current students were asked “In Spring 2020, how (if at all) did your student status change because of the COVID-19 pandemic?” The outcome of interest was dropping classes or experiencing any other major change to student status. The second educational item asked everyone “How (if at all), has the COVID-19 pandemic changed your student status or educational plans for Fall 2020?” Here, the outcome of interest was any major effect on educational plans defined by a change in course load (either reduced or increased), enrollment (dropping out, taking a gap term, or enrolling at a new school), or any other component of Fall 2020 educational plans. We examined employment with the item “Since March 2020, how has the COVID-19 pandemic affected your work?” before providing a list of potential effects that covered working remotely, changes to hours worked, loss of employment, issues with compensation

or benefits, or difficulty arranging or affording childcare. Responses were condensed into “Any significant effect on employment since March 2020” vs. “None”. Additional COVID-related experiences were examined with the item “Have you experienced any of the following as a result of the COVID-19 pandemic?”. Responses were dichotomized into “Lacked a place to sleep or money for rent, gas, or food due to COVID-19” vs. “None of the above”.

**Coping**—Respondents were asked “To cope with social distancing and isolation, are you doing any of the following?” and then given a list of fourteen coping strategies, providing yes or no answers to each. These potential coping strategies included physical behaviors (e.g., taking care of your body, such as taking deep breaths, stretching, or meditating), social practices (e.g., connecting with others, including talking with people you trust about your concerns and how you are feeling), substance use (e.g., vaping more), and dietary behavior (e.g., eating more food than usual).

## Analyses

All analyses incorporated survey weights that addressed oversampling of those who vaped, as well as other elements of the complex survey design. Weights were calculated as the inverse of the probability of participation at follow-up based on covariates measured in 12th grade: birth-assigned sex, race/ethnicity, college plans, truancy, high school grades, number of parents in the home, religiosity, parental education, substance use, region, cohort, and sampling weight. Implementation of these weights, which account for non-response bias between grade 12 and follow-up, allows for generalization back to the grade 12 MTF sample, which is nationally representative.

We analyzed the weighted prevalence of those with each symptom over the past two weeks and then divided those with each symptom based on whether respondents reported that their symptom was more than somewhat pandemic-attributable. In order to examine the association between COVID-19 and depressive or anxious symptoms overall, we examined multinomial logistic regressions for each category of symptoms separately in Stata, producing relative risk ratios (RRR). That is, for depressive and anxious symptoms, multinomial logistic regressions examined the outcome of membership in three groups: those without high symptoms, those with high symptoms more than somewhat attributed to the pandemic, and those with high symptoms *not* attributed to the pandemic more than somewhat. First, symptom outcomes were predicted by each demographic factor separately to examine group differences in depressive and anxious symptoms. Then, separate models linked pandemic-related experiences to these outcomes, adjusting for demographics. Lastly, a final group of models predicted these symptom group outcomes using coping behaviors, adjusting for demographics, to examine how coping practices were associated with symptom outcomes. Analyses used pairwise deletion, given minimal missingness (<10% overall for any given model).

## Results

Sample characteristics for the 1,244 young adults in this study are reported in Table 1. This sample was predominantly male sex assigned at birth (55.4%), white (57.2%), had a

parent who completed college (55.2%), and were most likely to be found in an urban setting compared to suburban or rural settings (41.2% in cities of any size).

Depressive symptoms were highly prevalent in this sample (Figure 1), ranging from 20.6% prevalence in the past two weeks (moving or speaking slowly or being fidgety and restless) up to 67.6% (feeling tired or having little energy). Six out of eight symptoms were prevalent in a majority of respondents (little interest or pleasure in things, feeling down/depressed/hopeless, sleep issues, feeling tired, poor appetite or overeating, and feeling bad about yourself). Dividing those with symptoms based on whether the symptoms were attributed more than somewhat to the pandemic, results showed that pandemic-attributed depressive symptoms were most common for little interest/pleasure in things (14.7% of respondents had this symptom attributed more than somewhat to the pandemic) and feeling down, depressed, or hopeless (18.8% had this symptom attributed more than somewhat to the pandemic).

Anxious symptoms were also highly prevalent (Figure 2), ranging from 31.6% of respondents (being so restless that it's hard to sit still) to 56.7% (feeling nervous, anxious, or on edge). Pandemic-attributed anxious symptoms were most common for worrying too much about different things (14.8% had this symptom attributed more than somewhat to the pandemic) and feeling afraid, as if something awful might happen (14.4% had this symptom more than somewhat due to the pandemic). Clinically high depressive symptoms were evident among 28.0% of respondents, while clinically high anxiety symptoms were seen among 20.8%. Relative to meta-analytic estimates in this population, studies using the PHQ-9 with the appropriate threshold for clinically high depression had an average prevalence of 28%, which matches our estimate precisely.<sup>3</sup> While the meta-analytic estimate for GAD-7 studies predicting prevalence of anxiety still produced a relatively higher estimate (32% vs our estimate of 20.8%), several included studies featured an estimate in the low twenties, suggesting that our results fall within the natural heterogeneity of studies on this topic.<sup>3</sup>

Table 2 shows results for multinomial regressions predicting the three membership groups (those without high symptoms, those with high symptoms more than somewhat attributed to the pandemic, and those with high symptoms *not* attributed to the pandemic more than somewhat) by demographic factors of sex, race/ethnicity, parental education and urbanicity. For sex, the risk of depressive and anxious symptoms, either primarily attributed to the pandemic or not, was higher for female respondents (e.g., RRR for high depressive symptoms more than somewhat attributed to the pandemic [female vs. male]: 1.74 [95% CI: 1.08, 2.82]). For race and ethnicity, risk of membership in either high symptom group was particularly elevated for respondents in the "Other" group (e.g., American Indian, Alaskan Native, Pacific Islander, and Middle Eastern respondents) compared to Black peers who had the lowest rates of significant depressive or anxious symptoms. Depressive and anxious symptoms were also elevated for white, Hispanic/Latino, and Asian respondents compared to Black respondents. For urbanicity, risk of depressive and anxious symptoms, either attributed to the pandemic or not, was consistently highest in the "Can't say"/mixed group.

The risk profile for high internalizing symptoms attributed to the pandemic across sex, race/ethnicity, and urbanicity was similar to the risk profile for high internalizing symptoms without major pandemic attribution. However, for parental education, there was evidence of heterogeneous risk profiles for depressive symptoms. Risk of depressive symptoms that were not/minimally COVID-attributed was lower for those with higher parental education (RRR [College or more vs. some college or less]: 0.67 [95% CI: 0.45, 0.98]). Risk for depressive symptoms that were highly COVID-related, however, was higher for those with higher parental education.

Table 3 demonstrates the amplified risk of depressive and anxious symptoms due to major shifts in education, employment, and other economic effects of the pandemic. Changes to student status and educational plans in both Spring 2020 and Fall 2020 were each associated with elevated risk for depressive and anxious symptoms, especially clinically high symptoms attributed more than somewhat to the pandemic. For instance, dropping classes or other major change to student status in Spring 2020 was linked to a demographic-adjusted RRR of 2.57 (95% CI: 1.37, 4.81) for depressive symptoms attributed more than somewhat to the pandemic. The RRR was even higher for those who experienced resource scarcity due to the pandemic, such as lacking a place to sleep or money for rent, gas, or food elevated risk of internalizing symptoms (aRRR for high depressive symptoms attributed more than somewhat to the pandemic: 4.43 [95% CI: 2.59, 7.55]). While these experiences posed a particularly major risk for anxious and depressive symptoms among young adults, these outcomes were also elevated among those who experienced more general changes to employment, such as shifts in hours worked or compensation (aRRR for high anxious symptoms attributed more than somewhat to the pandemic due to major effect on employment since March 2020: 2.24 [95% CI: 1.27, 3.94]).

Table 4 shows the associations between the utilization of coping practices and internalizing symptom outcomes. Notably, many forms of healthy coping rooted in physical or social behaviors were linked to a reduced risk of high symptoms (e.g., aRRR for “Taking care of your body, such as taking deep breaths, stretching, or meditating” predicting high depressive symptoms more than somewhat attributed to the pandemic: 0.57 [95% CI: 0.36, 0.92]). Meanwhile, coping methods rooted in substance use and dietary changes were typically linked to higher symptoms, especially for high symptoms more than somewhat attributed to the pandemic (e.g., aRRR for “Vaping more” predicting high anxious symptoms more than somewhat attributed to the pandemic: 4.76 [95% CI: 2.08, 10.93]).

While the link between coping strategies and internalizing symptoms was typically similar regardless of whether symptoms were primarily attributed to the pandemic, there were two notable coping strategies with heterogeneous links to depressive symptoms based on whether those symptoms were pandemic-attributed. Taking breaks from watching, reading, or listening to news stories, including social media was associated with mildly decreased risk of depressive symptoms not attributed to the pandemic (aRRR for high depressive symptoms = 0.88 [95% CI: 0.59, 1.31]), but increased the risk for depressive symptoms attributed to the pandemic (aRRR for high depressive symptoms more than somewhat attributed to the pandemic: 1.60 [95% CI: 1.00, 2.57]). A similar dynamic occurred for the coping mechanism of contacting a healthcare provider. For anxious symptoms, contacting a

healthcare provider was strongly linked to anxious symptoms attributed more than somewhat to the pandemic (aRRR = 2.14 [95% CI: 1.03, 4.47]) but not for anxious symptoms not/somewhat attributed to the pandemic (aRRR = 1.03 [95% CI: 0.47, 2.28]).

## Discussion

In the current SARS-CoV-2 pandemic, young adults have experienced high levels of depressive and anxious symptoms, and a portion of these symptoms could be attributed to the effects of the pandemic (e.g., educational, financial, housing-related). Risk of these symptoms was overall higher for female (vs. male) respondents, and for white, Hispanic/Latino, Asian, or “Other” race/ethnicity (vs. Black) respondents. Pandemic-related depressive symptoms specifically were slightly elevated for those with high parental education, while the opposite was true for depressive symptoms not attributed to the pandemic more than somewhat. Depressive and anxious symptoms, particularly when pandemic-attributable, were strongly predicted by significant changes to educational plans or practices, employment, or resource availability due to the pandemic. Healthy physical or social coping mechanisms were linked to reduced risk of these symptoms, while substance use and dietary changes were linked to elevated risk. Taking breaks from the news and contacting healthcare professionals were common coping behaviors highly prevalent among those whose depressive and anxious symptoms were mainly due to the pandemic. Ultimately, these findings reflect the need for urgent, major structural and social changes to improve the mental health of young adults in the pandemic.

Still, most people with each depressive or anxious symptom in this study reported that they were minimally attributable to the pandemic. These substantial symptoms unrelated to the pandemic reflect consistently increasing pre-pandemic trends in depressive and anxious symptoms.<sup>19,20</sup> However, adverse pandemic-related experiences were linked to elevated risk for clinically high depressive and anxious symptoms, regardless of whether young adults attributed their symptoms primarily to the pandemic. Thus, benefits of pandemic-related interventions and structural changes to address pandemic-related stressors may benefit all young people with clinically high depressive and anxious symptoms.

The expansion of accessible, affordable, equitable mental healthcare with remote telehealth options that address pandemic-related stressors would be useful, alleviating psychological distress for any number of young adults.<sup>21</sup> Additionally, interventions are needed to support productive coping methods and address maladaptive coping approaches that are consequences of the pandemic. Based on the coping mechanism results of our analyses, public health professionals should work towards the promotion of physical wellness and social connection for those distressed by the pandemic, while still emphasizing the value of social distancing and protective measures. As pandemic-related clinically high depressive and anxious symptoms were linked to higher rates of contacting healthcare providers and taking breaks from the news, there may also be an unmet need for succinct, fact-based, readily accessible information on pandemic-related news and recommendations. Additionally, just as healthy coping behaviors should be promoted, we should also monitor elevated substance use and disrupted eating patterns which, in this study, were associated with elevated depressive and anxious symptoms, pandemic-related or not. Public health



organizations should boost visibility for accessible resources to address those behaviors, such as text or video-based therapies or guided treatment plans from a connected, collaborative network of informed clinicians.<sup>22,23</sup>

The suggestions described above would help young adults to healthily cope with the pandemic, but our results also highlight that young adults who report struggling with pandemic-related effects, like not having a place to sleep or money to buy gas or food, have greater risk for depressive and anxious symptoms. Therefore, there is also a need for stronger structural supports to address the socioeconomic and educational effects of COVID. Given the associations between educational changes and clinically high depressive and anxious symptoms, further research should examine whether the widespread implementation of specific policies, such as flexibility of coursework and grading, support for effective remote learning, and school-based support for mental health counselling, would have mental health benefits.<sup>24,25</sup> To address the harmful effects of employment/income shifts and resource scarcity, it is likely worthwhile to examine the potential benefits of increased governmental economic stimulus sent directly to citizens, as well as expansion of insurance coverage, unemployment benefits, housing support, and employment opportunities.<sup>26-29</sup> These economic resources are vital for the improvement of young adults' mental health, though notably young people with higher parental education, a proxy for socioeconomic status, had higher pandemic-related depressive symptoms. Further research should examine this interplay of diverse socioeconomic metrics and depressive and anxious symptoms.

This study had several strengths, including the use of thorough, validated measures for depressive and anxious symptoms. In addition to these scales, the measures of pandemic-related educational, employment, economic experiences, and coping were thorough, covering a broad array of potential experiences and behaviors. Still, as with any study, there were limitations. This measure of pandemic attribution for each symptom was self-reported, and the true extent to which a depressive or anxious symptom is due primarily to the pandemic may not align with a respondent's perception. Future research may include a clinician interview or more thorough questions that examine participant history, which may influence the perception of the extent to which symptoms were truly attributable to the pandemic. We were also unable to examine mental health treatment in this sample, which may impact the relationship between pandemic-related stressors and internalizing symptom outcomes. Given the cross-sectional nature of these data, we were unable to disentangle the directionality of associations or the durations of symptoms. These results do not necessarily apply to other age groups or countries, as pandemic-related experiences likely differ substantially. Additionally, people who reported substance use at baseline were oversampled, though survey weighting accounted for this oversampling and adjusted results accordingly.

## Conclusion

The COVID-19 pandemic added to the already substantial depressive and anxious symptoms of young adults. This psychiatric harm warrants urgent interventions that not only support effective coping practices, but provide tangible socioeconomic resources.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Funding Statement:

Data collection and manuscript preparation were supported by research grants from the National Institute on Drug Abuse: R01DA001411, R01DA016575, and R01DA048853; additional funding was provided by National Center for Injury Prevention and Control grant R49-CE003094. The study sponsors had no role in the study design, collection, analysis or interpretation of the data, writing of the manuscript, or the decision to submit the paper for publication. The content is solely the responsibility of the authors and does not necessarily represent the official views of the study sponsor.


## Data Availability:

De-identified Monitoring the Future data are publicly available for download.

## References


- Vahratian A, Blumberg SJ, Terlizzi EP, Schiller JS. Symptoms of Anxiety or Depressive Disorder and Use of Mental Health Care Among Adults During the COVID-19 Pandemic - United States, August 2020-February 2021. *MMWR Morb Mortal Wkly Rep.* 2021;70(13):490-494. doi:10.15585/mmwr.mm7013e2 [PubMed: 33793459]
- Kujawa A, Green H, Compas BE, Dickey L, Pegg S. Exposure to COVID-19 pandemic stress: Associations with depression and anxiety in emerging adults in the United States. *Depress Anxiety.* 2020;37(12):1280-1288. [PubMed: 33169481]
- Deng J, Zhou F, Hou W, et al. The prevalence of depressive symptoms, anxiety symptoms and sleep disturbance in higher education students during the COVID-19 pandemic: A systematic review and meta-analysis. *Psychiatry Res.* 2021;301:113863. doi:10.1016/j.psychres.2021.113863 [PubMed: 33984824]
- Ganson KT, Tsai AC, Weiser SD, Benabou SE, Nagata JM. Job Insecurity and Symptoms of Anxiety and Depression Among U.S. Young Adults During COVID-19. *J Adolesc Heal.* 2021;68(1):53-56. doi:10.1016/j.jadohealth.2020.10.008
- Clabaugh A, Duque JF, Fields LJ. Academic Stress and Emotional Well-Being in United States College Students Following Onset of the COVID-19 Pandemic. *Front Psychol.* 2021;12:628787. doi:10.3389/fpsyg.2021.628787 [PubMed: 33815214]
- Waselewski EA, Waselewski ME, Chang T. Needs and Coping Behaviors of Youth in the U.S. During COVID-19. *J Adolesc Heal.* 2020;67(5):649-652. doi:10.1016/j.jadohealth.2020.07.043
- Fullana MA, Hidalgo-Mazzei D, Vieta E, Radua J. Coping behaviors associated with decreased anxiety and depressive symptoms during the COVID-19 pandemic and lockdown. *J Affect Disord.* 2020;275:80-81. doi:10.1016/j.jad.2020.06.027 [PubMed: 32658829]
- Patrick ME, Parks MJ, Fairlie AM, Kreski NT, Keyes KM, Miech R. Using substances to cope with the CoViD-19 pandemic: US national data at age 19 years. *J Adolesc Heal.* 2021.
- Miech RA, Johnston LD, O'Malley PM, Bachman JG, Schulenberg JE, Patrick ME. Monitoring the Future national survey results on drug use, 1975-2020: Volume I, Secondary school students. *Ann Arbor Inst Soc Res Univ Michigan*, 571 pp. 2021.
- Kroenke K, Strine TW, Spitzer RL, Williams JBW, Berry JT, Mokdad AH. The PHQ-8 as a measure of current depression in the general population. *J Affect Disord.* 2009;114(1-3):163-173. [PubMed: 18752852]
- Spitzer RL, Kroenke K, Williams JBW, Löwe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med.* 2006;166(10):1092-1097. [PubMed: 16717171]
- Shin C, Lee S-H, Han K-M, Yoon H-K, Han C. Comparison of the usefulness of the PHQ-8 and PHQ-9 for screening for major depressive disorder: analysis of psychiatric outpatient data. *Psychiatry Investig.* 2019;16(4):300.

13. Johnson SU, Ulvenes PG, Øktedalen T, Hoffart A. Psychometric Properties of the General Anxiety Disorder 7-Item (GAD-7) Scale in a Heterogeneous Psychiatric Sample . *Front Psychol* . 2019;10. <https://www.frontiersin.org/article/10.3389/fpsyg.2019.01713>.
14. Arias-de la Torre J, Vilagut G, Ronaldson A, et al. Prevalence and variability of current depressive disorder in 27 European countries: a population-based study. *Lancet Public Heal*. 2021;6(10):e729–e738.
15. Löwe B, Decker O, Müller S, et al. Validation and standardization of the Generalized Anxiety Disorder Screener (GAD-7) in the general population. *Med Care*. 2008;266–274. [PubMed: 18388841]
16. Wu Y, Levis B, Riehm KE, et al. Equivalency of the diagnostic accuracy of the PHQ-8 and PHQ-9: a systematic review and individual participant data meta-analysis. *Psychol Med*. 2020;50(8):1368–1380. [PubMed: 31298180]
17. Jardon C, Choi KR. COVID-19 Experiences and Mental Health Among Graduate and Undergraduate Nursing Students in Los Angeles. *J Am Psychiatr Nurses Assoc*. January 2022;10783903211072222. doi:10.1177/10783903211072222
18. Haikalas M, Doucette H, Meisel MK, Birch K, Barnett NP. Changes in College Student Anxiety and Depression From Pre- to During-COVID-19: Perceived Stress, Academic Challenges, Loneliness, and Positive Perceptions. *Emerg Adulthood*. 2021;10(2):534–545. doi:10.1177/21676968211058516
19. Goodwin RD, Weinberger AH, Kim JH, Wu M, Galea S. Trends in anxiety among adults in the United States, 2008–2018: Rapid increases among young adults. *J Psychiatr Res*. 2020;130:441–446. doi:10.1016/j.jpsychires.2020.08.014 [PubMed: 32905958]
20. Keyes KM, Gary D, O'Malley PM, Hamilton A, Schulenberg J. Recent increases in depressive symptoms among US adolescents: trends from 1991 to 2018. *Soc Psychiatry Psychiatr Epidemiol*. 2019. doi:10.1007/s00127-019-01697-8
21. Pfender E. Mental Health and COVID-19: Implications for the Future of Telehealth. *J Patient Exp*. 2020;7(4):433–435. doi:10.1177/2374373520948436 [PubMed: 33062854]
22. Oesterle TS, Kolla B, Risma CJ, et al. Substance Use Disorders and Telehealth in the COVID-19 Pandemic Era: A New Outlook. *Mayo Clin Proc*. 2020;95(12):2709–2718. doi:10.1016/j.mayocp.2020.10.011 [PubMed: 33276843]
23. Johns G, Taylor B, John A, Tan J. Current eating disorder healthcare services – the perspectives and experiences of individuals with eating disorders, their families and health professionals: systematic review and thematic synthesis. *BJPsych Open*. 2019;5(4):e59. doi:DOI: 10.1192/bjo.2019.48 [PubMed: 31530301]
24. Lederer AM, Hoban MT, Lipson SK, Zhou S, Eisenberg D. More Than Inconvenienced: The Unique Needs of U.S. College Students During the COVID-19 Pandemic. *Heal Educ Behav*. 2020;48(1):14–19. doi:10.1177/1090198120969372
25. Halliburton AE, Hill MB, Dawson BL, Hightower JM, Rueden H. Increased stress, declining mental health: Emerging adults' experiences in college during COVID-19. *Emerg Adulthood*. 2021;9(5):433–448.
26. Masand P, Patkar A, Tew C, Hoerner A, Szabo ST, Gupta S. Mental Health and COVID-19: Challenges and Multimodal Clinical Solutions. *J Psychiatr Pract*. 2021;27(4):254–264. doi:10.1097/PRA.0000000000000560 [PubMed: 34398575]
27. Woolhandler S, Himmelstein DU. Intersecting U.S. Epidemics: COVID-19 and Lack of Health Insurance. *Ann Intern Med*. 2020;173(1):63–64. doi:10.7326/M20-1491 [PubMed: 32259195]
28. Cuellar A, Mark TL, Sharfstein SS, Huskamp HA. How to mitigate the mental health care consequences of the COVID-19 financial crisis. *Psychiatr Serv*. 2020;71(12):1317–1319. [PubMed: 32988321]
29. Kim D. Financial hardship and social assistance as determinants of mental health and food and housing insecurity during the COVID-19 pandemic in the United States. *SSM - Popul Heal*. 2021;16:100862. doi:10.1016/j.ssmph.2021.100862



■ Has Symptom, More Than Somewhat Attributed to the Pandemic
 ■ Has Symptom, Not or Somewhat Attributed to the Pandemic
 ■ No Symptom

**Figure 1:**  
Depressive Symptoms and Pandemic Attribution (with Weighted Percentages)



**Figure 2:**  
Anxiety Symptoms and Pandemic Attribution (with Weighted Percentages)

**Table 1:**

Sample Characteristics (Overall N = 1,244 young adults, ages: 17-22, MTF Supplement 2020)

Demographic Characteristics (Weighted %)		
Sex *		
	Female	44.1
	Male	55.4
Race/Ethnicity *		
	White	57.2
	Hispanic/Latino	18.0
	Black	7.9
	Asian	5.4
	Other	10.7
Parental Education *		
	College or more	55.2
	Some college or less	39.8
Urbanicity *		
	Rural	16.2
	A small or medium city	22.8
	A suburb of a small or medium city	18.7
	A large or very large city	18.4
	A suburb of a large or very large city	14.3
	Can't say, mixed	8.4
Mental Health Characteristics (%)		
	Clinically High Depressive Symptoms, PHQ-8 (Score 10)	28.0
	Clinically High Anxiety Symptoms, GAD-7 (Score 10)	20.8
Coping Strategies (%)		
	Making time to relax	61.9
	Connecting with others, including talking with people you trust about your concerns and how you are feeling	53.1
	Taking breaks from watching, reading, or listening to news stories, including social media	52.6
	Taking care of your body, such as taking deep breaths, stretching, or meditating	46.4
	Engaging in healthy behaviors like trying to eat healthy, well-balanced meals, exercising regularly, getting plenty of sleep, or avoiding alcohol and drugs	45.1
	Eating more food than usual	26.7
	Eating less food than usual	19.7
	Using cannabis or marijuana	15.4
	Vaping more	8.5
	Drinking more alcohol	7.8
	Contacting a healthcare provider	5.4
	Using prescription drugs (like valium, etc.)	0.9

<b>Demographic Characteristics (Weighted %)</b>	
Using non-prescription drugs	0.9
Smoking more cigarettes	0.8
<b>Pandemic-Related Experiences (%)</b>	
Dropping any classes or other change to student status in Spring 2020	18.7
Major change to student status and educational plans in Fall 2020	46.4
Lacked a place to sleep or money for rent, gas, or food due to COVID-19	22.5
Significant impact on employment since March 2020	70.3

\* Demographic Factors may not add to 100% due to data missingness

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Depressive and Anxious Symptom Group Membership Predicted by Demographic Factors, Model Sample Size Range: 1,146-1,189

Table 2:

Relative Risk Ratios for Depressive Symptom Group Membership (ref = Not clinically high PHQ score) Predicted by Demographic Factors		Clinically High Symptoms Not/Only Somewhat Attributed to the Pandemic	Clinically High Symptoms More Than Somewhat Attributed to the Pandemic
Demographic	Group	RRR (95% CI)	RRR (95% CI)
Sex	Female (vs Male)	1.35 (0.92, 1.99)	<b>1.74 (1.08, 2.82)</b>
Race/Ethnicity	white (vs Black)	2.30 (0.95, 5.54)	1.96 (0.71, 5.42)
	Hispanic/Latino (vs Black)	2.16 (0.85, 5.50)	1.17 (0.38, 3.67)
	Asian (vs Black)	1.75 (0.58, 5.27)	1.41 (0.34, 5.92)
	Other (vs Black)	<b>2.86 (1.06, 7.74)</b>	<b>3.47 (1.02, 11.82)</b>
Parental Education	College or more (vs Some College or Less)	<b>0.67 (0.45, 0.98)</b>	1.25 (0.77, 2.04)
Urbanicity	A small or medium city (vs Rural)	0.99 (0.56, 1.73)	0.79 (0.39, 1.61)
	A suburb of a small or medium city (vs Rural)	0.67 (0.37, 1.21)	0.90 (0.42, 1.90)
	A large or very large city (vs Rural)	0.69 (0.37, 1.27)	0.61 (0.30, 1.26)
	A suburb of a large or very large city (vs Rural)	0.74 (0.35, 1.57)	0.46 (0.20, 1.08)
	Can't say, mixed (vs Rural)	1.68 (0.71, 3.96)	1.46 (0.48, 4.41)
Relative Risk Ratios for Anxious Symptom Group Membership (ref = Not clinically high GAD score) Predicted by Demographic Factors		Clinically High Symptoms Not/Only Somewhat Attributed to the Pandemic	Clinically High Symptoms More Than Somewhat Attributed to the Pandemic
Demographic	Group	RRR (95% CI)	RRR (95% CI)
Sex	Female (vs Male)	<b>2.11 (1.33, 3.33)</b>	<b>2.03 (1.16, 3.55)</b>
Race/Ethnicity	white (vs Black)	1.97 (0.72, 5.38)	<b>3.66 (1.22, 11.03)</b>
	Hispanic/Latino (vs Black)	1.51 (0.50, 4.59)	2.63 (0.82, 8.45)
	Asian (vs Black)	0.87 (0.23, 3.32)	2.96 (0.76, 11.59)
	Other (vs Black)	1.83 (0.55, 6.05)	<b>4.72 (1.31, 16.95)</b>
Parental Education	College or more (vs Some College or Less)	0.91 (0.58, 1.43)	0.87 (0.52, 1.48)
Urbanicity	A small or medium city (vs Rural)	1.18 (0.59, 2.34)	1.43 (0.67, 3.03)
	A suburb of a small or medium city (vs Rural)	1.19 (0.59, 2.41)	1.41 (0.58, 3.44)



Relative Risk Ratios for Depressive Symptom Group Membership (ref = Not clinically high PHQ score) Predicted by Demographic Factors		
	Clinically High Symptoms Not/Only Somewhat Attributed to the Pandemic	Clinically High Symptoms More Than Somewhat Attributed to the Pandemic
Demographic	RRR (95% CI)	RRR (95% CI)
Group		
A large or very large city (vs Rural)	0.66 (0.32, 1.38)	0.95 (0.44, 2.05)
A suburb of a large or very large city (vs Rural)	0.90 (0.37, 2.14)	0.97 (0.33, 2.85)
Can't say, mixed (vs Rural)	1.23 (0.53, 2.87)	1.77 (0.58, 5.43)

\* Bolding signifies p<.05

**Table 3:** Associations (aRRR\* and 95% CI's) for Pandemic-Related Experiences Predicting Depressive and Anxious Symptom Group Membership, Model Sample Size Range: 975\*\* -1,142

Associations (aRRR and 95% CI's) for Pandemic-Related Experiences Predicting Depressive Symptom Group Membership (ref = Not clinically high PHQ score)		Clinically High Symptoms Not/Only Somewhat Attributed to the Pandemic	Clinically High Symptoms More Than Somewhat Attributed to the Pandemic
<b>Pandemic-Related Experience</b>		aRRR (95% CI)	aRRR (95% CI)
Dropping any classes or other change to student status in Spring 2020		1.88 (1.00, 3.51)	2.57 (1.37, 4.81)
Major change to student status and educational plans in Fall 2020		1.39 (0.94, 2.06)	1.82 (1.14, 2.89)
Lacked a place to sleep or money for rent, gas, or food due to COVID-19		1.74 (1.11, 2.73)	4.43 (2.59, 7.55)
Significant impact on employment since March 2020		0.97 (0.64, 1.46)	2.02 (1.15, 3.52)
Associations (aRRR and 95% CI's) for Pandemic-Related Experiences Predicting Anxious Symptom Group Membership (ref = Not clinically high GAD score)		Clinically High Symptoms Not/Only Somewhat Attributed to the Pandemic	Clinically High Symptoms More Than Somewhat Attributed to the Pandemic
<b>Pandemic-Related Experience</b>		aRRR (95% CI)	aRRR (95% CI)
Dropping any classes or other change to student status in Spring 2020		2.07 (0.95, 4.50)	1.97 (0.98, 3.99)
Major change to student status and educational plans in Fall 2020		1.26 (0.80, 2.00)	1.81 (1.07, 3.07)
Lacked a place to sleep or money for rent, gas, or food due to COVID-19		1.85 (1.11, 3.10)	3.31 (1.92, 5.71)
Significant impact on employment since March 2020		1.50 (0.88, 2.53)	2.24 (1.27, 3.94)

\* Adjusted for sex, race/ethnicity, parental education, and urbanicity; Bolding signifies p<.05

\*\* Sample size for spring 2020 educational change models significantly reduced due to questionnaire structure, item was only asked to current students

**Table 4:**

Associations (aRRR\* and 95% CI's) for Coping Behaviors Predicting Depressive and Anxious Symptom Group Membership, Model Sample Size Range: 1,126-1,127

Associations (aRRR and 95% CI's) for Coping Behaviors Predicting Depressive Symptom Group Membership (ref = Not clinically high PHQ score)			
	Clinically High Symptoms Not/Only Somewhat Attributed to the Pandemic	Clinically High Symptoms More Than Somewhat Attributed to the Pandemic	aRRR (95% CI)
<b>Coping Behavior</b>			
Taking breaks from watching, reading, or listening to news stories, including social media	0.88 (0.59, 1.31)		<b>1.60 (1.00, 2.57)</b>
Taking care of your body, such as taking deep breaths, stretching, or meditating	0.71 (0.47, 1.06)		<b>0.57 (0.36, 0.92)</b>
Engaging in healthy behaviors like trying to eat healthy, well-balanced meals, exercising regularly, getting plenty of sleep, or avoiding alcohol and drugs	<b>0.60 (0.40, 0.89)</b>	0.76 (0.46, 1.26)	
Making time to relax	<b>0.55 (0.37, 0.82)</b>	0.94 (0.58, 1.52)	
Connecting with others, including talking with people you trust about your concerns and how you are feeling	0.69 (0.46, 1.03)	0.80 (0.50, 1.29)	
Contacting a healthcare provider	0.62 (0.24, 1.62)	<b>2.89 (1.39, 6.00)</b>	
Smoking more cigarettes	0.34 (0.04, 2.84)	0.91 (0.15, 5.51)	
Vaping more	<b>2.92 (1.41, 6.09)</b>	<b>4.36 (2.09, 9.12)</b>	
Drinking more alcohol	1.02 (0.55, 1.89)	<b>3.15 (1.60, 6.22)</b>	
Using prescription drugs (like valium, etc.)	2.98 (0.72, 12.39)	<b>5.72 (1.66, 19.68)</b>	
Using non-prescription drugs	0.29 (0.03, 2.85)	4.06 (0.92, 17.97)	
Using cannabis or marijuana	<b>2.43 (1.36, 4.33)</b>	<b>4.00 (2.26, 7.08)</b>	
Eating more food than usual	<b>1.87 (1.23, 2.85)</b>	<b>3.55 (2.20, 5.75)</b>	
Eating less food than usual	<b>2.35 (1.50, 3.68)</b>	<b>3.29 (1.98, 5.45)</b>	
<b>Associations (aRRR and 95% CI's) for Coping Behaviors Predicting Anxiety Symptom Group Membership (ref = Not clinically high GAD score)</b>			
	Clinically High Symptoms Not/Only Somewhat Attributed to the Pandemic	Clinically High Symptoms More Than Somewhat Attributed to the Pandemic	
<b>Coping Behavior</b>			
Taking breaks from watching, reading, or listening to news stories, including social media	1.34 (0.86, 2.10)	<b>2.40 (1.45, 3.97)</b>	
Taking care of your body, such as taking deep breaths, stretching, or meditating	1.04 (0.66, 1.66)	0.78 (0.45, 1.34)	

Associations (aRRR and 95% CIs) for Coping Behaviors Predicting Depressive Symptom Group Membership (ref = Not clinically high PHQ score)		
Coping Behavior	Clinically High Symptoms Not/Only Somewhat Attributed to the Pandemic aRRR (95% CI)	Clinically High Symptoms More Than Somewhat Attributed to the Pandemic aRRR (95% CI)
Engaging in healthy behaviors like trying to eat healthy, well-balanced meals, exercising regularly, getting plenty of sleep, or avoiding alcohol and drugs	0.73 (0.46, 1.14)	0.88 (0.53, 1.45)
Making time to relax	0.66 (0.42, 1.04)	0.91 (0.54, 1.52)
Connecting with others, including talking with people you trust about your concerns and how you are feeling	0.67 (0.42, 1.07)	1.01 (0.61, 1.70)
Contacting a healthcare provider	1.03 (0.47, 2.28)	<b>2.14 (1.03, 4.47)</b>
Smoking more cigarettes	0.60 (0.08, 4.61)	0.87 (0.15, 5.13)
Vaping more	<b>3.35 (1.51, 7.44)</b>	<b>4.76 (2.08, 10.93)</b>
Drinking more alcohol	1.36 (0.69, 2.69)	<b>3.19 (1.61, 6.30)</b>
Using prescription drugs (like valium, etc.)	0.60 (0.07, 5.32)	3.27 (0.93, 11.47)
Using non-prescription drugs	0.56 (0.07, 4.75)	2.58 (0.40, 16.73)
Using cannabis or marijuana	1.93 (1.00, 3.75)	<b>5.24 (2.81, 9.79)</b>
Eating more food than usual	<b>2.21 (1.36, 3.59)</b>	<b>2.27 (1.37, 3.76)</b>
Eating less food than usual	1.41 (0.87, 2.28)	<b>2.55 (1.43, 4.57)</b>

\* Adjusted for sex, race/ethnicity, parental education, and urbanicity; Bolding signifies p<.05