

Drug Poisoning Deaths in the United States, 1980–2008

Margaret Warner, Ph.D.; Li Hui Chen, Ph.D.; Diane M. Makuc, Dr.P.H.,
Robert N. Anderson, Ph.D.; and Arialdi M. Miniño, M.P.H.

Key findings

Data from the National Vital Statistics System Mortality File


- In 2008, poisoning became the leading cause of injury death in the United States and nearly 9 out of 10 poisoning deaths are caused by drugs.
- During the past three decades, the number of drug poisoning deaths increased sixfold from about 6,100 in 1980 to 36,500 in 2008.
- During the most recent decade, the number of drug poisoning deaths involving opioid analgesics more than tripled from about 4,000 in 1999 to 14,800 in 2008.
- Opioid analgesics were involved in more than 40% of all drug poisoning deaths in 2008, up from about 25% in 1999.
- In 2008, the drug poisoning death rate was higher for males, people aged 45–54 years, and non-Hispanic white and American Indian or Alaska Native persons than for females and those in other age and racial and ethnic groups.

In 2008, over 41,000 people died as a result of a poisoning. One of the Healthy People 2020 objectives, retained from *Healthy People 2010*, is to reduce fatal poisonings in the United States (1). However, poisoning mortality increased during the *Healthy People 2010* tracking period. Drugs—both legal and illegal—cause the vast majority of poisoning deaths. Misuse or abuse of prescription drugs, including opioid analgesic pain relievers, is responsible for much of the increase in drug poisoning deaths (see “Definitions” section). This report highlights trends in poisoning deaths, drug poisoning deaths, and the type of drugs involved in drug poisoning deaths and updates a previous data brief on this topic (2).

Keywords: opioid analgesics • overdose • prescription pain relievers • National Vital Statistics System Mortality File

Poisoning is now the leading cause of death from injuries in the United States and nearly 9 out of 10 poisoning deaths are caused by drugs.

Figure 1. Motor vehicle traffic, poisoning, and drug poisoning death rates: United States, 1980–2008



NOTE: In 1999, the *International Classification of Diseases, Tenth Revision (ICD-10)* replaced the previous revision of the ICD (ICD-9). This resulted in approximately 5% fewer deaths being classified as motor-vehicle traffic-related deaths and 2% more deaths being classified as poisoning-related deaths. Therefore, death rates for 1998 and earlier are not directly comparable with those computed after 1998. Access data table for Figure 1 at http://www.cdc.gov/nchs/data/databriefs/db81_tables.pdf#1.
SOURCE: CDC/NCHS, National Vital Statistics System.



In 2008, the number of poisoning deaths exceeded the number of motor vehicle traffic deaths for the first time since at least 1980. In 2008, there were more than 41,000 poisoning deaths, compared with about 38,000 motor vehicle traffic deaths. In 2008, 89% of poisoning deaths were caused by drugs.


During the past three decades, the poisoning death rate per 100,000 population nearly tripled from 4.8 in 1980 to 13.5 in 2008, while the motor vehicle traffic death rate decreased by almost one-half from 22.9 in 1980 to 12.5 in 2008 (Figure 1). In the most recent decade, from 1999 to 2008, the poisoning death rate increased 90%, while the motor vehicle traffic death rate decreased 15%.

From 1980 to 2008, the percentage of poisoning deaths caused by drugs increased from 56% to 89%. In 2008, about 77% of the drug poisoning deaths were unintentional, 13% were suicides, and 9% were of undetermined intent (see Appendix table).

Poisoning is the leading cause of death from injury in 30 states.

In 2008, poisoning was the leading cause of injury death in the following 30 states: Alaska, Arizona, California, Colorado, Connecticut, Delaware, Florida, Hawaii, Illinois, Indiana, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Nevada, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oregon, Pennsylvania, Rhode Island, Utah, Vermont, Washington, West Virginia, and Wisconsin (Figure 2).

Figure 2. Age-adjusted poisoning death rates: Comparison of state and U.S. rates: United States, 2008



NOTES: The poisoning death rate for Georgia may not be based on the final numbers of poisoning deaths. See "Data source and methods" for details. Figure 2 at http://www.cdc.gov/nchs/data/databriefs/db81_tables.pdf#2.
SOURCE: CDC/NCHS, National Vital Statistics System.


In 2008, age-adjusted poisoning death rates varied by state, ranging from 7.6 to 30.8 per 100,000 population. In 20 states, the age-adjusted poisoning death rate was significantly higher than the U.S. rate of 13.4 deaths per 100,000 population.

The five states with the highest poisoning death rates were New Mexico (30.8), West Virginia (27.6), Alaska (24.2), Nevada (21.0), and Utah (20.8). In 43 states over 80% of poisoning deaths were caused by drugs (data not shown).

Opioid analgesics were involved in more than 40% of drug poisoning deaths in 2008.

Of the 36,500 drug poisoning deaths in 2008, more than 40% (14,800) involved opioid analgesics (Figure 3). For about one-third (12,400) of the drug poisoning deaths, the type of drug(s) involved was specified on the death certificate but it was not an opioid analgesic. The remaining 25% involved drugs, but the type of drugs involved was not specified on the death certificate (for example, “drug overdose” or “multiple drug intoxication” was written on the death certificate).

Figure 3. Number of drug poisoning deaths involving opioid analgesics and other drugs: United States, 1999–2008



¹Opioid analgesics include natural and semi-synthetic opioid analgesics (for example, morphine, hydrocodone, and oxycodone) and synthetic opioid analgesics (for example, methadone and fentanyl). Some deaths in which the drug was poorly specified or unspecified may involve opioid analgesics.

NOTES: Drug categories are mutually exclusive. Access data table for Figure 3 at http://www.cdc.gov/nchs/data/databriefs/db81_tables.pdf#3.

SOURCE: CDC/NCHS, National Vital Statistics System.

From 1999 to 2008, the number of drug poisoning deaths involving opioid analgesics increased from about 4,000 to 14,800, more rapidly than deaths involving only other types of drugs or only nonspecified drugs.


From 1999 to 2008, the number of drug poisoning deaths involving only nonspecified drugs increased from about 3,600 to about 9,200. Some drug poisoning deaths for which the drug was not specified may involve opioid analgesics.

Natural and semi-synthetic opioid analgesics such as morphine, hydrocodone, and oxycodone were involved in over 9,100 drug poisoning deaths in 2008, up from about 2,700 in 1999.

Of the 14,800 drug poisoning deaths involving opioid analgesics in 2008, the majority involved natural and semi-synthetic opioid analgesics such as morphine, hydrocodone, and oxycodone. The number of drug poisoning deaths involving natural and semi-synthetic opioid analgesics increased steadily each year from about 2,700 deaths in 1999 to over 9,100 deaths in 2008 (Figure 4).

The number of drug poisoning deaths involving methadone, which is a synthetic opioid analgesic used to treat opioid dependency as well as pain, increased sevenfold from about 800 deaths in 1999 to about 5,500 in 2007. Between 2007 and 2008, the number of deaths involving methadone decreased by nearly 600 deaths, the first decrease since 1999.

Figure 4. Number of drug poisoning deaths involving opioid analgesic by opioid analgesic category: United States, 1999–2008




NOTES: Opioid analgesic categories are not mutually exclusive. Deaths involving more than one opioid analgesic category shown in this figure are counted multiple times. Natural and semi-synthetic opioid analgesics include morphine, hydrocodone, and oxycodone; and synthetic opioid analgesics include fentanyl. Access data table for Figure 4 at http://www.cdc.gov/nchs/data/databriefs/db81_tables.pdf#4. SOURCE: CDC/NCHS, National Vital Statistics System.

The number of drug poisoning deaths involving synthetic opioid analgesics other than methadone, such as fentanyl, tripled from about 700 in 1999 to 2,300 in 2008.

In 2008, the drug poisoning death rate was higher among those aged 45–54 years than among those in other age groups.

From 1999 to 2008, the drug poisoning death rate increased among all age groups. In 2004, the drug poisoning death rate among those aged 45–54 years surpassed the rate among those aged 35–44 years, and became the age group with the highest drug poisoning death rate (Figure 5).

Figure 5. Drug poisoning death rates by age: United States, 1999–2008



NOTE: Access data table for Figure 5 at http://www.cdc.gov/nchs/data/databriefs/db81_tables.pdf#5. SOURCE: CDC/NCHS, National Vital Statistics System.

From 1999 to 2008, the age-adjusted drug poisoning death rate increased for males and females and for all race and ethnicity groups (Appendix table). In 2008, the rate was higher for males than

for females, and higher for non-Hispanic American Indian or Alaska Native and non-Hispanic white persons than for those in other race and ethnicity groups.

Appendix table. Age-adjusted drug poisoning death rates, by demographic characteristics and intent: United States, 1999–2008

	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Age-adjusted death rate										
Total	6.1	6.2	6.8	8.1	8.9	9.3	10.0	11.4	11.8	11.9
Sex										
Male	8.2	8.3	9.0	10.5	11.4	11.7	12.7	14.6	14.8	14.8
Female	3.9	4.1	4.7	5.8	6.4	6.9	7.3	8.2	8.9	9.0
Race and ethnicity										
Hispanic	5.5	4.6	4.4	5.3	5.6	5.2	5.7	6.2	5.9	5.9
Non-Hispanic white	6.1	6.6	7.4	9.2	10.1	10.9	11.7	13.5	14.4	14.7
Non-Hispanic Black	7.5	7.3	7.6	8.2	8.2	8.3	9.4	10.9	9.8	8.5
Non-Hispanic American Indian or Alaska Native	6.0	5.5	6.8	8.4	10.6	12.3	12.9	13.9	13.9	15.6
Non-Hispanic Asian or Pacific Islander	1.2	1.0	1.2	1.4	1.4	1.5	1.7	1.9	1.9	1.8
Intent¹										
Unintentional	4.0	4.2	4.6	5.7	6.3	6.8	7.5	8.8	9.1	9.2
Suicide	1.1	1.2	1.2	1.3	1.3	1.4	1.4	1.5	1.5	1.6
Undetermined	0.9	0.9	1.0	1.1	1.2	1.1	1.0	1.1	1.2	1.1

¹Age-adjusted drug poisoning rates for homicides, legal interventions, and operations of war rates are less than 0.1 per 100,000 population each year and are not shown.

SOURCE: CDC/NCHS, National Vital Statistics System.

Summary

In 2008, the number of poisoning deaths exceeded the number of motor vehicle traffic deaths and was the leading cause of injury death for the first time since at least 1980. During the past three decades, the poisoning death rate nearly tripled, while the motor vehicle traffic death rate decreased by one-half. During this period, the percentage of poisoning deaths that were caused by drugs increased from about 60% to about 90%.

The population groups with the highest drug poisoning death rates in 2008 were males, people aged 45–54 years, and non-Hispanic white and American Indian or Alaska Native persons. The vast majority of drug poisoning deaths are unintentional (see [Appendix table](#)). Opioid analgesics were involved in more drug poisoning deaths than other specified drugs, including heroin and cocaine. Opioid analgesics were involved in nearly 15,000 deaths in 2008, while cocaine was involved in about 5,100 deaths and heroin was involved in about 3,000 deaths (data not shown). Deaths involving opioid analgesics may involve other drugs as well, including benzodiazepines (2).

In addition to an increase in the number of deaths caused by drug poisoning, increases in drug use, abuse, misuse, and nonfatal health outcomes have been observed. In the past two decades, there has been an increase in the distribution and medical use of prescription drugs, including opioid analgesics (3). From 1999 to 2008, the use of prescription medications increased (4). In 2007–2008, 48% of Americans used at least one prescription drug in the past month and 11% of Americans used five or more prescriptions in the past month. Analgesics for pain relief were

among the common drugs taken by adults aged 20–59 years (4). In 2009–2010, over 5 million Americans reported using prescription pain relievers nonmedically in the past month (that is, without a doctor’s prescription or only for the experience or feeling they caused), and the majority of people using prescription pain relievers nonmedically reported getting the drugs from friends or family (5,6). From 2004 to 2008, the estimated rate of emergency department visits involving nonmedical use of opioid analgesics doubled from 49 per 100,000 to 101 per 100,000 (7).

Government agencies and other organizations joined together to achieve great reductions in the number of deaths from motor vehicle crashes in the past three decades (8,9). A comprehensive approach, including improvements in the safety of vehicles; improvements in roadways; increased use of restraint systems, such as seat belts and child safety seats; reductions in speed; and also efforts to reduce driving under the influence of alcohol and drugs, contributed to the decline in motor vehicle related deaths (8,9). Using a comprehensive, multifaceted approach, it may be possible to reverse the trend in drug poisoning mortality.

Definitions

Injury deaths: Include deaths that are caused by forces external to the body. Examples of causes of injury death include drowning, fall, firearm, fire or burn, motor vehicle traffic, poisoning, and suffocation.

Poisoning deaths: Include drug poisonings resulting from unintentional or intentional overdoses of a drug, being given the wrong drug, taking the wrong drug in error, or taking a drug inadvertently. Poisoning deaths also include poisoning resulting from other toxic substances, gases, or vapors.

Opioid analgesics: Drugs that are usually prescribed to relieve pain and include: Natural and semi-synthetic opioid analgesics such as morphine, codeine, hydrocodone, and oxycodone; methadone, which is a synthetic opioid analgesic used to treat opioid dependency as well as pain; and other synthetic opioid analgesics (excluding methadone) such as fentanyl and propoxyphene. Opium and heroin are not included in this class of drugs.

Data source and methods

Estimates are based on the National Vital Statistics System multiple cause of death mortality files (10). Deaths were classified using the *International Classification of Diseases (ICD), Tenth Revision (ICD–10)* in 1999–2008 and the *Ninth Revision of the ICD (ICD–9)* in 1980–1998. Poisoning deaths were defined as having ICD–10 underlying cause of death code (UCOD): X40–X49, X60–X69, X85–X90, Y10–Y19, Y35.2, or *U01(.6–.7) and ICD–9 UCOD: E850.0–E869.9, E950.0–E952.9, E962(.0–.9), E972, or E980.0–E982.9. Drug poisoning deaths were defined as having ICD–10 UCOD: X40–X44 (unintentional), X60–X64 (suicide), X85 (homicide), Y10–Y14 (undetermined intent) and ICD–9 UCOD: E850–E858, E950.0–E950.5, E962.0, or E980.0–E980.5. Motor vehicle traffic deaths were defined as having ICD–10 UCOD: V02–V04(.1,.9), V09.2, V12–V14(.3–.9), V19(.4–.6), V20–V28(.3–.9), V29–V79(.4–.9), V80(.3–.5), V81–V82(.1), V83–V86(.0–.3), V87(.0–.8), or V89.2 and ICD–9 UCOD: E810.0–E819.9, E958.5, or E988.5. When the ICD–10 replaced ICD–9 in 1999, approximately 5% fewer deaths were classified as motor vehicle deaths and 2% more deaths were classified as poisoning deaths (11).

The identification of individual drugs and drug classes involved in drug poisoning deaths is limited by the classification structure of the ICD. Trends involving individual drugs and drug classes begin with 1999 when the ICD–10 replaced ICD–9, because the classification of individual drugs and drug classes is not comparable between ICD revisions. Among deaths with drug poisoning as the underlying cause, the following ICD–10 codes indicate the type of drug(s) involved: only nonspecified drug(s) (only T50.9); specified drug(s) other than opioid analgesic (any of the codes T36–T50.8 other than T40.2–T40.4); and any opioid analgesic (any of the codes T40.2–T40.4); and natural and semi–synthetic opioid analgesic (T40.2); methadone (T40.3); synthetic opioid analgesic, excluding methadone (T40.4); heroin (T40.1); and cocaine (T40.5).

Age-adjusted death rates were calculated using the direct method and the 2000 standard population (10). To identify state rates that were significantly higher or lower than the overall U.S. rate, differences between national and state estimates were evaluated using two-sided significance tests at the 0.01 level. Georgia was excluded from this comparison because the cause of death was inconclusive for a high proportion of deaths in Georgia in the 2008 NVSS mortality file. When the national mortality file was closed to updates, the manner of death was pending for 8.8% of deaths and was assigned an ill-defined cause in 3.5% of deaths for Georgia as compared with 0.5% pending and 0.3% ill-defined for the nation. Poisoning deaths, which require lengthy investigations, are typically among the causes that remain pending at the close of the file.

Several factors related to death investigation and reporting may affect measurement of death rates involving specific drugs. At autopsy, toxicological lab tests may be performed to determine the type of legal and illegal drugs present. The substances tested for and circumstances in which the tests are performed vary by jurisdiction. Measurement errors related to these factors are more likely to affect substance specific death rates than the overall drug poisoning death rate.

About the authors

Margaret Warner, Li Hui Chen, and Diane M. Makuc are with the Centers for Disease Control and Prevention’s (CDC) National Center for Health Statistics (NCHS), Office of Analysis and Epidemiology. Arialdi M. Miniño and Robert N. Anderson are with CDC, NCHS, Division of Vital Statistics.

References

1. Healthy People 2020 Topics and Objectives. Available from: <http://www.healthypeople.gov/2020/>. Accessed September 30, 2011.
2. Warner M, Chen LH, Makuc DM. Increase in fatal poisonings involving opioid analgesics in the United States, 1999–2006. NCHS data brief, no 22. Hyattsville, MD: National Center for Health Statistics. 2009. Available from: <http://www.cdc.gov/nchs/data/databriefs/db22.htm>.
3. Paulozzi L, Mack K, Rudd R, Jones C. Vital signs: Overdoses of prescription opioid pain relievers—United States, 1999–2008. MMWR, vol 60, 1487–92. 2011. Available from: <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6043a4.htm>.
4. Gu Q, Dilon CF, Burt VL. Prescription drug use continues to increase: U.S. prescription drug data for 2007–2008. NCHS data brief, no 42. Hyattsville, MD: National Center for Health Statistics. 2009. Available from: <http://www.cdc.gov/nchs/data/databriefs/db42.htm>.

**U.S. DEPARTMENT OF
HEALTH & HUMAN SERVICES**

Centers for Disease Control and Prevention
National Center for Health Statistics
3311 Toledo Road
Hyattsville, MD 20782

FIRST CLASS MAIL
POSTAGE & FEES PAID
CDC/NCHS
PERMIT NO. G-284

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE, \$300

NCHS Data Brief ■ No. 81 ■ December 2011

5. Substance Abuse and Mental Health Services Administration. The NSDUH Report: Trends in nonmedical use of prescription pain relievers: 2002 to 2007. Rockville, MD. 2009. Available from: <http://oas.samhsa.gov/2k9/painRelievers/nonmedicalTrends.pdf>.
6. Substance Abuse and Mental Health Services Administration. Results from the 2010 National Survey on Drug Use and Health: Summary of National Findings, NSDUH Series H-41, HHS Publication No. (SMA) 11-4658. Rockville, MD. 2011. Available from: <http://oas.samhsa.gov/NSDUH/2k10NSDUH/2k10Results.htm>.
7. Cai R, Crane E, Poneleit K, Paulozzi L. Emergency department visits involving nonmedical use of selected prescription drugs: United States, 2004–2008 (Reprinted from MMWR, vol 59, 705–9. 2010). JAMA-J Am Med Assoc 304(5):514–6. 2010. Available from: <http://jama.ama-assn.org/content/304/5/514.full>.
8. CDC. Motor-vehicle safety: A 20th century public health achievement (Reprinted from MMWR, vol 48, 369. 1999). JAMA-J Am Med Assoc 281(22):2080–82. 1999. Available from: <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm4818a1.htm>.
9. Longthorne A, Subramanian R, Chen CL. An analysis of the significant decline in motor vehicle traffic fatalities in 2008. DOT HS 811 346. Washington, DC 2010. Available from: <http://www-nrd.nhtsa.dot.gov/Pubs/811346.pdf>.
10. Miniño AM, Murphy SL, Xu JQ, Kochanek KD. Deaths: Final data for 2008. National vital statistics reports; vol 59, no 10. Hyattsville, MD: National Center for Health Statistics. 2011. http://www.cdc.gov/nchs/data/nvsr/nvsr59/nvsr59_10.pdf.
11. Bergen G, Chen L, Warner M, Fingerhut L. Injury in the United States: 2007 Chartbook. Hyattsville, MD: National Center for Health Statistics. 2008. Available from: <http://www.cdc.gov/nchs/data/misc/injury2007.pdf>.

Suggested citation

Warner M, Chen LH, Makuc DM, Anderson RN, Miniño AM. Drug poisoning deaths in the United States, 1980–2008. NCHS data brief, no 81. Hyattsville, MD: National Center for Health Statistics. 2011.

Copyright information

All material appearing in this report is in the public domain and may be reproduced or copied without permission; citation as to source, however, is appreciated.

National Center for Health Statistics

Edward J. Sondik, Ph.D., *Director*
Jennifer H. Madans, Ph.D., *Associate
Director for Science*

Office of Analysis and Epidemiology

Diane M. Makuc, Dr.P.H., *Acting Director*

For e-mail updates on NCHS publication releases, subscribe online at: <http://www.cdc.gov/nchs/govdelivery.htm>.

For questions or general information about NCHS: Tel: 1-800-232-4636
E-mail: cdcinfo@cdc.gov
Internet: <http://www.cdc.gov/nchs>

ISSN 1941-4927 (Print ed.)
ISSN 1941-4935 (Online ed.)

CS228727
DHHS Publication No. (PHS) 2012-1209