

Supplementary Table S3. Weighted use of each clinical preventive services among Medicaid expansion and non-expansion states by income level from 2011-2013

Expansion states*Year	Colon cancer screening		Breast cancer screening		Cervical cancer screening		HIV testing		Influenza vaccination	
	Lower-income	Higher-income	Lower-income	Higher-income	Lower-income	Higher-income	Lower-income	Higher-income	Lower-income	Higher-income
Expansion*2011	NA	NA	NA	NA	NA	NA	Reference	Reference	Reference	Reference
Expansion*2012	Reference	Reference	Reference	Reference	Reference	Reference	-0.001	-0.05	0.06	-0.01
(95% CI)							(-0.14 - 0.14)	(-0.13 - 0.02)	(-0.08 - 0.21)	(-0.08 - 0.06)
Expansion*2013	-0.23	0.07	-0.19	0.55**	-0.27	0.52**	-0.03	-0.01	0.0543	0.04
(95% CI)	(-0.52 - 0.07)	(-0.07 - 0.21)	(-0.81 - 0.44)	(0.18 - 0.93)	(-0.72 - 0.18)	(0.21 - 0.82)	(-0.17 - 0.11)	(-0.08 - 0.07)	(-0.09 - 0.20)	(-0.03 - 0.11)

\*\*p<0.01.

Abbreviations: HIV=human immunodeficiency virus; 95% CI=95% confidence interval.

Note: Lower-income group: household income<\$20,000; Higher-income group: household income ≥ \$20,000. Adjusted by age, sex, race/ethnicity, education level, marital status and health status. For sex-related services (breast cancer screening and cervical cancer screening), sex was not included for adjusting.