

RESEARCH ARTICLE

Open Access

Isolation and genomic characterization of Sfl, a serotype-converting bacteriophage of *Shigella flexneri*

Qiangzheng Sun^{1†}, Ruiting Lan^{2†}, Yiting Wang^{1†}, Jianping Wang^{1†}, Yan Wang¹, Peijing Li¹, Pengcheng Du¹ and Jianguo Xu^{1*}

Abstract

Background: All *Shigella flexneri* serotypes except serotype 6 share a common O-antigen tetrasaccharide backbone and nearly all variations between serotypes are due to glucosyl and/or O-acetyl modifications of the common O unit mediated by glycosyltransferases encoded by serotype-converting bacteriophages. Several *S. flexneri* serotype-converting phages including SfV, SfX, Sf6 and SflI have been isolated and characterized. However, *S. flexneri* serotype-converting phage Sfl which encodes a type I modification of serotype 1 (1a, 1b, 1c and 1d) had not yet been characterized.

Results: The Sfl phage was induced and purified from a *S. flexneri* serotype 1a clinical strain 019. Electron microscopy showed that the Sfl phage has a hexagonal head and a long contractile tail, characteristic of the members of *Myoviridae* family. Sfl can convert serotype Y to serotype 1a and serotype X to serotype 1d, but cannot convert 10 other *S. flexneri* serotypes (1a, 1b, 2a, 2b, 3a, 3b, 4a, 4b, 5a, Xv) tested, suggesting that Sfl has a narrow host range. Similar to other *S. flexneri* serotype-converting phages, Sfl integrates into the *tRNA-thrW* gene adjacent to *proA* of the host chromosome when lysogenized. The complete sequence of the Sfl genome was 38,389 bp, encoding 66 open reading frames and two tRNA genes. Phage Sfl shares significant homology with *S. flexneri* phage SfV, *Escherichia coli* prophage e14 and lambda, and is classified into the lambdoid phage family. Sfl was found to use a *cos* mechanism for DNA packaging similar to that of phage SfV.

Conclusions: Sfl contains features of lambdoid phages and is closely related to *S. flexneri* phage SfV, *E. coli* prophage e14 and lambda. The characterization of Sfl enhances our understanding of serotype conversion of *S. flexneri*.

Background

Shigella is the major cause of endemic bacillary dysentery (shigellosis) in developing countries. It is estimated that there are about 164.7 million cases of shigellosis annually worldwide, of which 163.2 million were in developing countries, resulting in 1.1 million deaths, most of which were children under 5 years of age [1]. Among the four *Shigella* species, *S. dysenteriae*, *S. flexneri*, *S. boydii*, and *S. sonnei*, *S. flexneri* is the predominant species.

Based on the combination of antigenic determinants present in the O-antigen of the cell envelope

lipopolysaccharide (LPS), *S. flexneri* is further divided into various serotypes. To date, at least 16 serotypes have been recognized [2-4]. Except for serotype 6, all share a basic repeating tetrasaccharide unit, comprised of one GlcNAc and three rhamnosides [4]. Modifications to the side chain of the tetrasaccharide by the addition of glucosyl and/or O-acetyl groups give rise to various antigenic determinants [3]. The genes responsible for the O-antigen modification are always either the gene cluster *gtrABC* for glucosyl groups or the single *oac* gene for the O-acetyl group; all encoded by serotype-converting bacteriophages [3,5-10]. In all glucosylation modification phages, the *gtrABC* gene cluster is always located immediately upstream of the *attP* site, followed by the *int* and *xis* genes [6].

Up to now, four *S. flexneri* serotype-converting bacteriophages, SfV, SfX, Sf6 and SflI, have been induced

* Correspondence: xujianguo@icdc.cn

†Equal contributors

¹State Key Laboratory for Infectious Disease Prevention and Control, National Institute for Communicable Disease Control and Prevention, China CDC, P.O. Box 5, Beijing, Changping, China

Full list of author information is available at the end of the article

and purified by different groups [8,11-13]. Morphologically, SfV and SflI, which have an isometric head and a long tail, belong to Group A in the family of *Myoviridae* [8,11]; while SfX and Sf6, which possess a short tail linked to an isometric head, belong to the family of *Podoviridae* [12,13]. The complete genome sequences of phage SfV and Sf6 have been obtained by directly sequencing the phage DNA purified from phage particles, and their genetic features have been well characterized [9,10]. Recently, the prophage genome of SfX was determined from the sequenced *S. flexneri* serotype Xv strain 2002017; which is presumably the whole genome of phage SfX, because a SfX phage particle can be induced and isolated from 2002017 [2]. The SfX genome is 37,355 bp length, encoding 59 ORFs (unpublished data). The genome of SflI has not yet been sequenced from free phage particles, but prophage genomes can be derived from sequenced *S. flexneri* serotype 2a strains Sf301 and 2457T [14,15], which show considerable variation with one or both being prophage remnants.

S. flexneri serotype 1 is defined by reaction with type I antisera. A total of 4 subtypes, 1a, 1b, 1c and 1d have been recognized [16-18]. In serotype 1, a glucosyl group is attached to the GlcNac residue of the repeating unit by an alpha-1, 4 linkage, which results in the presence of serotype 1-specific I antigen. The type I modification is mediated by an O-antigen glucosylation locus (*gtrI*, *gtrA*, *gtrB*) encoded on the SflI prophage genome [5]. The glucosylation genes and flanking partial SflI sequences were previously obtained from a serotype 1a strain Y53 [17]. However, the free phage particle of SflI had not been isolated, and its full genomic characteristics have not yet been elucidated [5].

In this study, we induced and purified the free SflI phage particles from *S. flexneri* serotype 1a clinical strain 019 and characterized its morphology, host range and genomic features.

Results and discussion

Isolation of phage SflI from *S. flexneri* serotype 1a strain 019

Using the conditions described in Methods, we induced the SflI phage from serotype 1a strain 019. Plaques were observed on the semi-solid LB agar when the host strain 036 was infected with induced products from strain 019. Lysogens isolated from plaques were serologically identified as serotype 1a, characterized by agglutination with both typing sera I and grouping sera 3;4. PCR amplification indicated that the SflI specific gene *gtrI* is present on both phage particles and the lysogens. These results suggest that phage SflI has been successfully induced and isolated from strain 019. This is the first report of isolation of free SflI particles from *S. flexneri*.

The morphology of SflI is characteristic of the *Myoviridae* family

The purified SflI phage particles were morphologically analyzed using electron microscopy. The phage has a hexagonal head of ca. 55 nm in diameter, a knob-like neck, a contractile tail of ca. 110 nm, and a tail sheath of ca. 55 nm (Figure 1). There are indications of a baseplate-like structure and long tail fibers, but no other distinctive features could be seen (Figure 1). These characteristics suggest that phage SflI is a member of the *Myoviridae* family in the order *Caudovirales* [19].

In comparison to other morphologically characterized serotype-converting phages Sf6, SfV, SflI and SfX, SflI has a very similar appearance to SflI and SfV [8,11], but distinctive from SfX and Sf6 [12,20]. The microscopic difference reflected the genetic divergence among them in that the SflI packaging and structure genes were identical to those of phage SfV, but divergent from those of SfX and Sf6 (see below, Figure 2).

Phage SflI has a very narrow host range

Host specificity of serotype-converting bacteriophages has long been recognized, which results in the specific lytic spectrum and serotype conversion of *S. flexneri* in nature [20]. The recognition between the O-antigen of host bacterium and the tail component of a phage is the key mechanism of host specificity [20]. To determine the host range of SflI, 132 *S. flexneri* strains of 12 serotypes (1a, 1b, 2a, 2b, 3a, 3b, 4a, 4b, 5a, Y, X and Xv) were tested following the methods described in the Methods. Apart from 10 serotype Y strains, which were all converted to serotype 1a as expected, the 24 serotype X strains tested





Figure 1 Electron micrograph of *S. flexneri* bacteriophage SflI stained with phosphotungstic acid.



were also lysogenized, and converted to a newly named serotype 1d [16]. The serotype 1d strains were serologically characterized as reacting with both serotype 1 specific I typing sera and serotype X specific 7;8 grouping sera [16]. Interestingly, such a serotype has already appeared in natural infections in Anhui and Henan provinces, China [21]. Except for serotypes Y and X, the other serotypes could not be lysogenized by phage Sfl. A possible explanation for the host range restriction of phage Sfl is phage immunity due to modification of the O-antigen as phage receptors [22].

Sfl uses a site-specific mechanism for DNA packaging and has the same *attP* core sequence as SflI, SflV, SflV and SflX Restriction enzyme analysis revealed that phage Sfl has a linear but not circular genome (data not shown). Genomic comparison found that the Sfl prophage genome has similar packaging genes to that of phage SflV; and the fragments adjacent to them were also highly similar to the cohesive end site (*cos*) of phage SflV [9], with only one base difference at the 5' end (T versus A). These data suggest that Sfl may use the same site-specific mechanism as SflV for packaging. Direct sequencing of the putative termini of the Sfl genome extracted from free phage particles and

comparison of the corresponding regions with the Sfl prophage genome in strain 019 revealed a 10 nucleotide (5'-TGCCCGCCCC-3') gap in the Sfl phage genome. Therefore, we conclude that Sfl uses a *cos* mechanism for DNA packaging as postulated for phage SflV [9], and does not use a head full mechanism (*pac*) as for phage Sfl6 and SflX [10,12].

Integration of lambdoid phages into the bacterial chromosome generally occurs by site-specific recombination between the phage *attP* and the bacterial *attB* sites [23]. In all serotype-converting phages except for Sfl6, the *attP* site is always found located immediately downstream of the O-antigen modification genes, and preceded by the *int* and *xis* genes [6]. To determine the *attP* site of phage Sfl, the region between genes *gtrA* and *intI* of Sfl was PCR amplified and sequenced and a 261 bp sequence was obtained, in which, 46 bases, ATTCGTAATGCGAAGGTCGTAGGTTCTCGACTCCTAT TATCGGCACCA, were found to be identical to the *attR/attL* core sequence of prophage Sfl in strain Y53 [5] (Figure 3). In the lysogen of 036_1a, the 261 nucleotide sequence was divided into two parts, located at opposite ends of the Sfl prophage genome (Figure 3). Evidently, site-specific recombination occurred at this *attP*

site. The *attP* core sequence of Sfl is identical to that of *S. flexneri* serotype-converting phage SfII, SfV and SfX, as well as that of serotype-converting phages p22 of *Salmonella typhimurium* and DLP12 of *E. coli* [5,8,24].

Characterization of Sfl genome sequence

The complete genome sequence of Sfl was obtained by combining the Sfl prophage genome of host strain 019 with the *attP* site obtained by PCR sequencing as above. Firstly, the whole genome sequence of host strain 019 was sequenced using Illumina Solexa sequencing. A total of 4,382,674 reads were generated to reach about 110-fold coverage and assembled *de novo* into 376 contigs and scaffolds. The Sfl prophage genome located between genes *int* and *gtrIA* was extracted from one of the contigs which was further assembled with the *attP* site sequence obtained above to construct a circular phage Sfl genome. To revert to the linear organization as usual practice, we artificially linearised the sequence starting from the terminase small subunit gene and ending with the *cos* site (Figure 2).

The genome size of Sfl is 38,389 bp similar to that of sequenced *S. flexneri* serotype-converting phages Sf6 (39,043 bp) [9], SfV (37,074 bp) [10] and SfX (37,355

(unpublished data). The overall G + C content is 50.12%, which is very similar to that of its host (50.9%) [25]. Sixty-six putative ORFs (including one pseudogene) were predicted and their functions are listed in the Additional file 1: Table S1.

The genetic architecture of the Sfl genome is similar to that of sequenced *S. flexneri* serotype-converting phages SfV, Sf6 and SfX: the left-most region encodes genes for phage packaging and structure, followed by the middle region with genes involved in serotype conversion, integration/excision, recombination, immunity and regulation, replication and the Nin region, and then the putative lysis cassette at the right-most region ending with the *cos* site of the phage genome (Figure 2). The genomic structure of Sfl is also similar to that of phage SfV and lambda. Thus it belongs to the family of lambdaoid phages.

tRNAscan was used to find tRNA genes. Two tRNA genes in tandem, with anticodons GUU for asparagine (Asn) and UGU for threonine (Thr), were found to be located downstream of gene Q (35,738 – 35,809 for Asn, and 35,818 – 35,890 for Thr). One or both of these tRNA genes were also to be found located at this position in phage Sf6, ST64T, PS3 and p21 [10,26,27]. A recent study



Figure 3 DNA sequences of chromosomal integration site of *S. flexneri* phage Sfl. Sequences obtained by PCR and sequencing of junction regions using a series of primers across the integration site. (A) *attP* in phage Sfl. (B) *attB* in strain 036. (C) *attL* in strain 036_1a. (D) *attR* in 036_1a. Sequences in box are DNA regions between conserved genes; Underlined sequences are *tRNA-thrW*; Sequences in blue are *att* core sequence; Conserved genes are shaded and their transcription orientation is marked by an arrow.

suggested that phage-encoded tRNA could serve to supplement the host tRNA reservoir, allowing the rare codons in the phage to be more efficiently decoded [28]. Codon analysis indeed found a convincing bias of ACA (anticodon UGU) in the Sfl genome when compared to its *S. flexneri* host (with 17.3% in phage Sfl, and 7.1% in strain Sf301), but no obvious bias was observed on CAA (anticodon GUU), and the significance of the *tRNA-Asn* in Sfl is not clear.

Genomic comparison reveals that Sfl is genetically related to *Shigella* phage SfV, *E. coli* prophage e14 and lambda

The ORFs encoded in the Sfl genome were searched against the GenBank database at both DNA and amino acid levels. Sfl encoded proteins exhibited homology to various phages and prophages originating from various hosts, including *Shigella* (SfV, Sf6 and SfX), *E. coli* (lambda, phip27, VT1-sakai, BP-4795, 933 W, 1717, 2851, Stx1, Stx2, VT2-Sa, YYZ-2008, 86, M27 and e14) and *Salmonella* (ST64B, p22-pbi, SE1, ST104, ST64T and epsilon34). Figure 2 displays the homologies of phage Sfl to other phages. The Sfl genes involved in phage packaging and morphogenesis are homologous and organized in a similar manner to those of phage SfV, phi-p27, ST64B and prophage e14. As reported earlier [6], the O- antigen modification and integration and excision modules (*gtrA*, *gtrB*, *int* and *xis*) are homologous to that of serotype-converting bacteriophages from *S. flexneri* (SfV and SfX) and *Salmonella* (p22-pbi, SE1, ST104, ST64T and epsilon34). However, the early and regulatory regions located in the right half of the genome were homologous to that of lambda and Shiga toxin-1 and Shiga toxin-2 phages (phip27, VT1-sakai, BP-4795, 933 W, 1717, 2851, Stx1, Stx2, VT2-Sa, YYZ-2008, 86 and M27).

Therefore Sfl is a mosaic phage with its left half most homologous to phage SfV (91.6% - 100% identity at protein level, and 89-98% at DNA level [ORF by ORF comparison]) and *E. coli* prophage e14 (94.0% - 100% identity at protein level, and 97% at DNA level) and right half most homologous to Lambda (67% - 100% identity at protein level, and 80 - 98% at DNA level). Homology to SfV encompasses at least 23 ORFs encoding functions for morphology (*orf1*, *orf2* and *orf9* - *orf24*), O-antigen modification (*orf26*, *orf27*), integration/excision (*orf28* to *orf29*) (Figure 2, Table 1). The homologous ORFs are located in four contiguous regions, amounting to 17,487 bp nucleotides and accounting for 45.6% of the entire phage genome (Table 1). Sfl also shared genetic relatedness with the *E. coli* prophage e14. The homologous regions mainly encode proteins responsible for phage assembly and morphogenesis and are located in the left half of the Sfl genome (Figure 2 and Table 1). The homologous regions account for 46% of the Sfl genome. Based on the homology of the first 22 ORFs (Additional file 2: Figure S1), it

seems that Sfl is closer to e14 than to SfV since 5 ORFs (Sfl *orf3* to *orf7*) are highly homologous between Sfl and e14, but share little homology between Sfl and SfV. For the remaining 17 ORFs except *orf8*, the pairwise percentage identities are very similar between Sfl, SfV and e14. On the other hand, the homology between Sfl and SfV extends further to *orf28* with high homology of *orf23*, *orf24* and *orf26* to *orf28*. Similarly, six contiguous DNA segments, which account for 28.4% of the Sfl genome, were found to be homologous to the corresponding regions of lambda. These homologous regions are mainly located in the early and regulatory regions, and encode functional modules for phage recombination (*orf35* to *orf43*), immunity and regulation (*orf45* to *orf50*), replication (*orf51*, *orf52*), Nin region (*orf53* to *orf55*, *orf57* to *orf60*), and part of the lysis module (*orf64*) (Figure 2 and Table 1). Thus a total of 72.9% of the Sfl genome is homologous to either SfV, e14 or lambda.

Conclusions

The serotype-converting bacteriophage Sfl was isolated from a *S. flexneri* serotype 1a strain. It had a narrow lytic pattern and converted only serotype Y to serotype 1a and serotype X to serotype 1d. Morphologically Sfl is a member of the *Myoviridae* family in the order of *Caudovirales*. Genomic analysis revealed that Sfl contains features of lambdaoid phages and is closely related to *S. flexneri* phage SfV, *E. coli* prophage e14 and lambda. The characterization of serotype-converting phage Sfl enhances our understanding of serotype conversion of *S. flexneri*.

Methods

Bacterial strains, media and culture

S. flexneri serotype 1a strain 019 [16] was used as the source for induction of phage Sfl. *S. flexneri* strain 036 (serotype Y) was used as the host for phage infection and large volume propagation of Sfl [16]. One hundred and thirty two *S. flexneri* strains of 12 serotypes (17 serotype 1a, 5 serotype 1b, 10 serotype 2a, 10 serotype 2b, 10 serotype 3a, 2 serotype 3b, 5 serotype 4a, 5 serotype 4b, 4 serotype 5a, 10 serotype Y, 24 serotype X and 30 serotype Xv) were used for phage host range detection. All *S. flexneri* strains used in this study were isolated from diarrheal patients in China, or purchased from National Collection of Type Cultures (NCTC), UK. *S. flexneri* strains were serologically identified using *Shigella* antisera Kits (Denka Seiken, Japan) and monoclonal antibody reagents (Reagensia AB, Sweden). *S. flexneri* strains were routinely cultured on LB agar or in LB broth with shaking at 37°C.

Induction of phage Sfl

Induction of phage Sfl was performed as methods described by Mavris *et al.* [8]. Briefly, a freshly grown colony of strain 019 was incubated in 10 ml LB broth

Table 1 Homology of Sfl to *S. flexneri* phage SflV and *E. coli* prophage e14 and lambda

Phage or prophage	Nucleotide position	Homologous nucleotide position in Sfl (total length [bp])	% identity at nucleotide level	Sfl ORFs ^a	% of Sfl genome
SflV	9 – 2,211	2 – 2,194 (2,193)	98	<i>orf1</i> , (<i>orf2</i>)	45.6
	5,793 – 17,782	6,053 – 18,042 (11,990)	97	<i>orf9</i> - <i>orf24</i>	
	19,146 – 22,042	19,787 – 22,681 (2,895)	98	(<i>orf26</i>), <i>orf2</i> - <i>orf29</i> , <i>attP</i>	
	36,666 – 37,074	37,964 – 38,372 (409)	89	(<i>orf66</i>)	
Lambda	30,418 – 30,910	23,002 – 23,493 (491)	95	(<i>orf31</i>), <i>orf32</i> , (<i>orf33</i>)	28.4
	31,206 – 34,381	24,281 – 27,456 (3,176)	98	(<i>orf35</i>), <i>orf36</i> - <i>orf43</i>	
	35,104 – 35,386	27,708 – 27,990 (283)	98	(<i>orf45</i>)	
	35,496 – 41,084	28,052 – 33,640 (5,590)	98	<i>orf46</i> - <i>orf55</i>	
	42,097 – 43,068	2 – 2,194 (2,193)	97	<i>orf57</i> - <i>orf59</i> , (<i>orf60</i>)	
	45,966 – 46,361	6,053 - 18,042 (11,990)	80	(<i>orf64</i>)	
e14	2,840,259 - 2,859,298 ^b	1 - 17,234, 36,721 - 38,389 (17,660)	97	<i>orf1</i> - <i>orf22</i> , (<i>orf66</i>)	46%

^a Parentheses indicate that the region of homology starts or ends within an ORF.

^b *E. coli* S88 strain genome (accession no. CU928161).

overnight with vigorous shaking. After being induced for 30 min at 56°C with aeration, the cultures were centrifuged, and the supernatants were filtered through a 0.22 mm membrane filter (Promega) to remove bacterial cells. The filtrates were either used directly for phage infection assay or stored at 4°C with addition of 10% (v/v) chloroform.

Phage infection and lysogenization

S. flexneri strain 036 cells were prepared using the methods for phage lambda [29]. Phage infection and lysogenization were performed using the methods described previously [16]. The serotypes of isolated colonies were identified by slide agglutination assay. Large volume phage purification was performed on *S. flexneri* strain 036, according to the methods for phage SflII [8].

Electron microscopy

The purified phages were absorbed on carbon-coated copper grids (300 mesh) and negatively stained with 2% (w/v) sodium phosphotungstate (pH 7.0). Samples were visualized with a Hitachi 600 electron microscope at 80 kV.

Host range detection

To determine the host range of phage Sfl, one hundred and thirty two *S. flexneri* strains of 12 serotypes were infected with Sfl. The preparation of component cells, phage infection and lysogen isolation were performed as methods for strain 036 above. The Sfl host range was determined by observing the presence of plaques and serologically identification of the lysogens.

Identification of the chromosomal integration site and cohesive ends (*cos* sites) of phage Sfl

Oligonucleotide primers *gtrI*-F (5'- ATTGAACGCCTCC TTGCTATGC -3'), *intI*-R (5'- AGTGTTCAGGAA

ATGGGAGGC -3'), *proA*-F (5'- ACAAAGCGAAATCA TCCTCAA -3'), and *yaiC*-R (5'- GCAGGAAACCACC ATCAACACC -3'), which are complementary to the genes *gtrI* and *intI* in phage Sfl, and *proA* and *yaiC* in *S. flexneri* chromosome, respectively, were used to identify the *attP* and *attB* sites of phage Sfl and strain 036, as well as the *attR* and *attL* regions of the Sfl lysogen. PCR was conducted using the Sensoquest labcycler PCR System (SENSO, German) under standard protocol. The PCR products were either cloned into TA vector pMD20-T (TaKaRa) for sequencing or sequenced directly.

To determine the cohesive ends of the Sfl phage, two primers, *cos*-F: 5'- ATGCCACCACGAACCCCAAAAAG -3' (nt 37,964 - 37,985, complementary to Sfl genome sequence), *cos*-R: 5'- GGCTTGGGGCGACGCCCGGA -3' (nt 72-91, complementary to Sfl genome), were designed to sequence the putative termini of the Sfl genome directly using phage DNA as the template. The phage genome ends obtained were further compared to the corresponding regions of the Sfl prophage genome in strain 019. The missing region in the former sequence is the putative *cos* site of phage Sfl.

Genome sequencing and analysis

To obtain the entire phage genome sequence of Sfl, the whole genome of source strain 019 was sequenced by Illumina Solexa sequencing. A paired-end (PE) library with an average insertion length of between 500 bp and 2,000 bp was constructed. Reads were generated with Illumina Solexa GA Iix (Illumina, San Diego, CA) and assembled into scaffolds using SOAP denovo (Release1.04). The sequence between genes *intI* and *gtrA* was extracted for further analysis. By assembling with the sequence amplified from Sfl DNA using primer pair *gtrI*-F and *intI*-R mentioned above, the entire sequence of Sfl genome in its circular state was obtained. Open reading frames (ORFs) of

SfI were determined using the ORF Finder program, which is accessible through the National Center for Biotechnology Information (NCBI). Searches for homologous DNA and protein sequences were conducted with the BLAST software against the non-redundant GenBank database (<http://www.ncbi.nlm.nih.gov/blast/blast/>). tRNA genes were determined with tRNAscan-SE Search server (<http://lowelab.ucsc.edu/tRNAscan-SE>).

Nucleotide accession number

The genomic sequence of phage SfI has been deposited in GenBank as accession number JX509734.

Additional files

Additional file 1: Table S1. Analysis of predicted ORFs and proteins of SfI.

Additional file 2: Figure S1. Gene by gene comparison of homologous regions of SfI with *S. flexneri* phage SfV and *E. coli* prophage e14. The arrows indicate the predicted proteins and orientation of the ORFs. The regions marked with a lightly red rectangle represent >50% sequence identity at amino acid level.

Competing interests

The authors declare that they have no competing interest.

Authors' contributions

JX, QS and RL designed the study, and co-drafted the manuscript. YW participated in the induction of the phage. JW carried out the PCR amplification and DNA sequencing. PL participated in the phage induction and infection. YW and PD participated in the sequence alignment and genome annotation. All authors read and approved the final manuscript.

Acknowledgements

This work was supported by grants from the National Natural Science Foundation of China (No. 81271788), the National Basic Research Priorities Program (2011CB504901), the Project of State Key Laboratory for Infectious Disease Prevention and Control (2011SKLID203, 2008SKLID106), the National Key Program for Infectious Diseases of China (2013ZX10004221, 2013ZX10004216-001-002) and the Special Project of Beijing Educational Committee (YB20098450101).

Author details

¹State Key Laboratory for Infectious Disease Prevention and Control, National Institute for Communicable Disease Control and Prevention, China CDC, P.O. Box 5, Beijing, Changping, China. ²School of Biotechnology and Biomolecular Sciences, University of New South Wales, Sydney, NSW 2052, Australia.

Received: 24 August 2012 Accepted: 31 January 2013

Published: 17 February 2013

References

1. Kotloff KL, Winickoff JP, Ivanoff B, Clemens JD, Swerdlow DL, Sansonetti PJ, Adak GK, Levine MM: **Global burden of *Shigella* infections: implications for vaccine development and implementation of control strategies.** *Bull World Health Organ* 1999, **77**(8):651–666.
2. Ye C, Lan R, Xia S, Zhang J, Sun Q, Zhang S, Jing H, Wang L, Li Z, Zhou Z, *et al*: **Emergence of a new multidrug-resistant serotype X variant in an epidemic clone of *Shigella flexneri*.** *J Clin Microbiol* 2010, **48**(2):419–426.
3. Stagg RM, Tang SS, Carlin NI, Talukder KA, Cam PD, Verma NK: **A novel glucosyltransferase involved in O-antigen modification of *Shigella flexneri* serotype 1c.** *J Bacteriol* 2009, **191**(21):6612–6617.
4. Simmons DA, Romanowska E: **Structure and biology of *Shigella flexneri* O antigens.** *J Med Microbiol* 1987, **23**(4):289–302.
5. Adhikari P, Allison G, Whittle B, Verma NK: **Serotype 1a O-antigen modification: molecular characterization of the genes involved and their novel organization in the *Shigella flexneri* chromosome.** *J Bacteriol* 1999, **181**(15):4711–4718.
6. Allison GE, Verma NK: **Serotype-converting bacteriophages and O-antigen modification in *Shigella flexneri*.** *Trends Microbiol* 2000, **8**(1):17–23.
7. Adams MM, Allison GE, Verma NK: **Type IV O antigen modification genes in the genome of *Shigella flexneri* NCTC 8296.** *Microbiology* 2001, **147**(Pt 4):851–860.
8. Mavris M, Manning PA, Morona R: **Mechanism of bacteriophage SfII-mediated serotype conversion in *Shigella flexneri*.** *Mol Microbiol* 1997, **26**(5):939–950.
9. Allison GE, Angeles D, Tran-Dinh N, Verma NK: **Complete genomic sequence of SfV, a serotype-converting temperate bacteriophage of *Shigella flexneri*.** *J Bacteriol* 2002, **184**(7):1974–1987.
10. Casjens S, Winn-Stapley DA, Gilcrease EB, Morona R, Kuhlewein C, Chua JE, Manning PA, Inwood W, Clark AJ: **The chromosome of *Shigella flexneri* bacteriophage Sf6: complete nucleotide sequence, genetic mosaicism, and DNA packaging.** *J Mol Biol* 2004, **339**(2):379–394.
11. Allison GE, Angeles DC, Huan P, Verma NK: **Morphology of temperate bacteriophage SfV and characterisation of the DNA packaging and capsid genes: the structural genes evolved from two different phage families.** *Virology* 2003, **308**(1):114–127.
12. Guan S, Bastin DA, Verma NK: **Functional analysis of the O antigen glucosylation gene cluster of *Shigella flexneri* bacteriophage SfX.** *Microbiology* 1999, **145**(5):1263–1273.
13. Gernski P Jr, Koeltzow DE, Formal SB: **Phage conversion of *Shigella flexneri* group antigens.** *Infect Immun* 1975, **11**(4):685–691.
14. Wei J, Goldberg MB, Burland V, Venkatesan MM, Deng W, Fournier G, Mayhew GF, Plunkett G 3rd, Rose DJ, Darling A, *et al*: **Complete genome sequence and comparative genomics of *Shigella flexneri* serotype 2a strain 2457T.** *Infect Immun* 2003, **71**(5):2775–2786.
15. Jin Q, Yuan Z, Xu J, Wang Y, Shen Y, Lu W, Wang J, Liu H, Yang J, Yang F, *et al*: **Genome sequence of *Shigella flexneri* 2a: insights into pathogenicity through comparison with genomes of *Escherichia coli* K12 and O157.** *Nucleic Acids Res* 2002, **30**(20):4432–4441.
16. Sun Q, Lan R, Wang Y, Wang J, Luo X, Zhang S, Li P, Ye C, Jing H, Xu J: **Genesis of a novel *Shigella flexneri* serotype by sequential infection of serotype-converting bacteriophages SfX and SfI.** *BMC Microbiol* 2011, **11**:269.
17. Bastin DA, Lord A, Verma NK: **Cloning and analysis of the glucosyl transferase gene encoding type I antigen in *Shigella flexneri*.** *FEMS Microbiol Lett* 1997, **156**(1):133–139.
18. Stagg RM, Cam PD, Verma NK: **Identification of newly recognized serotype 1c as the most prevalent *Shigella flexneri* serotype in northern rural Vietnam.** *Epidemiol Infect* 2008, **136**(8):1134–1140.
19. Ackermann HW: **Tailed bacteriophages: the order caudovirales.** *Adv Virus Res* 1998, **51**:135–201.
20. Lindberg AA, Wollin R, Gernski P, Wohlhieter JA: **Interaction between bacteriophage Sf6 and *Shigella flexneri*.** *J Virol* 1978, **27**(1):38–44.
21. Luo X, Sun Q, Lan R, Wang J, Li Z, Xia S, Zhang J, Wang Y, Jin D, Yuan X, *et al*: **Emergence of a novel *Shigella flexneri* serotype 1d in China.** *Diagn Microbiol Infect Dis* 2012, **74**(3):316–319.
22. Lindberg AA (Ed): *Bacterial surface polysaccharides and phage adsorption.* New York: Academic; 1977.
23. Campbell AM: **Chromosomal insertion sites for phages and plasmids.** *J Bacteriol* 1992, **174**(23):7495–7499.
24. Huan PT, Bastin DA, Whittle BL, Lindberg AA, Verma NK: **Molecular characterization of the genes involved in O-antigen modification, attachment, integration and excision in *Shigella flexneri* bacteriophage SfV.** *Gene* 1997, **195**(2):217–227.
25. Blattner FR, Plunkett G 3rd, Bloch CA, Perna NT, Burland V, Riley M, Collado-Vides J, Glasner JD, Rode CK, Mayhew GF, *et al*: **The complete genome sequence of *Escherichia coli* K-12.** *Science* 1997, **277**(5331):1453–1462.
26. Mmolawa PT, Schmieger H, Tucker CP, Heuzenroeder MW: **Genomic structure of the *Salmonella enterica* serovar Typhimurium DT 64 bacteriophage ST64T: evidence for modular genetic architecture.** *J Bacteriol* 2003, **185**(11):3473–3475.
27. Guo HC, Kainz M, Roberts JW: **Characterization of the late-gene regulatory region of phage 21.** *J Bacteriol* 1991, **173**(4):1554–1560.

28. Schmidt H, Scheef J, Janetzki-Mittmann C, Datz M, Karch H: **An *ileX* tRNA gene is located close to the Shiga toxin II operon in enterohemorrhagic *Escherichia coli* O157 and non-O157 strains.** *FEMS Microbiol Lett* 1997, **149**(1):39–44.
29. Sambrook J, Fritsch EF, Maniatis T: *Molecular Cloning: a Laboratory Manual*, 2nd edn. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory; 1989.

doi:10.1186/1471-2180-13-39

Cite this article as: Sun *et al.*: Isolation and genomic characterization of Sfi, a serotype-converting bacteriophage of *Shigella flexneri*. *BMC Microbiology* 2013 **13**:39.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

