

Respiratory Illnesses



Respiratory Illnesses

[Respiratory Illnesses Home](#)

Respiratory Virus Activity Levels

Provides an update on how COVID-19, influenza, and RSV may be spreading nationally and in your state.

Activity Levels Update:


- The amount of respiratory illness (fever plus cough or sore throat) causing people to seek healthcare is elevated across most areas of the country, though some decrease in activity is noted. This week, 37 jurisdictions experienced high or very high activity compared to 39 jurisdictions last week.
- Nationally, emergency department visits with diagnosed influenza, COVID-19, and RSV remain elevated in many areas. The decreases seen across all three conditions in the past week might be affected by holiday-related healthcare seeking behavior and will be monitored.
- National COVID-19 test positivity remained stable at 12.7%. For influenza, the percentage of specimens testing positive in clinical labs decreased nationally to 14.0%. National RSV positivity remained stable at 9.9%.
- Nationally, COVID-19 [wastewater viral activity levels](#) are very high. All regions show high and increasing levels, with the highest regional levels in the South and Midwest. There are early indications that wastewater activity levels are slowing their rate of increase in the Midwest and Northeast.
- After several weeks of increases in influenza indicators, a single week of decrease in most indicators have been noted. CDC will continue to monitor for a second period of increased influenza activity that often occurs after the winter holidays.

Reported on Friday, January 12th, 2024.

Level of Respiratory Illness Activity


Activity levels determined weekly based on the percentage of visits to enrolled outpatient healthcare providers or emergency departments for fever and cough or sore throat reported to ILINet. Visits can be attributed to a variety of respiratory pathogens that cause these symptoms. Activity levels reflect how the percentage in the most recent week compares to what that jurisdiction typically experiences during low circulation periods. Trend information for the percentages used to calculate activity levels can be found at: [National, Regional, and State Level Outpatient Illness and Viral Surveillance \(cdc.gov\)](#).





Territories PR VI

Respiratory Illness Activity Level



[Dataset on data.cdc.gov](#) | [Link to Dataset](#)

Data Table

Location	Respiratory Illness Level	Respiratory Illness Level
Alabama	Level 12	Very High
Alaska	Level 5	Low
Arizona	Level 10	High
Arkansas	Level 10	High
California	Level 11	Very High
Colorado	Level 11	Very High
Connecticut	Level 9	High
Delaware	Level 6	Moderate

● District Of Columbia	Level 10	High
● Florida	Level 10	High
● Georgia	Level 12	Very High
● Hawaii	Level 5	Low
● Idaho	Level 10	High
● Illinois	Level 7	Moderate
● Indiana	Level 9	High
● Iowa	Level 6	Moderate
● Kansas	Level 7	Moderate
● Kentucky	Level 10	High
● Louisiana	Level 13	Very High
● Maine	Level 7	Moderate
● Maryland	Level 10	High
● Massachusetts	Level 10	High
● Michigan	Level 8	High
● Minnesota	Level 1	Minimal
● Mississippi	Level 11	Very High
● Missouri	Level 6	Moderate
● Montana	Level 8	High
● Nebraska	Level 10	High
● Nevada	Level 9	High
● New Hampshire	Level 9	High
● New Jersey	Level 11	Very High
● New Mexico	Level 13	Very High
● New York	Level 8	High
● New York City	Level 12	Very High
● North Carolina	Level 11	Very High
● North Dakota	Level 10	High
● Ohio	Level 10	High
● Oklahoma	Level 7	Moderate
● Oregon	Level 4	Low
● Pennsylvania	Level 9	High
● Puerto Rico	Level 2	Minimal
● Rhode Island	Level 8	High
● South Carolina	Level 13	Very High
● South Dakota	Level 5	Low

● Tennessee	Level 13	Very High
● Texas	Level 11	Very High
● U.S. Virgin Islands	Level 4	Low
● Utah	Level 5	Low
● Vermont	Level 4	Low
● Virginia	Level 10	High
● Washington	Level 8	High
● West Virginia	Level 4	Low
● Wisconsin	Level 8	High
● Wyoming	Level 11	Very High

Data table showing data for the United States Map figure.

Data Notes: Level of Respiratory Illness Activity



- **Source:** U.S. Outpatient Influenza-Like Illness Surveillance Network (ILINet).
- Additional information available at: [Outpatient Illness Surveillance methods section](#).
- This system monitors visits for respiratory illness that includes fever plus a cough or sore throat, (also referred to as influenza-like illness, or ILI), not laboratory confirmed infections; therefore, patient visits due to a variety of respiratory pathogens that cause similar symptoms may be captured.
- The activity levels compare the mean reported percent of visits due to ILI during the current week to the mean reported percent of visits due to ILI during non-influenza weeks. The 13 activity levels correspond to the number of standard deviations below, at, or above the mean for the current week compared with the mean during non-influenza weeks.
- This map uses the proportion of visits to enrolled outpatient healthcare providers or emergency departments for respiratory illness to measure the activity level within a state. It does not, however, measure the extent of geographic spread of respiratory illness within a state. Therefore, outbreaks occurring in a single city could cause the state to display high activity levels.
- Data collected in ILINet may disproportionately represent certain populations within a state depending on enrolled providers, and therefore may not accurately depict the full picture of respiratory virus activity for the whole state.
- The data presented in this map is preliminary and may change as more data is received.
- Differences in the data presented by CDC and state health departments likely represent differing levels of data completeness with data presented by the state likely being the more complete.

Emergency Department Visits for Viral Respiratory Illness

Weekly percent of total emergency department visits associated with COVID-19, influenza, and RSV.

State

United States

County


All

Health Service Area

Counties Represented

All

All



● Combined ● COVID-19 ● Influenza ● RSV

Data presented through: 01/06/2024; Data as of: 01/10/2024

Dataset on data.cdc.gov | Link to Dataset


Data Table				
Week Ending	Combined	COVID-19	Influenza	RSV
10/01/2022	2.9%	1.8%	0.5%	0.6%
10/08/2022	2.9%	1.7%	0.7%	0.6%
10/15/2022	3.3%	1.6%	0.9%	0.7%
10/22/2022	3.7%	1.6%	1.3%	0.9%
10/29/2022	5.1%	1.7%	2.5%	1.0%
11/05/2022	6.5%	1.7%	3.7%	1.2%
11/12/2022	6.7%	1.7%	3.9%	1.2%
11/19/2022	7.0%	1.8%	4.3%	1.1%
11/26/2022	9.8%	2.2%	6.7%	1.0%
12/03/2022	9.4%	2.6%	6.2%	0.8%
12/10/2022	9.0%	2.7%	5.7%	0.7%
12/17/2022	8.5%	2.9%	5.2%	0.5%
12/24/2022	8.2%	3.3%	4.6%	0.5%
12/31/2022	7.7%	3.8%	3.6%	0.4%
01/07/2023	5.6%	3.3%	2.0%	0.3%
01/14/2023	3.8%	2.6%	1.0%	0.2%
01/21/2023	3.2%	2.3%	0.7%	0.2%
01/28/2023	2.8%	2.1%	0.6%	0.2%
02/04/2023	2.6%	2.1%	0.5%	0.1%
02/11/2023	2.5%	2.0%	0.4%	0.1%
02/18/2023	2.3%	1.8%	0.4%	0.1%
02/25/2023	2.1%	1.7%	0.3%	0.1%
03/04/2023	1.9%	1.5%	0.3%	0.1%
03/11/2023	1.7%	1.3%	0.3%	0.1%
03/18/2023	1.6%	1.2%	0.3%	0.1%
03/25/2023	1.5%	1.2%	0.3%	0.1%
04/01/2023	1.4%	1.1%	0.3%	0.0%
04/08/2023	1.3%	1.0%	0.3%	0.0%
04/15/2023	1.2%	0.9%	0.3%	0.0%
04/22/2023	1.0%	0.8%	0.2%	0.0%
04/29/2023	1.0%	0.7%	0.3%	0.0%
05/06/2023	1.0%	0.7%	0.3%	0.0%
05/13/2023	0.9%	0.6%	0.3%	0.0%
05/20/2023	0.9%	0.6%	0.3%	0.0%
05/27/2023	0.9%	0.6%	0.3%	0.0%
06/03/2023	0.9%	0.6%	0.3%	0.0%
06/10/2023	0.8%	0.5%	0.3%	0.0%
06/17/2023	0.7%	0.5%	0.2%	0.0%

06/24/2023	0.7%	0.5%	0.2%	0.0%
07/01/2023	0.7%	0.5%	0.2%	0.0%
07/08/2023	0.8%	0.6%	0.2%	0.0%
07/15/2023	0.9%	0.7%	0.2%	0.0%
07/22/2023	1.0%	0.8%	0.2%	0.0%
07/29/2023	1.2%	1.0%	0.2%	0.0%
08/05/2023	1.5%	1.3%	0.2%	0.0%
08/12/2023	1.8%	1.6%	0.2%	0.0%
08/19/2023	2.2%	2.0%	0.2%	0.1%
08/26/2023	2.7%	2.4%	0.2%	0.1%
09/02/2023	2.7%	2.5%	0.2%	0.1%
09/09/2023	2.7%	2.4%	0.3%	0.1%
09/16/2023	2.4%	2.1%	0.3%	0.1%
09/23/2023	2.3%	1.9%	0.3%	0.1%
09/30/2023	2.2%	1.7%	0.3%	0.2%
10/07/2023	2.1%	1.5%	0.4%	0.2%
10/14/2023	2.1%	1.4%	0.5%	0.3%
10/21/2023	2.3%	1.4%	0.6%	0.4%
10/28/2023	2.5%	1.4%	0.7%	0.5%
11/04/2023	2.7%	1.3%	0.9%	0.6%
11/11/2023	3.4%	1.5%	1.2%	0.7%
11/18/2023	3.8%	1.6%	1.4%	0.8%
11/25/2023	4.5%	1.9%	1.7%	1.0%
12/02/2023	4.8%	2.0%	1.9%	1.0%
12/09/2023	5.4%	2.1%	2.5%	0.9%
12/16/2023	6.8%	2.3%	3.7%	0.9%
12/23/2023	8.7%	2.8%	5.2%	0.9%
12/30/2023	9.8%	3.3%	5.7%	0.9%
01/06/2024	7.4%	2.9%	4.0%	0.7%

Data Notes: Emergency Department Visits for Viral Respiratory Illness ^

Percent of Tests Positive for Respiratory Viruses

Weekly percent of tests positive for the viruses that cause COVID-19, influenza, and RSV at the national level. Preliminary data are shaded in gray.



● COVID-19 ● Influenza ● RSV

Data presented through: 01/06/2024; Data as of: 01/11/2024

[Dataset on data.cdc.gov](#) | [Link to Dataset](#)

Data Table			
Week Ending	COVID-19	Influenza	RSV
10/01/2022	6.9%	2.0%	10.2%
10/08/2022	6.9%	2.7%	12.5%
10/15/2022	7.1%	3.5%	14.6%
10/22/2022	6.8%	5.4%	16.5%
10/29/2022	6.9%	8.4%	18.3%
11/05/2022	6.5%	12.8%	19.1%
11/12/2022	6.8%	14.9%	19.5%
11/19/2022	6.7%	18.3%	16.7%

11/26/2022	7.6%	25.8%	13.4%
12/03/2022	8.6%	25.6%	10.5%
12/10/2022	8.7%	26.3%	7.9%
12/17/2022	9.4%	24.5%	6.7%
12/24/2022	9.8%	20.8%	5.6%
12/31/2022	10.6%	14.9%	5.1%
01/07/2023	10.3%	8.6%	4.9%
01/14/2023	8.8%	4.6%	4.0%
01/21/2023	8.5%	2.9%	3.3%
01/28/2023	8.3%	2.1%	2.5%
02/04/2023	8.0%	1.7%	2.2%
02/11/2023	8.0%	1.4%	1.8%
02/18/2023	8.1%	1.1%	1.6%
02/25/2023	7.9%	1.0%	1.4%
03/04/2023	7.4%	0.9%	1.2%
03/11/2023	6.8%	1.0%	1.1%
03/18/2023	6.8%	0.9%	0.9%
03/25/2023	6.9%	0.9%	0.8%
04/01/2023	6.5%	1.0%	0.7%
04/08/2023	6.2%	1.1%	0.6%
04/15/2023	5.7%	1.0%	0.6%
04/22/2023	5.0%	0.8%	0.5%
04/29/2023	4.9%	1.0%	0.5%
05/06/2023	5.0%	0.9%	0.5%
05/13/2023	4.8%	1.0%	0.5%
05/20/2023	4.6%	1.2%	0.5%
05/27/2023	4.6%	1.2%	0.4%
06/03/2023	4.2%	1.3%	0.5%
06/10/2023	4.1%	1.1%	0.5%
06/17/2023	4.3%	1.1%	0.5%
06/24/2023	4.4%	0.9%	0.6%
07/01/2023	4.9%	1.1%	0.6%
07/08/2023	5.7%	1.2%	0.5%
07/15/2023	6.5%	1.0%	0.6%
07/22/2023	7.4%	1.0%	0.8%
07/29/2023	8.9%	1.0%	0.8%

08/05/2023	10.5%	1.1%	0.9%
08/12/2023	12.2%	0.9%	0.8%
08/19/2023	13.3%	0.9%	0.8%
08/26/2023	14.5%	0.8%	1.2%
09/02/2023	14.1%	0.7%	1.1%
09/09/2023	13.8%	0.7%	1.3%
09/16/2023	12.6%	0.8%	1.4%
09/23/2023	12.0%	0.9%	2.1%
09/30/2023	10.9%	0.9%	2.9%
10/07/2023	10.0%	1.1%	3.7%
10/14/2023	9.1%	1.3%	4.8%
10/21/2023	9.4%	1.5%	6.0%
10/28/2023	9.5%	1.8%	7.7%
11/04/2023	9.2%	2.6%	9.2%
11/11/2023	10.0%	4.0%	10.7%
11/18/2023	9.5%	4.9%	12.0%
11/25/2023	10.9%	6.1%	13.0%
12/02/2023	11.7%	7.0%	11.9%
12/09/2023	11.6%	8.4%	12.1%
12/16/2023	12.0%	13.3%	12.2%
12/23/2023	12.4%	17.2%	11.2%
12/30/2023	12.8%	18.0%	10.5%
01/06/2024	12.7%	14.0%	9.9%

Data Notes: Percent of Tests Positive for Viral Respiratory Pathogens



- **Sources:** COVID-19 and RSV: National Respiratory and Enteric Virus Surveillance System (NREVSS), a sentinel network of laboratories located through the US, includes clinical, public health and commercial laboratories; additional information available at: <https://www.cdc.gov/surveillance/nrevss/index.html>. Influenza: Clinical laboratory test results from NREVSS and U.S. World Health Organization collaborating laboratories; more details about influenza virologic surveillance are available here: <https://www.cdc.gov/flu/weekly/overview.htm>.
- Data for recent weeks in gray may be incomplete due to delays in reporting. These preliminary may change as more data become available.
- Data represent laboratory tests performed, not individual people.
- The data are from across the country in all regions.
- The percent of tests positive is calculated by dividing the number of positive tests by the total number of tests administered, then multiplying by 100 [(# of positive tests/total tests) x 100].
- COVID-19: The condition caused by infection with severe acute respiratory syndromic coronavirus type-2 (SARS-CoV-2).
- RSV and COVID-19 are limited to nucleic acid amplification tests (NAATs), also listed as polymerase chain reaction tests (PCR).
- Participating laboratories report weekly to CDC the total number of RSV tests performed that week and the number of those tests that were positive. The RSV trend graphs display the national average of the weekly % test positivity for the current, previous, and following weeks in accordance with the recommendations for assessing RSV trends by percent (<https://academic.oup.com/jid/article/216/3/345/3860464>).
- COVID-19 laboratory data are available for download here: <https://data.cdc.gov/Laboratory-Surveillance/Percent-Positivity-of-COVID-19-Nucleic-Acid-Amplif/gvzb-yw6g>
- RSV laboratory data are available for download here: <https://data.cdc.gov/Laboratory-Surveillance/Percent-Positivity-of-Respiratory-Syncytial-Virus-/3cxc-4k8q>

Explore deeper data

[State Map of Outpatient Respiratory Illness Activity](#)

[Outpatient Illness and Viral Surveillance](#)

[State Map of Emergency Department Use for COVID-19](#)

[COVID-19 Testing, Hospitalization, and Death Trends](#)

[Weekly U.S. Influenza Surveillance Report](#)

[RSV Testing Trends in the U.S.](#)



PREVIOUS

[Weekly Viral Respiratory Illness Snapshot](#)

NEXT

[Severe Viral Respiratory Illness](#)

