

TABLE 9. Estimated percent using an inhaled corticosteroid in the past 3 months among children¹ with current² or active³ asthma status by state/territory - BRFSS Asthma Call-back Survey, United States, 2006-2010

State/Territory	Current				Active			
	Sample size [±]	%	SE [§]	95% CI [†]	Sample size	%	SE	95% CI
Arizona	125	25.9	6.0	(14.1 - 37.7)	149	23.4	5.3	(13.0 - 33.8)
California	170	28.0	4.4	(19.4 - 36.6)	206	23.9	3.6	(16.8 - 31.0)
Connecticut	347	26.8	3.0	(20.9 - 32.7)	413	23.2	2.6	(18.1 - 28.3)
District of Columbia	114	25.5	4.5	(16.7 - 34.3)	140	23.3	3.9	(15.7 - 30.9)
Georgia	341	24.6	3.2	(18.3 - 30.9)	399	22.0	2.8	(16.5 - 27.5)
Hawaii	417	20.0	2.6	(14.9 - 25.1)	444	19.8	2.4	(15.1 - 24.5)
Illinois	168	25.8	4.4	(17.2 - 34.4)	196	22.8	3.9	(15.2 - 30.4)
Indiana	348	30.2	3.2	(23.9 - 36.5)	391	25.8	2.9	(20.1 - 31.5)
Iowa	249	20.0	2.9	(14.3 - 25.7)	274	17.8	2.6	(12.7 - 22.9)
Kansas	588	32.1	2.3	(27.6 - 36.6)	657	29.3	2.1	(25.2 - 33.4)
Louisiana	107	19.0	4.1	(11.0 - 27.0)	122	17.1	3.7	(9.8 - 24.4)
Maine	284	34.8	3.3	(28.3 - 41.3)	332	30.1	3.0	(24.2 - 36.0)
Maryland	385	21.7	2.5	(16.8 - 26.6)	467	17.8	2.1	(13.7 - 21.9)
Massachusetts	269	31.7	4.1	(23.7 - 39.7)	315	27.6	3.7	(20.3 - 34.9)
Michigan	480	30.5	2.8	(25.0 - 36.0)	533	28.7	2.6	(23.6 - 33.8)
Mississippi	80	21.4	6.2	(9.2 - 33.6)	94	18.3	5.4	(7.7 - 28.9)
Missouri	230	31.6	4.4	(23.0 - 40.2)	255	28.3	4.2	(20.1 - 36.5)
Montana	210	28.4	3.9	(20.8 - 36.0)	253	26.0	3.5	(19.1 - 32.9)
Nebraska	467	30.2	3.7	(22.9 - 37.5)	528	27.8	3.4	(21.1 - 34.5)
New Hampshire	150	26.3	4.1	(18.3 - 34.3)	173	23.6	3.7	(16.3 - 30.9)
New Jersey	296	21.7	3.2	(15.4 - 28.0)	337	19.5	3.0	(13.6 - 25.4)
New Mexico	234	21.5	3.8	(14.1 - 28.9)	262	19.5	3.4	(12.8 - 26.2)
New York	276	23.3	3.3	(16.8 - 29.8)	336	19.2	2.7	(13.9 - 24.5)
North Dakota	98	25.4	5.4	(14.8 - 36.0)	120	19.6	4.6	(10.6 - 28.6)
Ohio	222	25.3	4.4	(16.7 - 33.9)	258	21.1	3.8	(13.7 - 28.5)
Oklahoma	250	32.1	3.5	(25.2 - 39.0)	256	31.7	3.5	(24.8 - 38.6)
Oregon	193	19.6	2.9	(13.9 - 25.3)	224	18.5	2.7	(13.2 - 23.8)
Pennsylvania	134	28.2	4.7	(19.0 - 37.4)	156	23.9	4.0	(16.1 - 31.7)
Rhode Island	153	32.6	4.2	(24.4 - 40.8)	180	30.2	3.8	(22.8 - 37.6)
Texas	479	27.6	3.3	(21.1 - 34.1)	558	23.2	2.9	(17.5 - 28.9)
Utah	368	30.6	3.8	(23.2 - 38.0)	414	28.9	3.5	(22.0 - 35.8)
Vermont	389	36.0	2.8	(30.5 - 41.5)	443	33.1	2.7	(27.8 - 38.4)
Virginia	88	35.7	8.7	(18.6 - 52.8)	100	33.7	8.1	(17.8 - 49.6)
Washington	341	23.7	2.9	(18.0 - 29.4)	389	22.8	2.9	(17.1 - 28.5)
West Virginia	193	22.2	3.4	(15.5 - 28.9)	213	20.1	3.1	(14.0 - 26.2)
Wisconsin	147	28.6	4.8	(19.2 - 38.0)	206	22.2	3.7	(14.9 - 29.5)
36 state total	9,390	26.8	1.0	(24.8 - 28.8)	10,793	23.4	0.9	(21.6 - 25.2)
Median		26.6				23.3		
Range		19.0-36.0				17.1-33.7		

¹ Aged < 18 years

² "Yes" response to "Do you still have asthma?"

³ Reported any of the following during the past year: asthma medication use, asthma symptoms, or MD visits for asthma

[±] Sample size is for multiple years of participation. Responses of "Don't Know" and "Refused" are excluded.

[§] Standard error

[†] Confidence interval