Morbidity and Mortality

Weekly Report

OPUBLICISHEALTH SERVICE

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE

CDC LIBRARY

ATLANTA 22, GA. Prepored by the

COMMUNICABLE DISEASE CENTER

MElrose 4-5131

For release October 27, 1961

Atlanta 22, Georgia

Vol. 10, No. 42

Provisional Information on Selected Notifiable Diseases in the United States and on Deaths in Selected Cities for Week Ended October 21, 1961

Poliomyelitis — There has been an increase in the number of poliomyelitis cases reported during the current week ending October 21. A total of 59 cases, 39 paralytic, was reported. This is in contrast to the low report of 35 cases, including 20 paralytic, for the previous week.

The 1961 incidence through the current week is compared below with the corresponding time period in the Past four years.

POLIO (CUMULATED WEEKLY) THROUGH THE 42ND WEEK FOR PAST FIVE YEARS

	1961	1960	1959	1958	1957
Paralytic	693	1,850	4,709	2,300	1,838
Total	1,080	2,680	7,204	4,744	5,385

The increase in reported cases during the current week is scattered throughout the nation with no evidence of localization. The paralytic case reported this week from South Carolina was the 21st case (17 paralytic) to occur in Newberry County. This case, a 32 year old white male, had onset on October 16 and died on October 20. On Thursday, October 19, 22,600 doses of Type III oral polio vaccine were administered throughout the county in a mass immunization program.

Only six States reported as many as three paralytic cases. Of the six, the cases were scattered geographically in New York, Missouri, West Virginia, Louisiana, and

Table I. Cases of Specified Notifiable Diseases: United States

(Cumulative totals include revised and delayed reports through previous week)

Disease		42nd Wee	≥k	Cumulative						
(Seventh Revision of International Lists, 1955)	Ended	Ended Oct. 22, 1960	Median 1956-60	F	irst 42 wee	ks	Since s	Approxi- mate seasonal		
 Weekly incidence low or sporadic Data not available Quantity zero 	Oct-			1961	1960	Median 1956-60	1960-61	1959-60	Median 1955-56 to 1959-60	low point
nthrax062	_	_	*	6	15	*	*	*	*	*
O40 1	_	-	*	5	10	*	*	*	*	*
"MCCLI Daig (undulant farem) O44 (3	20	15	484	648	653	*	*	*	*
	12	26	36	451	551	627	142	206	235	July 1
CDhalitta imeasticus 000	47	23	52	1,346	1,557	1,725	1,346	1,557	1,725	Jan. 1
TPG-LLUS. infections and				-,	-,	-,		i -,	-,5	
	1,191	944	309	60,761	31,259	15,980	7,943	5,641	1,960	Sept. 1
	1	1	*	46	59	*	*	*	*	*
	1,336	1,331	1,331	391,530	406,153	453,826	5,902	6,229	6,755	Sept. 1
	97	98		2.639	2,474		2,639	2,474		Jan. 1
	43	51	43	1,734	1,781	1,961	219	244	259	Sept. 1
	59	140	248	1,080	2,680	5,373	977	2,471	4,845	Apr. 1
	39	98	194	693	1,850	1,828	633	1,691	1,553	Apr. 1
Paralytic	14	21	42	256	560	2,676	231	535	2,515	Apr. 1
	6	21	1 12	131	270	869	113	245	777	Apr. 1
	_	2	*	54	82	*	*	*	*	*
treptococoni como Abaset	-	-	*	3	2	*	*	*	*	*
	4,752	4.016		258,816	250,141		39,605			Aug. 1
	19	28	28	657	678	861	543	548	681	Apr. 1
rever, endemic	1	-	*	36	55	*	*	*	*	*
ables in animals	57	43	67	2,833	2,888	3,590	199	117	190	Oct. 1

California. In Wisconsin, 3 of the 4 paralytic cases reported were from Milwaukee County. Investigations are underway.

Hepatitis - There were 1,191 cases of hepatitis reported for the current week ending October 21. This is 42 cases more than was reported the preceding week. The level of hepatitis cases has remained approximately the same since July.

EPIDEMIOLOGICAL REPORTS

Nicotinic Acid Poisoning - Pennsylvania

Two cases of nicotinic acid (niacin) poisoning involving 5 persons were recently investigated in Philadelphia, Pennsylvania. In both instances ground meats including beef and lamb were the foods implicated. Symptoms exhibited by all 5 persons were essentially the same – intense flushing of the skin, a feeling of warmth, itching, and some abdominal discomfort. The ill effects lasted from 45 minutes to an hour. Onset of symptoms occurred within 15 minutes following the ingestion of the meat. All affected individuals ate the ground meats twice and on both occasions became ill with the same syndrome. Samples of meat obtained at the home of the patients and the market where meats were purchased revealed niacin to be present in the following amounts.

Chopped lamb patties 87 mg/100 grams of meat Ground beef patties 112 mg/100 grams of meat Ground beef stuffed into

peppers 50 mg/100 grams of meat Ground beef 145 mg/100 grams of meat

The niacin had been added to the meat to assure retention of its color for longer periods of time, thereby increasing its shelf life. It is felt by the Philadelphia officials that the addition of niacin to ground meats is more universal than is generally suspected.

(Reported by Milton Werrin, V.M.D., Chief, Veterinary Public Health Section, Philadelphia Department of Public Health)

Hepatitis from A Chimpanzee - Connecticut

Another report has been received on human hepatitis cases possibly transmitted from a chimpanzee.

On August 8, the Director of the Children's Zoo in New London, Connecticut, purchased a chimpanzee from an importer in New York City and brought it to New London. The chimp was so young it was taken into the Director's home. He, his wife and their 16-year-old son diapered, nursed and cared for the animal.

On September 6 the wife became ill with a fever, anorexia and nausea. Five days later, she developed jaundice. On September 11, the son developed similar

symptoms, and both cases were diagnosed as infectious hepatitis. On September 15 the husband and an aunt living in the home received gamma globulin. Neither he nor the aunt developed symptoms.

Prior to the Director's purchase of the chimp, it had been "farmed out" to a pet shop in New York City for a two-week period since the animal required special care. Reportedly, the chimp originated from Liberia and arrived as a single chimpanzee in a shipment of animals. Specimens from the chimpanzee, the human cases, and the contacts are being tested.

Previous reports this year in MMWR on human hepatitis cases probably transmitted from chimpanzees have involved 8 laboratory workers in Miami, Florida (MMWR Vol. 10, No. 20) and 6 zoo workers in Miami, Florida (MMWR Vol. 10, No. 35).

A summary of human hepatitis in the United States probably transmitted from chimpanzees or monkeys appeared in CDC Hepatitis Surveillance Report No. 4 (March 21, 1961).

(Epidemic reported by Dr. Mila Rindge, M.P.H., Chief, Epidemiology Section, Connecticut State Department of Health and Dr. Robert McCollum, Yale University).

Bat Rabies - Massachusetts and South Dakota

Massachusetts recently discovered and reported its first case of rabies in bats. The bat was found in Harvard, Massachusetts, and was brought to the State laboratory on September 5, 1961, after having bitten a woman the previous evening. The bat apparently flew into the open bedroom window of a child visiting his grandparents. After discovering the bat in his bedroom the youngster drove the animal into a closet, slammed the door shut and injured he bat while doing so. The boy's mother hearing the commotion arose from her sleep to investigate. Upon opening the closet door she was bitten on the toe by the bat. The bat was carried by the husband to the State Diagnostic Laboratory the next morning.

Direct impression smears of the bat's brain and salivary glands stained for Negri bodies were negative. However, mouse inoculation studies were suspicious. Tissues from both the bat and the mouse brain were sent to the Communicable Disease Center in Atlanta and were found to be positive by fluorescent antibody techniques on October 9, 1961.

The description of the bat coincided most closely with that of the small brown Myotis lucifugus.

This represents the first case of animal rabies discovered in Massachusetts since 1949.

(Continued on page 8)

Table 2. CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES, EACH DIVISION AND STATE, AND PUERTO RICO, FOR WEEKS ENDED OCTOBER 22, 1960 AND OCTOBER 21, 1961

(By place of occurrence. Numbers under diseases are category numbers of the Seventh Revision of the International Lists, 1955)

				Po	liomyeli	tis 080					Menin-	Brucel losis
	(Include		otal t specified	by type)	Par	alytic C	80.0,080	.1	Nonparalytic		gitis, aseptic	(undu-
Area	42nd	Week		Cumulative, first 42 weeks		Week	Cumulative, first 42 weeks		080.2		340 pt.	fever)
	1961	1960	1961	1960	1961	1960	1961	1960	1961	1960	1961	1961
UNITED STATES	59	140	1,080	2,680	39	98	693	1,850	14	21	97	3
ŒW ENGLAND	-	8	27	207	-	7	19	162	4	1	6	-
Maine New Hampshire	-	4	3 1	38	-	4	3	38	-	-	-	0
Vermont	_		5	7	_	_	4	2	-	_	_	
Massachusetts	-	1	12	29	-	-	8	19	-	1	3	
Rhode Island	-	-	1	99	-	-	1	76	-	= -	3	-
Connecticut	-	3	5	34	П -	3	3	27	-	-	-	-
IDDLE ATLANTIC	10	22	282	391	8	10	186	275	-	6	2	-
New York	9 1	12 4	204 34	218 75	7	4 3	131	142	-	3	1	-
Pennsylvania	_	6	44	98	_	3	28	55 78		3	1	_
AST NORTH CENTRAL	16	26	135	453	10	20	83	273	3	5	31	
Ohio	5	4	41	106	2	4	19	55	1		1	
Indiana	4	9	16	102	2	6	9	75	1	2	1	-
Illinois	2	5 8	25 26	131 83	- 2	3	12	88	-	2	12	-
Michigan	5	-	27	31	4	7 -	22	46 9	1	1	11	-
EST NORTH CENTRAL	4									}	6	-
Minnesota	-	2 1	63 6	154 49	3 -	1	28	87 41		1	8	- 5
Iowa	-		18	21	_		و	4	_		6	- 0
Missouri	4	-10	20	32	3	_	6	22	-	-	1	
North Dakota	-	-	3	14	-	-	-	5	-	Ψ.	-	-
South Dakota	-	- 1	1 8	4 15			4	1 8	-	*	-	-
Kansas	_	- 1	7	19	_	_	3	6	1 12 -	1	1	
OUTH ATLANTIC	14	41	185	484	7	34	135	380	7	5	10	
Delaware			2	-		-	1	300	12		10	-
Maryland	1	18	28	126	1	14	27	115	1200	4	-	
District of Columbia		7	3 10	2	1	-	3	2		-		11112
West Virginia	5	6	29	31 46	3	7 6	8	28 38	2	121	8	-
North Carolina	4	4	21	80	1	4	11	60	3	1 4	-	_
South Carolina	1	1	32	123	1	1	24	81		-	: in:	-
Georgia	- 3	5	28	18	-	_	21	16	-		-	-
CERTAMENTO			32	58	1	2	21	40	2	1	2	-
Kentucky	-	17 10	74 27	210 117	_	7	42	85 5			2	
Tennessee	-	1	17	40	_	1	8	27	_			
Alabama	-	6	9	21	_	6	9	21	-	-	2	1
Mississippi	-	-	21	32	-		20	32	-	- 1	-	-
EST SOUTH CENTRAL	5	6	139	254	5	5	75	157		1	- 5	1
Arkansas	4	1	17 49	28		-	7	20		-	-	-
Louisiana	-	_	49	47 16	4	1.0	39	28 11		1	- 1	will
Texas	1	5	69	163	1	5	29	98	-		4	1
OUNTAIN	2	5	44	73	1	3	25	38	1		270	5 = 32
Montana	1	2	4	18	_	2	2	14	î	11 11-		
Idaho	-	2	14	7		-	6	1	-	10.7-	-	-
WyomingColorado	1	1	7	19 15	- 1	1	7	1 14	-	-	-	-
New Mexico	-] [3	5	-			14 2		21		-
Arizona	-	-	8	4	-	-	6	4		-	- 1	
Nevada	-	-	8	5	1 2	-	4	2	-	-	-	-
Nevada	-	-	- 121	-	-	-	-	-	-	-	-	-
ACIFIC	8 1	13 1	131 23	454 31	5 1	11 1	100 15	393 31	3	2	33	2
Oregon	2		16	32	1	-	8	18	1	-	1	
California	5	12	88	382	3	10	73	335	2	2	32	2
Alaska	-	-	1 1 -	2	-	-	1	2	-	-		-
Hawaii	-	73.	4	7	-		4	7	-		-	
uerto Rico	_	3	6	472	-	3	6	465				100 mm

Table 2. CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES, EACH DIVISION AND STATE, AND PUERTO RICO, FOR WEEKS ENDED OCTOBER 22, 1960 AND OCTOBER 21, 1961 - Continued

(By place of occurrence. Numbers under diseases are category numbers of the Seventh Revision of the International Lists, 1955)


		Diphthe	ria 055		Enceph infec	alitis, tious		titis, i rum 092	Measles			
Area	42nd	42nd Week		Cumulative, first 42 weeks		2	42nd	42nd Week		ative, 2 weeks	085	
	1961	1960	1961	1960	1961	1960	1961	1960	1961	1960	1961	1960
UNITED STATES	12	26	451	551	47	23	1,191	944	60,761	31,259	1,336	1,331
NEW ENGLAND	-	-	7	11	-	-	68	18	1,902	913	178	152
Maine New Hampshire	_	_	_	2 -	_	_	11 13	3 -	129 170	56 27	84 2	1 3
Vermont	-	-	-	-	-	-	2	1	184	13	-	1 2
MassachusettsRhode Island	_	_	6	8 1	_	-	31 3	7 2	773 216	453 179	64	130 3
Connecticut	-	-	1	-	-	-	8	5	430	185	28	15
MIDDLE ATLANTIC	-	-	20	13	13	5	158	183	8,282	3,805	133	265
New York		_	7 -	3 2	11	4	83	123	3,529	2,098	73	126
Pennsylvania	-	-	13	8	2	-	39 36	7 53	1,962 2,791	259 1,448	13 47	56 83
EAST NORTH CENTRAL	-	-	15	38	5	-	226	147	12,073	5,588	304	322
OhioIndiana	_	-	1 1	16 5	2	-	66	45	4,027	1,899	17	88
Illinois	_	-	10	6	2	_	22 56	14 52	1,812 2,171	631 1,205	37 153	38 29
MichiganWisconsin	- -	-	3	9	-	-	72	32	3,755	1,651	64	61
WEST NORTH CENTRAL	-	4	40	2 36	1 1	- /	10	4	308	202	33	106
Minnesota	-	3	24	17	-	4	81 25	68 24	5,812 1,281	2,217 320	43	22
Iowa	-	_	2	7	-	2	21	2	1,708	358	17	5
North Dakota	-	_	1 4	2 1	-	_	21	18 1	1,299 126	779 162	2 18	13
South Dakota	-	1	7	6	-	-	7	4	178	138	3	
Nebraska Kansas	_	_	2	1 2	- 1	- 2	3 4	10 9	584 636	228	3707	2
SOUTH ATLANTIC	4	11	105	178	5	5	174	93	7,661	232 3,641	NN 69	NN 85
Delaware	-	-	-	-	-	_	1	10	175	218	_	9
Maryland District of Columbia	_	-	1 2	1	-	2	13	10	681	363	31	9
Virginia	_	2	15	29	3	- 2	3 30	3	113 1,231	42 674	3	13
West Virginia	-	-	1	4	-	-	17	13	1,385	675	12	18
North Carolina	2 1	2	9 10	10 46	1 -	-	60 9	13 8	1,711 403	332 112	5	1
Georgia	-	5	31	33	-	-	ģ	4	684	239	6	5
Florida	1	2	36	55	1	1	32	32	1,278	986	7	30
EAST SOUTH CENTRAL	1 -	4	37 9	56 2	3	2 2	167 52	116	9,143	4,441	95	71
Tennessee	-	-	3	7	1	-	53	47 33	2,698 3,576	1,604	36 47	35 33
Alabama Mississippi	- 1	3 1	19 6	27	-	-	30	28	1,580	976	8	2
WEST SOUTH CENTRAL	7	7	209	20	2	-	32	8	1,289	421	4	1
Arkansas		_	4	183 14	9	1 -	83 15	56 21	4,418 859	2,460 157	150	77
Louisiana	1	5	26	40	-	-	9	-6	479	140		3.7
Texas	1 5	1 1	9 170	18 111	- 8	- 1	5 54	- 29	295 2,785	291 1,872	150	1 74
MOUNTAIN	_	_	10	35	1	_	50	47	ĺ	-	150	
Montana	-	-	2	3	-	-	4	9	3,606 311	2,432 121	51	130 11
Idaho	-	_	-	11 5	<u>-</u>	-	4	2 2	270	268	6	14
Colorado	_	-	4	3	_	_	21	15	148 1,220	26 894	15 8	79 8
New MexicoArizona	-	-	3	4	1	-	11	5	398	281	NN	-
Utah	-	5 90	- 1	3 6	_		6	10 4	573 563	539 223	13	7 11
Nevada	-	-	1	-	-	-	-	-	123	80	- 4	
PACIFIC	-	-	8	1	10	6	184	216	7,864	5,762	313	207
Washington	-	-	2	-	1	-	23	35	898	679	118	57
California	-	-	2	-	9	6	35 120	38 128	1,221 5,320	941 3,877	44 135	49 92
Alaska Hawaii	- 2	2	4	1	794	-	4	14	365	188	133	7
,	Ξ.	X.	7.0	~	:₩	-	2	1	60	77	3	2
Puerto Rico		5	52	118	-	-	16	8	812	629	69	24

NN-Not Notifiable

Table 2. CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES, EACH DIVISION AND STATE, AND PUERTO RICO, FOR WEEKS ENDED OCTOBER 22, 1960 AND OCTOBER 21, 1961 - Continued

(by place of occurrence. Numbers under diseases are category numbers of the Seventh Revision of the International Lists, 1955)

	Malaria	Meningo infec	ccocal	Psitta- cosis	Strepto- coccal sore throat,	T	yphoid i	ever 040)	Typhus fever, endemic		es in
Area	}				etc.			(3,000)	lative,	Cincento	ani	mals
	110-117	05	7	096.2	050,051	42nd	Week		42 weeks	101		
	1961	1961	1960	1961	1961	1961	1960	1961	1960	1961	1961	196
UNITED STATES	1	43	51	-	4,752	19	28	657	678	1	57	4:
EW ENGLAND	-	2	1	0.50	171	-	1	18	9	-		3
Maine	-	-	-	-	8	-	-	1	2	-	-	
New HampshireVermont	_	_	_	_	3 2	_			-	[-	
Massachusetts	_	1	1	-	41	_	1	12	4			
Rhode Island	-	-	-	-	11	-	-	2	-	-	-	
Connecticut	-	1	-	-	106	-	-	3	3	-	_	
IDDLE ATLANTIC	_	8	10	-	148	4	2	81	47	_	2	
New York	-	5	6	-	101	4	1	45	31	-	1	
New Jersey	-	1	1	-	22	-	<u> </u>	17	2	-	-	
Pennsylvania	-	2	3	-	25	-	1	19	14	-	1]
AST NORTH CENTRAL	-	15	15	-	320	2	3	83	90	-	8	
Ohio	_	_	4	_	48	-	2	28	26	-	2	
Illinois	_	5	3		81 53	1	1	20 26	23 21		1 1	1
Michigan-	_	8	6	-	63	1	_	6	13	[]	2	
Wisconsin	-	2	2	-	75	-	-	3	7	-	2	
EST NORTH CENTRAL	_	_	1	_	148	_	_	33	37	27	15	
Minnesota	_	_	_	-	9	-	_	5	1	_	6	1
Iowa	-	-	-	-	39	-	-	2	6		2	1
Missouri	-	-	-	-	1	-	-	20	22	-	3	
North Dakota	-	-	-	_	48	-	-	-	1	-	0,000	
Nebraska	_	_	1	_	i - 1	Ī		3	3 2	1	4	
Kansas	_	_	Î	_	51	-	-	2	2	2 1	_	
OUTH ATLANTIC	_	6	10	_	404	4	4	108	103	1	6	
Delavare-	S-2	_		-	1			1	1	2	_	
Maryland	-	1	2	-	8	-	-	2	5	[-]	-	
District of Columbia	-	-	1	-	3	-	1	10	8	-	296	1,0
Virginia	-	2	2	_	110 132	1	1 1	18	23 10		2	
North Carolina	-	2	i]	41	_		14	8	ī	1	
South Carolina	-	300	_	-	22	-	_	8	12	-	-	
Georgia-	-	-	-	-	13	2	1	34	24	-	1	
Florida	-	1	-	-	74	1	-	12	12	-	2	
AST SOUTH CENTRAL	-	2	3	-	979	1	8	74	102	-	4	
Nentucky	-		-	-	50	-	4	18	23	- Te -	3	
Tennessee	-	1	1	_	883 18	1	2 2	10	55 19	-	1	
Mississippi	_	1	1		28	-	-	2	5		_	
EST SOUTH CENTRAL		_		_	_	4	ł			1	17	
AIKANSAS-	_	-	-		518 -	3	6	142 28	188 46		17 4	
Louisiana					2		i	21	57	-	1	
Oklahoma	_	-	-	-	5	-	-	12	12	-	-	
Texas-	-	-	-	-	511	1	1	81	73	-	12	
OUNTAIN	12	1	5	_	1,094	2	2	56	38		1	
MUNTana	-	- 1	-	-	33	1	1	17	11	! -	-	l
Idaho		-	-	-	- 47 76	-	-	1 3	3 4	-	-	
Colorado	-		1	-	265		_	6	1	1 [-	
New Mexico	-	1	2		363	1	1	16	9	-	-	
Arizona	-	· **	1	_	161	-	_	7	9	-	1	-
UCBD	:-:	770	1		149		-	= 2	1	-	-	1
Nevada		75	*	-		-	-	4	-	-	-	
CIFIC	1	9	6	-	970	2	2	62	64	- 1	4	
"abungton	-	2	1	-	252	-	-	7	5	-	-	
oregon	-	1	2	-	31	-	2	1 5/	8		4	-23
California	1	8	3	-	632 14	2	2	54	50 1	- '	4	
Hawaii	-	-		1,550	41		_		-			
	-	_										
lerto Rico	-	-	1	-	7	- 1	1	18	19		-	1



The chart shows the number of deaths reported for 117 major cities of the United States by week for the current year, a 5-week moving average of these figures plotted at the central week, and an adjusted average for comparison. For each region the adjusted average was computed as follows: From the total deaths reported each week for the years 1956-1960, 3 central figures were selected by eliminating the highest and lowest figure reported for that week. A 5-week moving average of the arithmetic mean of the 3 central figures was then computed with adjustment to allow for population growth in each region. The average value of the regional increases was 2 percent which was incorporated in the adjusted average shown in the chart.

Table 4 shows the number of death certificates re-

ceived during the week indicated for deaths that occurred in selected cities. Figures compiled in this way, by week of receipt, usually approximate closely the number of deaths occurring during the week. However, differences are to be expected because of variations in the interval between death and receipt of the certificate and because of incomplete reporting due to holidays or vacations. If a report is not received from a city in time to be included in the total for the current week, an estimate is used.

The number of deaths in cities of the same size may also differ because of variations in the age, race, and sex composition of the populations and because some cities are hospital centers serving the surrounding areas. Changes from year to year in the number of deaths may be due in part to population increases or decreases.

Table 3. DEATHS IN SELECTED CITIES BY GEOGRAPHIC DIVISIONS

(By place of occurrence and week of filing certificate. Excludes fetal deaths. Data exclude figures shown in parentheses in table 4)

	42nd week	41st week	Adjusted	Percent change,	Cumulative, first 42 weeks			
Area	ended Oct. 21, 1961	ended Oct. 14, 1961	average, 42nd week 1956-60	adjusted average to current week	1961	1960	Percent change	
TOTAL, 117 REPORTING CITIES	11,820	10,613	11,222	+5.3	479,401	482,510	-0.6	
New England (14 cities) Middle Atlantic (20 cities) East North Central (21 cities) Bouth Atlantic (11 cities) East South Central (8 cities) West South Central (15 cities) Mountain (6 cities) Pacific (15 cities)	680 3,328* 2,471 821* 1,075 519 918 387 1,621	671 2,922 2,351 709 972 463 952 344 1,229	673 3,105 2,396 807 944 507 1,004 364 1,422	+1.0 +7.2 +3.1 +1.7 +13.9 +2.4 -8.6 +6.3 +14.0	29,393 136,448 102,459 32,697 41,466 21,661 41,020 15,320 58,937	30,286 133,993 104,326 33,525 41,357 21,861 42,084 15,107 59,971	-2.9 +1.8 -1.8 -2.5 +0.3 -0.9 -2.5 +1.4	

^{*}Includes estimate for missing reports.

Table 4. DEATHS IN SELECTED CITIES

(By place of occurrence and week of filing certificate. Excludes fetal deaths)

Area	42nd week ended Oct.	41st week ended Oct.		ative, 2 weeks	Area	42nd week ended Oct.	41st week ended Oct.	Cumula first 42	
	21, 1961	14,	1961	1960		21, 1961	14, 1961	1961	1960
			1001	1500				1501	1500
NEW ENGLAND:	ı				LIBOR NODEK CENTRAL C				
Boston, Mass	256	208	10 110	10 570	WEST NORTH CENTRAL—Con.	256	207	0 020	10 201
Bridgeport, Conn	30	38	10,110 1,597	10,579 1,714	St. Louis, Mo	77	57	9,838 2,763	10,301 2,907
Cambridge, Mass	34	32	1,225	1,305	Wichita, Kans	54	34	1,953	1,943
Fall River, Mass	31	33	1,139	1,181	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,		-	1,755	1,,,43
Hartford, Conn	51	57	2,014	2,047	SOUTH ATLANTIC:				
Lowell, Mass.	19	25	1,015	1,000	Atlanta, Ga	122	120	4,756	4,911
Lynn, Mass	20	24	913	1,023	Baltimore, Md	295	189	10,286	10,556
New Bedford, Mass	33	20	1,089	1,020	Charlotte, N.C	36	49	1,512	1,628
New Haven, Conn	39	37	1,901	1,876	Jacksonville, Fla	59	76	2,420	2,481
Providence, R.I.	44	54	2,590	2,664	Miami, Fla	73	83	3,167	3,038
Somerville, Mass	19	17	552	554	Norfolk, Va	49	47	2,081	1,676
Springfield, Mass	41	40	1,846	1,896	Richmond, Va	73	87	3,240	3,264
Waterbury, Conn Worcester, Mass	21 42	33 53	1,116	1,149	Savannah, Ga	32	28	1,370	1,398
"Olcestel, Mass.	42	23	2,286	2,278	St. Petersburg, Fla	(57)	(63)	(2,821)	(2,946)
MIDDLE ATLANTIC:					Tampa, Fla.	76	72 107	2,778	2,756
Albany, N.Y	53	40	1,920	1,791	Washington, D.C Wilmington, Del	208 52	197 24	8,195	8,059
Allentown, Pa	39	24	1,444	1,462		22	24	1,661	1,590
Buffalo, N.Y	170	98	6,111	6,072	EAST SOUTH CENTRAL:				
Camden, N.J	34	34	1,775	1,763	Birmingham, Ala	93	76	3,571	3,542
Elizabeth, N.J	35	29	1,233	1,224	Chattanooga, Tenn	43	31	1,953	1,982
Erie, Pa	19	35	1,620	1,619	Knoxville, Tenn	11	16	1,145	1,178
Jersey City, N.J	89	58	2,898	2,989	Louisville, Ky	118	93	4,757	4,813
Newark, N.J	100*	82	4,232	4,091	Memphis, Tenn	109	114	4,738	4,671
New York City, N.Y	1,606	1,688	69,665	68,288	Mobile, Ala	41	36	1,703	1,730
Paterson, N.J	35	40	1,628	1,608	Montgomery, Ala	26	30	1,338	1,450
Philadelphia, Pa	484	364	21,256	20,529	Nashville, Tenn	78	67	2,456	2,495
Pittsburgh, Pa	286	112	8,035	8,040	LECTI COLUMN CHANGES			100	
Reading, Pa	23	27	978	993	WEST SOUTH CENTRAL:	38	28	1 // 20	1 / 2/
Rochester, N.Y	108	81	4,223	4,201	Austin, Tex	44	37	1,428 1,169	1,434
Scranton, Pa	39	26	1,023	989	Corpus Christi, Tex	13	21	-	1,216
Syracuse, N.Y	41 64	33 63	1,485	1,569	Dallas, Tex	96	134	910	987
Trenton, N.J	45	31	2,560 1,891	2,614 1,733	El Paso, Tex	20	37	5,221 1,467	5,189 1,596
Utica, N.Y	28	30	1,180	1,136	Fort Worth, Tex	67	60	2,716	2,783
Yonkers, N.Y	30	27	1,291	1,282	Houston, Tex	155	170	6,920	7,076
E E	30		1,271	1,202	Little Rock, Ark	43	51	2,355	2,386
EAST NORTH CENTRAL:			1		New Orleans, La	177	155	7,062	7,474
Akron, Ohio	57	61	2,398	2,384	Oklahoma City, Okla	63	63	3,091	3,153
Canton, Ohio	34	35	1,301	1,461	San Antonio, Tex	88	104	4,264	4,224
Chicago, Ill	726	704	31,001	32,204	Shreveport, La	63	37	2,115	2,273
Cincinnati, Ohio	141	161	6,533	6,605	Tulsa, Okla	51	55	2,302	2,293
Cleveland, Ohio	219	210	8,504	8,821	MOURIMATA				
Dayton, Ohio	129	95	4,796	4,931	MOUNTAIN:	20	0,1	1 000	William William
Detroit, Mich	82 318	75 309	3,355 13,903	3,127 14,203	Albuquerque, N. Mex	39 12	34 16	1,320	1,293
Evansville, Ind	35	33	1,516	1,535	Colorado Springs, Colo Denver, Colo	125	110	686	702 5 005
Flint, Mich.	41	34	1,778	1,684	Ogden, Utah	123	12	4,841 689	5,005 700
Fort Wayne, Ind.	30	38	1,588	1,535	Phoenix, Ariz	98	87	3,460	3,195
Gary, Ind	34	37	1,283	1,299	Pueblo, Colo	19	15	710	689
Grand Rapids, Mich	52	50	1,902	1,730	Salt Lake City, Utah	37	37	2,008	2,038
Indianapolis, Ind	172	136	5,955	6,091	Tucson, Ariz	45	33	1,606	1,485
Madison, Wis	23	20	1,357	1,320			arts in the	n that a	_,
Milwaukee, Wis	132	117	5,145	5,249	PACIFIC:			en lide	
Peoria, Ill	16	35	1,186	1,272	Berkeley, Calif	27	12	728	712
Rockford, Ill	25	30	1,192	1,184	Fresno, Calif	(45)	(35)	(1,809)	(1,821)
South Bend, Ind.	32	26	1,204	1,223	Glendale, Calif	(42)	(18)	(1,387)	(1,588)
Toledo, Ohio	113	97	4,142	4,171	Honolulu, Hawaii	42	41	1,690	1,728
Youngstown, Ohio	60	48	2,420	2,297	Long Beach, Calif	67	48	2,329	2,272
WEST NORTH CENTRAL:				- 1	Los Angeles, Calif	560	441	20,814	21,087
Des Moines, Iowa			2 200	2 206	Oakland, Calif Pasadena, Calif	116	100	4,054	4,040
Duluth, Minn.	57	53	2,268	2,286	Portland, Oreg	47	25	1,397	1,444
Kansas City, Kans	28	18	1,077	1,070	Sacramento, Calif	92	95	4,457	4,624
Kansas City, Mo	35* 141	131	1,570 5,376	1,464 5,244	San Diego, Calif.	55 118	57	2,587	2,439
Lincoln, Nebr.	(30)	(27)	(1,171)	(1,090)	San Francisco, Calif	232	1/49	3,735	3,823
Minneapolis, Minn	112	79	4,947	5,237	San Jose, Calif	(31)	(29)	8,128	8,301 (1,450)
Omaha, Nebr	61	88	2,905	3,073	Seattle, Wash	167	97	(1,461) 5,439	5,798
			,		Spokane, Wash	55	42	1,975	1,995
							-74-		-9
*Estimate - based on average				1	Tacoma, Wash	43	34	1,604	1,708

⁽⁾ Figures shown in parenthesis are from cities which have been reporting less than five years and hence are not included in Table 3.

South Dakota has also reported its first case of bat rabies. On September 10, 1961, a man and his wife were trimming trees on their grounds. The wife was wearing a pullover-type sweater. In removing the sweater, she was bitten on the tip of the finger and her first impression was that she was stung by a wasp. The bat fluttered to the ground from the sweater and apparently made no attempt to escape when picked up and placed in a jar. A science teacher was asked if he would care to use the bat in any of his classroom studies, and since he was aware of the possibility of rabies in bats, suggested that it be submitted for diagnosis. The bat was presented dead to a veterinarian at Miller, South Dakota, on September 11th and was forwarded at once to South Dakota State College. It was identified as a hoary bat. A positive diagnosis of rabies was made on September 12. The woman was informed and was started on immunizations that day.

To date, bat rabies cases have been confirmed in 35 States.

(Reported by A. L. Frechette, M.D. Commissioner, Commonwealth of Massachusetts Department of Public Health; and Dr. G. S. Harshfield, South Dakota State College and B. Diamond, South Dakota State Department of Health.)

INTERNATIONAL NOTES - QUARANTINE MEASURES

Smallpox - Belgium

The World Health Organization reported a case of smallpox in Brussels, Belgium. The report indicated that a 17-month-old child had entered Brussels on October 12 on a commercial airlines from Leopoldville, The Congo. The child was hospitalized on October 14; the diagnosis of smallpox was made on October 16; and the child died on October 18. One traveler who accompanied the child from Leopoldville to Rome flew on to Idlewild Airport, New York City. This traveler had a valid vaccination certificate and his family has been vaccinated and will be kept under surveillance.

All travelers from Brussels are required to have valid smallpox certificates.

Cholera - Far East

The World Health Organization declared Hong Kong and Macao free of cholera on October 12. Sarawak was declared free of cholera on October 19. Valid cholera vaccination certificates and surveillance of unvaccinated persons are no longer required for travelers from these areas to the United States.

Unofficial information received indicates that vaccination against cholera is still being required for all arrivals into Formosa. SOURCE AND NATURE OF MORBIDITY DATA

These provisional data are based on reports to the Public Health Service from the health departments of each State and Puerto Rico. They give the total number of cases of certain communicable diseases reported during the week usually ended the preceding Saturday. Total figures for the United States and the Pacific Division include data for the States of Alaska and Hawaii. Cases of anthrax, botulism, and rabies in man are not shown in table 2, but a footnote to table 1 shows the States reporting these diseases. When diseases of rare occurrence are reported by a State (cholera, dengue, plague, louse-borne relapsing fever, smallpox, louse-borne epidemic typhus, and yellow fever) this is noted below table 1.

Hibrary 81 CDC 7 61 HEALTH, EDUCATION, AND WELFARE
PUBLIC HEALTH SERVICE
Communicable Disease Center
Atlanta 22, Georgia

Official Business

