Morbidity and Mortality

Weekly Report

PUBLIC HEALTH SERVICE

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE

Prepared by the

COMMUNICABLE DISEASE CENTER

MElrose 4-5131

For release November 24, 1961

Paral

Total

Atlanta 22, Georgia

Vol. 10 No 46

Provisional Information on Selected Notifiable Diseases in the United States and on Deaths in Selected Cities for Week Ended November 18, 1961

Poliomyelitis — There was no further decline in reports of poliomyelitis for the week ending November 18, 1961. A total of 30 cases, 23 paralytic, is to be compared to reports of 29 cases, 22 paralytic, during the previous week. The same period in 1960 brought reports of 60 total cases, 46 paralytic.

Cumulative totals for 1961 are shown below in comparison with the previous four years.

Polio (Cumulated Weekly) Through the 46th Week

	10	L USI I IV	e reurs		
	1961	1960	1959	19 58	1957
ytic	797	2101	5310	2790	2020
	1228	3025	8029	5520	5692

Only three States reported more than two cases: New York with nine cases, Michigan with four cases, and Louisiana with three. In each of these States, the origin of the cases has been from different counties.

Hepatitis - There have been 1215 cases of hepatitis reported for the week ending November 18. Though this is 173 more cases than the 1042 cases reported last week, it is too early to tell whether this reflects the beginning of a seasonal increase. The number of reported cases is still running several hundred a week more than that reported last year. This increase is generally reflected over the United States.

Table I. Cases of Specified Notifiable Diseases: United States

(Cumulative totals include revised and delayed reports through previous week)

Disease	4	6th Week		Cumulative						
(Seventh Revision of International Lists, 1955) * Weekly incidence low or sporadic Data not available Quantity zero	400	Ended	Median 1956-60	F1	rst 46 wee	eks .	Since s	Approxi- mate		
	Nov.	Nov. 19, 1960		1961	1960	Median 1956-60	1960-61	1959-60	Median 1955-56 to 1959-60	low point
Anthrax062	1	N. /	*	7	17	*	*	*	*	*
Botulism049.1	-	-	*	5	10		*	*	*	*
Brucellosis (undulant fever) 044	9	8	9	520	686	714	*	*	*	*
Ulphtheria055	12	44	44	508	692	758	199	347	374	July 1
Encephalitis, infectious	31	23	32	1,491	1,705	1,909	1,491	1,705	1,909	Jan. 1
Berum	1,215	1,039	334	65,246	34,787	17,227	12,429	9,098	3,190	Sept. 1
Malaria110-117	1	1	*	56	64	*	*	*	*	*
Measles	2,653	2,483	2,471	399,790	413,311	459,603	14,162	13,235	14,407	Sept. 1
Meningitis, aseptic340 pt.	48	58		2,909	2,751		2,909	2,751		Jan. 1
eningococcal infections	44	41	41	1,892	1,952	2,234	377	414	450	Sept. 1
Poliomyelitis080	30	6.3	162	1,228	3,025	5,680	1,125	2,814	5,229	Apr. 1
Paralytic	23	48	109	797	2,101	2,010	737	1,936	2,595	Apr. 1
Monparalytic080.2	3	11	37	291	614	2,748	266	588	1,844	Apr. 1
Unspecified080.3	4	4	16	140	310	922	122	290	790	Apr. 1
Bittacosis	2	2	*	58	90	*	*	*	*	*
Parties in man			*	3	2	*	*	*	*	*
***Prococcal sore throat.	10 -		44.04		0 -			. 1	14.23	- 33/01 14 15/4
14411101110 con-1 at favor 050 051	5,561	5,759		279,568	270,041		60,357			Aug. 1
TANDOID fever	29	12	22	745	746	949	631	615	769	Apr. 1
Trus fever, endemic101	2	1	*	39	59	*	*	*	E*:	5 Militar
Rabies in animals	59	41	58	3,048	3,072	3,814	414	301	431	Oct. 1

Pigures For Current Week Exclude Missing Report From Alaska

Anthrax - South Carolina I

Infectious Encephalitis - Cases of acute infectious encephalitis are reported on a weekly basis to the Communicable Disease Center. These cases, based largely on clinical diagnosis, include a variety of central nervous system infections and related diseases. The total of such cases through the 46th week of 1961 is 1491. This figure, in comparison with previous years, reflects a decline in incidence. Although a variety of factors can be postulated as contributing to this decline, the one most apparent is the absence of a summer-tall peak for reported cases. Such a peak has been found in past years, to correlate with outbreaks of arthropod-borne encephalitis. In 1960, and again this year, there has been very little activity of these viruses. With the onset of winter-like conditions throughout most of the country, the arbovirus season, from late July to early October, has passed and it is unlikely that further cases will occur. Review of special reports received from State epidemiologists and others reveals limited evidence of the presence of arbovirus activity but shows almost complete absence of human involvement this year. California and Utah the only States reporting human cases have each had three serologically confirmed cases. One of the California cases was Western encephalitis and the other two were St. Louis encephalitis; two of the Utah cases were Western encephalitis while the other one was a mixed St. Louis and Western encephalitis infection. Texas, with 22 cases last year, has yet to confirm its first arbovirus infection in 1961. Sporadic isolations of virus from mosquito pools and evidence serologically of equine illness in New Mexico, Arizona and Wyoming are testimony to the continued presence of the arboviruses. The outbreaks of Eastern encephalitis among pheasant flocks in Florida and Connecticut were reported in Morbidity and Mortality Report Vol. 10, No. 45, November 17, 1961.

EPIDEMIOLOGICAL REPORTS

Acute Respiratory Disease - Arizona

An epidemic of an influenza-like illness has been reported from Northeastern Arizona. The epidemic began in early November on the Hopi Indian Reservation, which has a population of approximately 3,500. An estimated 200-300 Hopi, primarily grade school children have been affected.

The disease is characterized by sudden onset of malaise with fever usually not above 103°. Muscle aches in the legs, arms, neck, and back are seen in most patients. A mild sore throat with cough is reported by a large percentage of patients. Frontal headache, vertigo,

and epistaxis are some of the other characteristic symptoms. Photophobia and diarrhea have not been present. Physical examination generally has been unrevealing with clear chests and no significant throat findings. As could best be determined, the illness lasts for five or six days. None of the cases has been hospitalized. There have been no fatalities.

The epidemic is most active in the grade schools on the reservation where school absenteeism over the last two weeks has exceeded 10 percent, a rate higher than recorded at any time since 1955. It has been difficult to define the extent of the involvement of the adult Hopi population. Laboratory studies will be performed at the International Influenza Center for the Americas at the Communicable Disease Center, Atlanta, Georgia.

Reports from the State of Arizona Health Department indicate a high incidence of upper respiratory disease in a number of areas throughout the State. However no reports of influenza-like illness have been received from any other State.

(Reported by Dr. Emory Johnson, Asst. Area Medical Officer, Indian Health Service; Dr. David Templin, Medical Officer in Charge at Keans Canyon Indian Hospital; and Dr. Jack Eason, Acting Director of Health, Arizona State Department of Health).

Anthrax - Massachusetts

On July 16 a seven year old boy was hospitalized in Beverly, Massachusetts because of a painful, black ulceration on the neck, which yielded Bacillus anthracis on culture. His fever and general intoxication responded promptly to parenteral penicillin therapy and he was discharged, recovered after 12 days. The infection was most probably contracted through his father's place of employment, a goat hide tannery, which the child had visited on numerous occasions. Exposure was possible both on the plant grounds and via his father's work clothes, which were worn home unchanged. Analysis of the work clothes and a small number of surface swabs obtained at the tannery failed to grow Bacillus anthracis.

The number of culture documented cases of B. anthracis infections in the United States is gradually falling (38 in 1955, 23 in 1960). Both the indolent cutaneous ulcers and the more virulent inhalation form of the disease usually are directly traceable to exposure during industrial processing of imported goat hides or hair. The processing mills importing these materials from Pakistan, India, and Africa, are located along the East Coast of the United States. Recent cases of anthrax have occurred in and near plants in Philadelphia, Pennsylvania, South Carolina, and New Hampshire. Because contamination with B. anthracis is present in as many as 50 percent of

(Continued on page 8)

Table 2. CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES, EACH DIVISION AND STATE, AND PUERTO RICO, FOR WEEKS ENDED NOVEMBER 19, 1960 AND NOVEMBER 18, 1961

(By place of occurrence. Numbers under diseases are category numbers of the Seventh Revision of the International Lists, 1955)

		Poliomyelitis 080										
*	(Include		otal ot specified	by type)	Par	alytic C	080.0,080	.1	Nonparalytic		Menin- gitis, aseptic	losis (undu-
Area	46th	Week		Cumulative, first 46 weeks		46th Week		Cumulative, first 46 weeks		080.2		lant fever)
	1961	1960	1961	1960	1961	1960	1961	1960	1961	1960	1961	1961
UNITED STATES	30	63	1,228	3,025	23	48	797	2,101	3	- 11	48	9
EW ENGLAND	2	1	35	226	2	1	25	178			3	
Maine		1	4	47	1	1	4	46	1	- 2	3	
New Hampshire	-	-	2		-] -		-			1	
Vermont	1	-	9	11	1	-	8	6	-	-	-	-
Massachusetts	1	-	14	31	1	-	9	20	-		2	-
Rhode Island		-	1 5	102	-	-	1	78		-		
12 1 TABLE 1 TO 1 T				35		-	3	28	-	-	-	
IDDLE ATLANTIC	10 9	10 4	328 241	451	6	9	219	320	1	-	3	
New York	-	-	35	245 82	5	3	155	159	1	- 4	1	
Pennsylvania	1	6	52	124	1	6	28 36	101			2	
	6								ļ.,	_		
AST NORTH CENTRAL	6 1	13 3	164 46	516 119	5	12	103	317	11-2	-	17	1
Indiana	_	4	18	129	1.5	2 4	21	62	124	-	-	
Illinois	1	3	34	141	1	3	10 17	97 94			3	
Michigan	4	3	34	92	4	3	30	53		- 1	9	3
Wisconsin	2.1	-	32	35		_	25	11	- 20		1	
EST NORTH CENTRAL	_	2	72	173		_	32	101		,		
Minnesota	-	ī	6	54			6	44		2	3 2	4
Iowa	-	-	18	21	-		9	4	-	-	2	
Missouri	-	1	24	44	-	-	7	32	20	1	1	1
North Dakota	-	- 1	4	14	-		1	5	-	- 2	-	art.
South Dakota	-	-	3	5	-	-	1	1	-			
Nebraska	-		8	16 19	-	1 5	4	9	-	-	-	1
Kansas	-	_	,	19	-		4	6	-	-	-	
SOUTH ATLANTIC	6	13	211	566	6	9	154	447	-	4	1	1
Delaware	-		2	1	- 5	-	1	-	-	7	-	
Maryland	2	6	41	150	2	5	31	135	-	1	- T	
District of Columbia		2	3 11	5 44	-	1	3	5	1200		-	
Virginia	1	1	32	58	1	1	11 22	40 47		1	To the	-
North Carolina	- 1	i	21	90	1 2	1	11	67	0.00	1	- 1 7752	
South Carolina	1	2	34	129	1	2	26	87		1		
Georgia		_	30	23	- 41	I	23	21	-			
Florida	2	1	37	67	2	- "	26	45	-	1	1	1
AST SOUTH CENTRAL	1 "	7	83	242	1 - 11	6	48	100	1	1	4 6 1 3	San a
Kentucky	_	-	27	130		-	5	5		-		9-11
Tennessee	-	3	20	48	-	2	9	31	-	1	-	1
Alabama	-	2	11	23	- 11	2	11	23	12	-	-	1 11
Mississippi	1	2	25	41	31 -01	2	23	41	1	-	1	4. 11
EST SOUTH CENTRAL	5	6	150	277	4	5	83	175	1	1	1	1
Arkansas		- 1	20	31	- 18	1 - 7	9	23	-	-	1	1
Louisiana	3	1	54	49	3	1	43	30		- "	-	LIA 215
Oklahoma	-	0	4	17	-	- 1	-	12	-	-	-	-
Texas	2	5	72	180	1	4	31	110	1	1	1	-
NUNTAIN	-	5	45	92	- 12	1	26	47	-	2	10	1
Montana	-	-	4	21	1 -3.		2	15	-	-	-	
Idaho		1	14	10	- 1	-	6	1	-	-	F 30	-4-00
Wyoming	- 1	1	-	20	-	1.	- 8	1	-	-		-
Colorado	- 1	1 1	8	19 7		4.5	8	18 4			1	-11
Arizona		3	8	8	1.7	1	6	6	1574	2	9	100
Utah		3	8	7	I T	_	4	2			-	1
Nevada		2.0	-		_	-		-	16.5-1	_		Alies
ACIFIC		6	140	482	23	5	107	416	_	1	9	-211
Washington		2	25	36	2.0	2	17	36	200		1	=160
Oregon			17	36	_	-	8	19			-	2.77
California		3	93	400	-	2	77	351	_	1	8	14
Alaska		-	-	2		-	II -	2		-		
Hawaii	-	1	5	8	-	1	5	8	1200	-	-	100
			~			1						
Puerto Rico	-	1	7	492	-	1	7	483		-	1	- 0-

Table 2. CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES, EACH DIVISION AND STATE, AND PUERTO RICO, FOR WEEKS ENDED NOVEMBER 19, 1960 AND NOVEMBER 18, 1961 - Continued

(By place of occurrence. Numbers under diseases are category numbers of the Seventh Revision of the International Lists, 1955)


	Diphtheria 055 Encephal infecti							titis, in		Measles		
Area	46th Week		Cumulative, first 46 weeks		082		46th Week		Cumulative, first 46 weeks		085	
	1961	1960	1961	1960	1961	1960	1961	1960	1961	1960	1961	1960
UNITED STATES	12	44	508	692	31	23	1,215	1,039	65,246	34,787	2,653	2,483
NEW ENGLAND			8	12	2	L	46	24	2,118	998	527	186
Maine	-	-	-	2	1	- 1	4	5	151	68	171	100
New Hampshire	-	-	-	-	-	-	6	3	193	30	59	
Vermont	-	'-	7	- 9	1 -	-	29	1 9	187 901	16 490	12 231	10
Rhode Island	-	_		í	311		1	3	233	190	4	1
Connecticut	-	-	1	-	-	-	6	3	453	204	50	4
MIDDLE ATLANTIC	-	1	20	16	9	3	125	153	8,804	4,300	231	640
New York] -	7 -	4	7	2	49	68	3,793	2,329	117	30
New JerseyPennsylvania		1 -	13	2 10	2	- 1	34 42	19 66	2,080 2,931	309 1,662	51 63	255
EAST NORTH CENTRAL	_		16	1		2						
Ohio		_ =	1 16	41 16	7 2	1 1	257 85	216 75	13,008 4,315	6,258 2,148	356 21	590 146
Indiana		-	2	7	-	1	24	29	1,914	717	49	54
Illinois	-	-	10	6	4	:	42	52	2,384	1,362	124	28
Michigan	-	_	3 -	10	1 -	1	96	58	4,067	1,811	102	114
			1	2		1	10	2	328	220	60	248
WEST NORTH CENTRAL		3	46 30	50	1	-	160	66	6,256	2,418	193	161
Iova		-	2	26 8	_	Ī.	62	31	1,462 1,815	403 386	11 102	68
Missouri	-	-	1	2	-	_	38	16	1,357	823	-	8
North Dakota	-	-	4	2	-	-	2	2	132	167	36	78
South Dakota	-	_	7	9		-	2	-	191	141	44	1 1
Kansas] [2 -	1 2	1 -] -	9 29	6 5	613 686	246 252	NN	NN
SOUTH ATLANTIC	6	7	124	209	2	2	167	112	8,281	4,017	184	229
Delaware	_		-	-	1	-	1 107	2	177	245	2	35
Maryland	-		1	1	1	-	11	21	718	406	25	
District of Columbia		- 2	3	-	-	-	3	1	124	54	3]
Virginia	1	3 -	15 1	39 4			37 26	13 30	1,373 1,454	716 760	51 76	27
North Carolina			10	14	-	_	54	10	1,915	367	11	15
South Carolina	3	3	13	50	-	-	9	3	446	146	1	76
Georgia	3	1	39	34	- 1	-	10	1	721	254	-	3
Florida	ĺ		42	67	- 1	2	16	31	1,353	1,069	15	7
EAST SOUTH CENTRAL	1 -	22 13	39 9	102 29	-	3 1	152	131	9,755	4,916	203	105
Tennessee	_	1	3	8		1 1	45 69	23 64	2,846 3,825	1,721 1,654	31 139	62
Alabama	1	1	20	33	-	-	16	26	1,666	1,068	12	1
Mississippi	9-	7	7	32	-	1	22	18	1,418	473	21	1
WEST SOUTH CENTRAL	5	11	235	225	4	3	83	72	4,718	2,685	255	96
Arkansas		- 5	4	16 61	1	<u> </u>	19	8	924	191	2	1
Oklahoma		1	28	25	_	[18	4 9	522 320	157 318	- 52	7
Texas	5	5	194	123	3	3	39	51	2,952	2,019	201	88
MOUNTAIN	-	:E _	12	36	1	_	60	63	3,804	2,693	263	151
Montana	-	-	2	3	[- II	-	5	12	325	146	100	22
Wyoming	-	-	:	11	_ =	-	2	8	279	295	11	28
Colorado	-		4	5 3	[[25	2 22	154 1,281	983	30 28	38
New Mexico	-	_	5	4	1	-	1	22	418	296	NN	3.
Arizona	-	-	-	3	-	-	15	14	618	592	26	40
Utah	_		1	7		Ī -	12	3	599	254	68	18
	_	_						-	130	96	-	- M
PACIFIC	-	-	8 2	1	5	10	165	202	8,502	6,502	441	325
Oregon	1,74	_	-		1	1	19 28	41 30	975	851 1,031	70	91
California	-	-	2		5	9	113	127	5,726	4,321	168	146
Alaska		-	4	1		-		4	391	221		100
Havai1	-	-	-	-	-	-	5	-	67	78	1	1
Puerto Rico	_	7	57	129	i .		19	23	897	684	34	42
	i	· ·	1 -		1		1 1	1	1 ",	004	J4	1 7

NN-Not Notifiable

Table 2. CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES, EACH DIVISION AND STATE, AND PUERTO RICO, FOR WEEKS ENDED NOVEMBER 19, 1960 AND NOVEMBER 18, 1961 - Continued

(By place of occurrence. Numbers under diseases are category numbers of the Seventh Revision of the International Lists, 1955)

	Malaria	Meningoccocal infections		Psitta- cosis	Strepto- coccal sore throat,	Т	yphoid i	fever 040		Typhus fever, endemic	Rabies in	
Area	110-117	05	57	096.2	etc. 050,051	46th Week		Cumulative, first 46 weeks		101	animals	
	1961	1961	1960			1961	1960	1961	1960	1961	1961	1960
UNITED STATES	1	44	41	2	5,561	29	12	745	746	2	59	41
							7				58	
NEW ENGLAND	71.5	4	6	-	239	_	and to	20	11	_		_
Maine	C 15	3	5	5.00	8	-	-	1	2	-	-	- ا
Vermont	100	1	1	121	2	-	-	-	-	-	-	-
Massachusetts		1	3	100	32	-	-	1 .7	-	-	7.00	-
Rhode Island		- 1	-	-	22	_	1 -	14	5	-		-
Connecticut		- 2	1		173	_	l -	3	4	1 -		-
IDDLE ATLANTIC	546	8	5	1	192		Ι,	Į.	1			-
New York		6			110	4	1 1	92 51	53		4	4
New Jersey	:=0	í	1	-	24	-	-	17	33 6	_	4	4
Pennsylvania	-	1	4	1	58	-	.70	24	14		24.0	
AST NORTH CENTRAL		10	9	_	294	7	_					_ ^
Ohio	:51	1	2	-	52	6	2 1	98 39	95	-	5	3
Indiana		î	-	-	56	1	-	21	27		1	1
Illinois	-	2	2	11.57	61	-	-	29	21		-	2
Michigan	-	4	5	-	51	_	1	6	16	1	3	_
Wisconsin	-	2	-	-	74		-	3	7	1 -	1	_
EST NORTH CENTRAL	*	2	1	-	204	-	1	34	46		20	7
Minnesota		1		-	23	_	11	- 5	1		3	3
Iowa	- 7	7	-	-	33	-	-	2	11	-	6	li
Missouri	-	-	-		14	-	-	20	24		5	2
South Dakota		- 5	-	7	85	-	-	-	1	371 -	1	
Nebraska		1	1	- 2	-	-V-	1	3	4	-	4	1
Kansas		-			49		_	1 3	3	-	1	-
SOUTH ATLANTIC	1	6	3		1				2			4 -
Delaware	-	3	-		514	10	2	126	113	1	7	7
Maryland	-	-		11.00	8		1	1 3	1	Cartes 1	10 To	2.70
District of Columbia	_	-	-	-	7 7	1	î	13	6 9			-
Virginia	-		1		107	1	1	20	24		15 15	3
West Virginia	-	2	-		139	-	-	10	12		6	4
North Carolina	-	-	2	-	40	2	- ·	16	9	-	1000	100
South Carolina	1 1		-		15	•	- 5	8	12		S 20 1	_2
Florida	ī	1			1 107	2		36	27	1	-	-
		1555	1.0	-	197	4		19	13		1	-
AST SOUTH CENTRAL	-	2	6	-	965	3	3	80	114	100 HDM 1 HI	5	9
KentuckyTennessee	-	-	2	-	62	-	1	18	29	1.00	3	1
Alabama-	- 5	1	3		846	3		50	55	1	2	5
Mississippi	-			_	40	-	1	10	23 7		-	3
EST SOUTH CENTRAL	2	3	5	5	743	5	1			-	-	-
Arkansas		1	-	-	17	2	2 2	155 31	201 52		16	9
Louisiana		2	4	TOTAL TO	4		10.2	25	52 59	-	2	2
Oklahoma	-	-	1	*:	23	2	2	12	12	2		-
Texas			1118	14.00	699	3	-	87	78		14	7
OUNTAIN-	-	1	-		1,405			62	45	40.0		- "
Montana-	-	11.			75	-	-	18	13	_	2	-
Idaho	- 47		10.00	-	74	-	-	1	3		3 -	
Wyoming	8-		-	30-	37	-	-	3	4	*	- 4	1.5
Colorado		1	100	•	411	7	7	7	1		*	
New Mexico-					351			17	12	-	-	-
Arizona	•		0.1		245	-		10	10	-	2	-
Nevada		- :	= 0		212			2	2	5 -	-	
			118 1	1.60	v (V =			4			-	-
ACIFIC	*	8	6	1	1,005	-	1	78	68	1	-	2
Washington Oregon	-	1	2	-	274		-	7	5	-		-
California	-	7	4	1	673		1	1	8		1	-
Alaska			- 4	1	673		1	68	54	-	- /	2
Revail	-		2		27			2	1	1		
	075110		-				1		-	1		
uerto Rico	n	1	1			1	1	22	20		2	2 .



The chart shows the number of deaths reported for 117 major cities of the United States by week for the current year, a 5-week moving average of these figures plotted at the central week, and an adjusted average for comparison. For each region the adjusted average was computed as follows: From the total deaths reported each week for the years 1956-1960, 3 central figures were selected by eliminating the highest and lowest figure reported for that week. A 5-week moving average of the arithmetic mean of the 3 central figures was then computed with adjustment to allow for population growth in each region. The average value of the regional increases was 2 percent which was incorporated in the adjusted average shown in the chart.

Table 4 shows the number of death certificates re-

ceived during the week indicated for deaths that occurred in selected cities. Figures compiled in this way, by week of receipt, usually approximate closely the number of deaths occurring during the week. However, differences are to be expected because of variations in the interval between death and receipt of the certificate and because of incomplete reporting due to holidays or vacations. If a report is not received from a city in time to be included in the total for the current week, an estimate is used.

The number of deaths in cities of the same size may also differ because of variations in the age, race, and sex composition of the populations and because some cities are hospital centers serving the surrounding areas. Changes from year to year in the number of deaths may be due in part to population increases or decreases.

Table 3. DEATHS IN SELECTED CITIES BY GEOGRAPHIC DIVISIONS

(By place of occurrence and week of filing certificate. Excludes fetal deaths. Data exclude figures shown in parentheses in table 4)

	46th week	45th week	Adjusted average, 46th week 1956-60	Percent change, adjusted average to current week	Cumulative, first 46 weeks			
Area	ended Nov. 18, 1961	ended Nov. 11, 1961			1961	1960	Percent change	
TOTAL, 117 REPORTING CITIES	11,859	11,161	11,559	+2.6	525,402	527,913	-0.5	
New England (14 cities) Middle Atlantic (20 cities) East North Central (21 cities) West North Central (9 cities) South Atlantic (11 cities) East South Central (15 cities) West South Central (15 cities) Mountain (8 cities) Pacific (15 cities)	739* 3,386* 2,621 769 987 560 987 368* 1,442*	768 2,970 2,453 789 961 512 874 341 1,493	684 3,110 2,474 845 981 531 1,062 368 1,504	+8.0 +8.9 +5.9 -9.0 +0.6 +5.5 -7.1 0.0	32,373 149,240 112,402 35,845 45,404 23,744 44,872 16,785 64,737	33,181 146,789 114,394 36,710 45,236 23,820 45,962 16,542 65,279	-2.4 +1.7 -1.7 -2.4 +0.4 -0.3 -2.4 +1.5 -0.8	

^{*}Includes estimate for missing reports.

Table 4. DEATHS IN SELECTED CITIES

(By place of occurrence and week of filing certificate. Excludes fetal deaths)

Area	46th week ended Nov.	45th week ended Nov.	Cumula first 4	ative, 6 weeks	Area	46th week ended Nov.	45th week ended Nov.	Cumula first 46	
	18, 1961	11, 1961	1961	1960		18, 1961	11,	1961	1960
			50,000				A F		10
EW ENGLAND:			Sec. 1		WEST NORTH CENTRAL-Con.:			1	
Boston, Mass	256	264	11,116	11,559	St. Louis, Mo	222	223	10,826	11,216
Bridgeport, Conn	55	60	1,793	1,877	St. Paul, Minn	82	61	3,028	3,182
Cambridge, Mass	24	25	1,338	1,419	Wichita, Kans	55	57	2,141	2,134
Fall River, Mass	35	32	1,273	1,290			i	i	
Hartford, Conn	46	69	2,235	2,251	SOUTH ATLANTIC:				
Lowell, Mass	38	46	1,150	1,093	Atlanta, Ga	120	118	5,226	5,393
Lynn, Mass	25	23	1,001	1,115	Baltimore, Md	225	249	11,247	11,504
New Hedford, Mass	25* 33	27 29	1,188 2,067	1,137 2,048	Charlotte, N.C Jacksonville, Fla	20 66	48 50	1,658 2,657	1,784
Providence, R.I	64	65	2,855	2,942	Miami, Fla	94	60	3,473	3,28
Somerville, Mass	13	7	600	606	Norfolk, Va	44	64	2,279	1,83
Springfield, Mass	48	38	2,030	2,083	Richmond, Va	89	74	3,560	3,56
Waterbury, Conn	20	25	1,209	1,272	Savannah, Ga	26	33	1,493	1,55
Worcester, Mass	57	58	2,518	2,489	St. Petersburg, Fla	(51)	(67)	(3,069)	(3,23
					Tampa, Fla	55	38	2,978	3,00
IDDLE ATLANTIC:	e /		2 120	1 007	Washington, D.C	206	186	9,008	8,85
Albany, N.Y.	54 30	57	2,130	1,987	Wilmington, Del	42	41	1,825	1,75
Allentown, Pa.	30 156*	32 141	1,567 6,673	1,600 6,616	FACE COURT CONTRACT				
Buffalo, N.Y	37	43	1,931	1,951	EAST SOUTH CENTRAL:				
Elizabeth, N.J	37	26	1,371	1,353	Birmingham, Ala	89	81	3,918	3,88
Erie, Pa.	34	32	1,783	1,775	Chattanooga, Tenn Knoxville, Tenn	40	46	2,144	2,15
Jersey City, N.J	95	49	3,175	3,244	Louisville, Ky	40 121	43 128	1,281	1,28
Newark, N.J	101*	86	4,648	4,469	Memphis, Tenn	119	84	5,199 5,163	5,20 5,07
New York City, N.Y	1,674	1,586	76,179	74,951	Mobile, Ala	38	35	1,844	1,89
Paterson, N.J	43	36	1,787	1,755	Montgomery, Ala	39	33	1,487	1,58
Philadelphia, Pa	512	.406_	23,176	22,316	Nashville, Tenn	74	62	2,708	2,73
Pittsburgh, Pa	233	90	8,751	8,830				1	
Reading, Pa	21	20	1,071	1,091	WEST SOUTH CENTRAL:	22	20		
Rochester, N.Y.	107	110	4,659	4,622	Austin, Tex	33	33	1,552	1,54
Schenectady, N.Y	23	22	1,110	1,076 1,727	Raton Rouge, La	23 27	31 30	1,290	1,31
Scranton, Pa	36 is 86	34 74	1,618 2,831	2,876	Dallas, Tex	125	98	1,014 5,711	1,07
Trenton, N.J	43	C 43	2,056	1,888	El Paso, Tex	37	25	1,603	5,71 1,74
Utica, N.Y	32	50	1,309	1,245	Fort Worth, Tex	72	66	2,970	3,04
Yonkers, N.Y	32	33	1,415	1,417	Houston, Tex	168	111	7,568	7,69
					Little Rock, Ark	39	57	2,576	2,62
AST NORTH CENTRAL:					New Orleans, La	198	140	7,726	8,20
Akron, Ohio	57	56	2,628	2,598	Oklahoma City, Okla	89	72	3,397	3,45
Canton, Ohio	38	38	1,449	1,596	San Antonio, Tex	95	92	4,642	4,57
Chicago, Ill.	817	751	34,045	35,237	Shreveport, La Tulsa, Okla	49	64	2,321	2,47
Cincinnati, OhioCleveland, Ohio	189 225	153 198	7,186 9,318	7,202 9,690	idisa, okia:	32	55	2,502	2,51
Columbus, Ohio	144	109	5,259	5,447	MOUNTAIN:			-0.736	
Dayton, Ohio	75	66	3,655	3,464	Albuquerque, N. Mex	47	35	1,465	1,46
Detroit, Mich	350	352	15,237	15,553	Colorado Springs, Colo	9	14	751	76:
Evansville, Ind	38	28	1,657	1,700	Denver, Colo	100	95	5,283	5,49
Flint, Mich	40	41	1,950	1,847	Ogden, Utah	20	19	776	75
Fort Wayne, Ind	36	38	1,744	1,687	Phoenix, Ariz	80	80	3,760	3,47
Gary, Ind.	27	27	1,406	1,405	Pueblo, Colo	17*	13	774	7.5
Grand Rapids, Mich	50	51	2,078	1,884	Salt Lake City, Utah	51	45	2,212	2,22
Indianapolis, Ind	126	149	6,540	6,670	Tucson, Ariz	44	40	1,764	1,62
Madison, Wis.	43	51	1,527	1,492	PACIFIC:				
Milwaukee, Wis	103	119	5,626	5,734	Berkeley, Calif	12	9	776	76
Peoria, Ill.	40	21	1,307	1,391	Fresno, Calif	(37)	(42)	(1,981)	76
Rockford, Ill	35	27	1,307	1,321	Glendale, Calif	(36)	(33)	(1,535)	(1,69
Toledo, Ohio	34	23	1,310 4,516	1,349 4,595	Honolulu, Hawaii	42	43	1,850	1,89
Youngstown, Ohio	96 58	100 55	2,657	2,532	Long Beach, Calif	69	54	2,555	2,50
	30	- 22	2,05.	-,55-	Los Angeles, Calif	514	536	22,912	22,95
EST NORTH CENTRAL:		ļ			Oakland, Calif	91	117	4,465	4,35
Des Moines, Iowa	55	43	2,468	2,518	Pasadena, Calif	35*	42	1,532	1,57
Duluth, Minn.	21	36	1,183	1,162	Portland, Oreg	115	122	4,906	5,02
Kansas City, Kans	33	40	1,728	1,611	Sacramento, Calif	61	69	2,847	2,65
Kansas City, Mo	112	132	5,855	5,767	San Diego, Calif.	103	56	4,082	4,19
Lincoln, Nebr	(28)	(33)	(1,279)	(1,196)	San Francisco, Calif.	169	201	8,896	9,03
Minneapolis, Minn	116	135	5,434	5,748	San Jose, Calif	(36)	(46)	(1,610)	(1,58
Omaha, Nebr	73	62	3,182	3,372	Seattle, Wash.	134	134	5,978	6,26
Estimate - based on average					Spokane, Wash.	61 36	61 49	2,186 1,752	2,20 1,86

⁽⁾ Figures shown in parenthesis are from cities which have been reporting less than five years and hence are not included in Table 3.

the imported goat hides and bales of goat hair tested, prophylactic immunization of plant personnel with vaccine proved 92 percent effective, has been used in many areas since 1955. Limited supplies of an experimental vaccine are available on request from the Anthrax Surveillance Unit, Communicable Disease Center, Atlanta, Georgia.

QUARANTINE MEASURES

Immunization Information for International Travel

No Changes Reported

SOURCE AND NATURE OF MORBIDITY DATA

These provisional data are based on reports to the Public Health Service from the health departments of each State and Puerto Rico. They give the total number of cases of certain communicable diseases reported during the week usually ended the preceding Saturday. Total figures for the United States and the Pacific Division include data for the States of Alaska and Hawaii. Cases of anthrax, botulism, and rabies in man are not shown in table 2, but a footnote to table 1 shows the States reporting these diseases. When diseases of rare occurrence are reported by a State (cholera, dengue, plague, louse-borne relapsing fever, smallpox, louse-borne epidemic typhus, and yellow fever) this is noted below table 1.

Library 81 CDC 7 61 U. S. DEPARTMENT OF
HEALTH, EDUCATION, AND WELFAR
PUBLIC HEALTH SERVICE
Communicable Disease Center
Atlanta 22, Georgia

Official Business

AUTUR MALITUR