

# Provisional Information on Selected Notifiable Diseases in the United States and on Deaths in Selected Cities for Week Ended December 9, 1961

**Respiratory Disease** – Type B influenza has been identified in outbreaks of respiratory disease in Arizona, Florida, Jamaica, Saskatchewan, and Aruba Island (N.W.I.). Respiratory illnesses of unidentified etiology reported in several other areas since mid-November have continued to spread in these areas. No new outbreaks have been reported.

#### Confirmed Type B Influenza Outbreaks

Arizono – The outbreak of respiratory disease which began in mid-November in the Hopi and Navajo Indian Reservation in northeastern Arizona has been identified as Type B influenza. Seven of 10 paired sera obtained during the outbreak show significant titer rises for Type B influenza. Positive sera were obtained from the Hopi Indian Reservation, as far west as Grand Canyon and from Fort Defiance, Arizona, on the New Mexico border.

Clinical characteristics of the illness included fever of over 100-101°, severe sore throat with cough, minimal conjunctivitis, absence of pulmonary involvement, myalgia and frontal headache. The acute symptoms last for two or three days.

The Keams Canyon School on the Reservation presented an unusual opportunity for study of vaccine effectiveness. Those children who board at the school had been vaccinated; the day school children had not been

## Table I. Cases of Specified Notifiable Diseases: United States

(Cumulative totals include revised and delayed reports through previous week)

Disease	142 224	49th Wee	₽k	Cumulative						
(Seventh Revision of International Lists, 1955)	Ended	Ended	himit	First 49 weeks			Since seasonal low week			Approxi- mate seasonal
<ul> <li>Weekly incidence low or sporadic</li> <li>Data not available</li> <li>Quantity zero</li> </ul>		Dec. 10, 1960	<b>Median</b> 1956-60	1961	1960	<b>Median</b> 1956-60	1960-61	1959-60	Median 1955-56 to 1959-60	low point
Anthrax062	100	1	*	7	18	*	*	*	*	*
Botulism049.1	1	1116-2-1	*	6	10	*	*	*	*	*
Brucellosis (undulant fever) 044	11	12	12	556	713	758	*	*	*	*
Diphtheria055	27	26	32	556	781	833	247	435	494	July 1
Encephalitis, infectious082	29	24	40	1,567	1,779	2,012	1,567	1,779	2.012	Jan. 1
Repatitis, infectious, and					10.0		-,507	1,775	2,012	San. 1
serum092,N998.5 pt.	1,367	1,159	346	68,926	37,945	18,153	16,223	12,203	4,116	Sept. 1
Malaria110-117	1	3	*	58	71	*	*	*	*	*
Measles085	3,646	4,092	4,060	409,236	423,356	467,143	23,608	23,234	24.357	Sept. 1
Meningitis, aseptic	28	43		2,998	2,886		2,998	2,886		Jan. 1
Meningococcal infections057	52	53	51	2,022	2,083	2,453	507	545	594	Sept. 1
Poliomyelitis080	15	49	92	1,280	3,170	8.347	1,177	2,958	5,576	Apr. 1
Paralytic080.0,080.1	10	33	61	837	2,215	5,548	777	2,049	2,816	Apr. 1
Nonparalytic080.2	3	6	12	304	628	2,122	279	602	1,909	Apr. 1
Unspecified080.3	2	10	19	139	327	677	121	307	851	Apr. 1
Psittacosis096.2	25	1	*	87	95	*	*	*	*	*
Rabies in man094	-	-	*	3	2	*	*	*	*	*
Streptococcal sore throat,		200 - C. L	ELINARDS	1 1 2 2 2 2	COLS	1. <u>1994</u> 10	1		-male ;	1976
including scarlet fever050,051	7,454	6,468		298,309	288,668		79.098			Aug. 1
Typhoid fever040	10	7	16	785	776	1,008	671	645	828	Apr. 1
Typhus fever, endemic101	1	10000	*	41	62	*	* *	*	*	*
Rabies in animals	53	64	81	3,231	3,212	4,001	597	441	636	Oct. 1


Botulism - Arkansas 1

vaccinated. All boarders are brought to the USPHS Indian, Hospital Clinic if there is any indication of illness. Of 320 boarders in the immunized group, only one case of clinical influenza has been noted by the clinic physicians. Of 120 day students who are not immunized, there have been 26 cases of clinical influenza.

Convert minister of	Attack Rate
Immunized	0.3%
Not Immunized	22.0%

Laboratory studies on these groups are in progress.

A Hopi Indian School on the Reservation illustrates well the epidemicity of the illness. The accompanying graph records the total number of absentees each day from this school which has a total enrollment of 190 students.



Canada - The epidemic of influenza which commenced during November in Saskatchewan has spread from the focus at Regina to involve a number of communities in the central and southern regions of that province. Absenteeism in the public schools has ranged from 30 to 60 percent. Clinically, the disease has been relatively mild. The course of illness has been described as being frequently biphasic in character with relapses noted following a four to six day course of the disease. It has been observed that the spread of disease among patients within the hospitals has been rare although many absences have been noted among hospital attendants. Two deaths, directly attributable to the disease, have resulted from a staphylococcus (phage type 80/81) pneumonia. Isolates obtained resemble the Great Lakes variant of Influenza B. No other outbreaks are presently known in Canada.

Aruba Island (N.W.I.) – In October 1961, approximately 200 cases of respiratory disease occurred. Clinical characteristics of the disease are not available; however, eight of nine paired sera obtained during this outbreak were positive for Type B Influenza. Miami, Florida – An outbreak of Type B influenza in a school-bus population was reported in MMWR, Vol. 10, No. 48, December 8, 1961.

Jamaica – A confirmed outbreak of Type B influenza in mining company personnel was reported in MMWR, Vol. 10, No. 48, December 8, 1961.

### **Respiratory Diseases of Unidentified Etiology**

California -. Acute upper respiratory illnesses occurring since the middle of November continue in the Los Angeles and San Francisco areas primarily in school children, and have appeared in most major centers of the State. School absenteeism as high as 30 percent in some schools is reported from northern coastal counties, mountain counties and the interior valley. Two illnesses are occurring, one characterized by malaise, sore throat, fever, cough and some myalgia, and the other by mild upper respiratory symptoms, diarrhea and fever. Laboratory and epidemiologic investigation of these illnesses is under way. There is no unusual absenteeism of teaching staff in the schools or increased absenteeism reported from industrial concerns.

**Oregon** – An outbreak of upper respiratory disease and gastroenteric disease which began in November in counties surrounding Portland has continued to spread, involving Washington County to the northwest and Grant's Pass in the southernmost portion of the State. Early reports of the appearance of the illness in the eastern part of the State have been received. In school children two types of illnesses continue to occur: 1) an upper respiratory disease with fever to 101°, sore throat and vomiting which lasts 3-4 days. 2) a gastrointestinal disease with mild cramps, diarrhea and fever to 101°.

Cases occurring in adults are reported in the Portland area, with upper respiratory symptoms, mild diarrhea and myalgia, and some temperatures of 103°. Several of these have had relapses after several days suggesting a biphasic pattern. There is no reported increase in industrial absenteeism. Laboratory studies are in progress. (Information for this summary was obtained from: Dr. Roslyn Robinson, Chief, International Influenza Center for the Americas, CDC, and a team from the Communicable Disease Center; Dr. Mike Segal, Dr. Ann Beasley, and Dr. Jim Conner from Variety Children's Research Foundation, Miami, Florida; Dr. Henry Renteln, Division of Preventive Medical Services, California State Department of Public Health; Dr. E. W. R. Best, Chief, Epidemiology, Division Department of National Health and Welfare, Otrawa, Canada; Dr. James O. Bond, Director, Bureau Preventable Diseases, Florida State Board of Health; Dr. Grant Skinner, State Epidemiologist, Oregon State Board of Health).

(Continued on page 8.)

# Table 2. CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES, EACH DIVISION AND STATE, ANDPUERTO RICO, FOR WEEKS ENDED DECEMBER 10, 1960 AND DECEMBER 9, 1961

(By place of occurrence. Numbers under diseases are category numbers of the Seventh Revision of the International Lists, 1955)

ter , unter		at 12 million		Po	liomyeli	tis 080						Brucel
To be set in the set of the set o	(Includ		otal ot specified	by type)	Par	alytic (	080.0,080	0.1			Menin- gitis,	losis (undu- lant
Area	49th Week		Cumulative, first 49 weeks		49th Week		Cumulative, first 49 weeks		Nonparalytic 080.2		aseptic 340 pt.	fever)
	1961	1960	1961	1960	1961	1960	1961	1960	1961	1960	1961	1961
UNITED STATES	15	49	1,280	3,170	10	33	837	2,215	3	6	28	11
NEW ENGLAND	- 23	1	40	232	-	1.2	27	183	1	1	2	1
Maine		-	5	49	-		5	48	-	-	-	-
New Hampshire Vermont	1.5	1	2	- 11	1.21	1.1	-		-	-	2	-
Massachusetts	120	1.258	18	34	100	1.12	8	6 23	·	-	1.1.1	-
Rhode Island	144	1	1	103	1.1.1	1.1.1	1	78	-	1	Trainin-	there
Connecticut		-1	5	35	1.00	-	3	28	1000			
MIDDLE ATLANTIC	5	9	336	480	4	6	224	341	1	2	3	A DOCT
New York	5	6	249	260	4	3	160	168	1.00	2	-	-1
New Jersey	1114	1	35	84	1 200	1	28	61		1. E.	-	-
Pennsylvania	1. E.	2	52	136	-	2	36	112		-	3	-
EAST NORTH CENTRAL	1	18	177	553	1	12	110	344	1.000	2	10	4
Ohio	1	1	48	127	1.1	1.1	22	65		1		
Indiana	1	8	21 36	141 150	1	7	12	108		1	-	-
Illinois Michigan	1.1	1	36	94	0	3 1	19	103	-	-	6	2
Wisconsin	1.1.1	5	34	41		1	30 27	55 13	-	in the second	4	2
	2	1	77	175	1	1	35			al. and		_
MINNESOTA	-	i	6	55	1	1	6	102	1	_	3	5
Iova	20120	_	19	22	1.0	÷.	10	45	1		3	2
Missouri	- 1.4	-50	26	44		-	8	32	1.000	1.1		-
North Dakota	1	-	5	14	1.549	1. 2	1	5	1	00000	Dotto	ined.
South Dakota	1	-	4	5	1	1.2-1	2	1			and the same	COURSE 1
NebraskaKansas	12		8	16 19	1.121	1.2	4	9			-	2
					120		The second second				_	1
SOUTH ATLANTIC	1	5	215 2	588	1	5	158	468	-	-	1	1
Delaware	1	2	42	153	1	2	1 32	138	1.0	-	-	-
District of Columbia	- 2.	2.	3	5	-	1.0	3	5		- letter	10 Do 198	and in
Virginia	1 45	3	12	55	1.721	3	12	50	( many	1.1.1	1	10.12
West Virginia	5 H-12	-	33	62	1.257	1.3.5	23	51	-	-	-ALCENTIV	723
North Carolina		-	21	92	-	-	11	69				1000
South Carolina	1.20	-	34 30	130 24		1.1	26	88	-	-	0.000	-
Georgia	1 1 10	0.028	38	67	00	1.620	23	22 45	1	1		1
EAST SOUTH CENTRAL	1 10	6	85	256	1.12	2	49				Constantion of the	-
Kentucky	0.00	4	27	134		-	5	110 5	-		1	
Tennessee	A. 1925	1.1.1	22	54		. 540	10	37	-	+	1	1.000
Alabama			11	23	-	1.250	11	23			-	1001
Mississippi		2	25	45	-	2	23	45	1000		100200	Stall
EST SOUTH CENTRAL	4	3	155	286	3	3	87	184	1		00.0	1
Arkansas	2	2	23	33	2	2	12	25				1000
Louisiana	1	-	55	50	1	-	44	31		-	- 11 - 11 - 11 - 11 - 11 - 11 - 11 - 1	-
Oklahoma		-	4	17				12	-	-	-	1
Texas	1	1	73	186	10.00	1	31	116	1	-	-	-
MOUNTA IN	1	1	48	99	1.22	1.0	28	51	-	-	1	1121
Montana	- 1.5 m	1	4	24	1.00	1.20	2	17	1.1.1		-	Contraction of the local division of the loc
Idaho			14	10 21	2.1		6	1	diam'ne	-		
Wyoming	1		10	21	1.000	1 (C)	10	19	-			1.00
New Mexico	1 22		3	7		1 14	-	4	1-101	1.00		
Arizona			8	9		1.1	6	7	12007	-	1	NAVE.
Utah	나 크		8	7		12	4	2		10.27		200
Nevada	1.00				-		-	-				
ACIFIC	1	5	147	501	-	4	119	432	1	1	7	ALL DAY
Washington		2	31	39 36		2	29 8	39 19	Guine			Same and
Oregon		-3	17 94	416		2	77	364	1	1	7	12.01
California	1	-	94	2	-	-	-	2	-	-		1114
Hawaii	1.1		5	8		1.25	5	. 8				10012
				501	and a	44	-	105	1			
Puerto Rico		1.1.1.1.1.6	7	504	-	1.00	7	495				

WINDOW FROM PARAMETER

# Table 2. CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES, EACH DIVISION AND STATE, AND<br/>PUERTO RICO, FOR WEEKS ENDED DECEMBER 10, 1960 AND DECEMBER 9, 1961 - Continued

(By place of occurrence. Numbers under diseases are category numbers of the Seventh Revision of the International Lists, 1955)

		Diphthe	eria 055		Encepha infect				nfectious ,N998.5 p		Mees	les
Area	49th Week		Cumulative, first 49 weeks		082		49th Week		Cumulative, first 49 weeks		085	
180 1 1900 1280 1 100	1961	1960	1961	1960	1961	<b>196</b> 0	1961	<b>196</b> 0	1961	1960	1961	1960
UNITED STATES	27	26	556	781	29	24	1,367	1,159	68,926	37,945	3,646	4.092
NEW ENGLAND	1.7.7	-	8	12	2		67	22	2,283	1,089	601	388
Maine	10.00	- 1	1. T-1	2	-	-	10	3	176	77	169	
New Hampshire	1.00	100	-	1.24	1		4		203	34	67	7
Vermont		1 1	7	- 9	51		2	-	192	27	28	
Rhode Island			<u> </u>	1	1	0.1	40	8	995 239	519 204	258 18	22
Connecticut	1.1		1	-			9	7	478	228	61	7
MIDDLE ATLANTIC	1.1	1	20	17	8	6	121	137	9,215	4,694	500	1,01
New York	-	1	7	5	8	3	42	64	3,972	2,529	329	39
New Jersey	-	11.20	-	2		2	27	1	2,181	333	70	14
Pennsylvania	1.12	1.1	13	10	0	1	52	72	3,062	1,832	101	47
EAST NORTH CENTRAL	1.0	1.1	16	41	3	1	315	236	13,847	6,890	819	1,090
Ohio	10	1/25	1	16			117	88	4,600	2,400	40	408
Indiana			2 10	7	-		40	39	2,036	797	51	6
Michigan	1.52		3	10	2 1	1	49 86	64 38	2,515	1,496	452	5 15
Wisconsin			1 -	2	-		23	7	4,326 370	232	115	42
WEST NORTH CENTRAL	5		53	63	1	_	128	85	6,626	2,650	77	10
Minnesota	5	-	38	36	-	_	51	22	1,622	483	7	10
Iowa	-	- 1	2	8	1	-	18	10	1,877	417	23	2
Missouri	-	-	1	2	12	-	41	26	1,435	886	4	3
North Dakota			4	3	1.20	- 1997 <b>-</b> 1	2	3	149	178	30	41
South Dakota Nebraska	1.1	1 1	2	10 2	1.5100	647 T.	47	8	198 632	154 271	13	
Kansas	-	-	-	2		-	5	4	713	261	NN	N
SOUTH ATLANTIC	2	10	132	227	3	3	176					372
Delaware	-	-			-	-	3	125	8,807	4,335	257	31
Maryland	1.00	-	1	1	1.1	C 1 -0	8	9	751	440	39	18
District of Columbia		-	3		1		5		135	60	10	
Virginia	1	1	18	40	1.15	3	36	18	1,461	763	95	19
West Virginia	1-12-5	4	11	4	1.5	12	20 62	28	1,534	835	60	3
South Carolina		1	13	50	1.27	1.1	12	25	2,087	423	3	20
Georgia	1	4	43	45	1.00		6	12	742	273	-	25
Florida	1.1	1	42	69	2	1.2	24	20	1,411	1,119	48	27
EAST SOUTH CENTRAL	1.1	4	46	128	2	1	208	192	10,318	5,492	301	429
Kentucky	1.10	4	9	49	1.00		73	88	3,021	1,933	45	326
Tennessee	1.1		3 27	9	1		83	55	4,039	1,853	231	84
Alabama Mississippi			7	37 33	- 1	1	25 27	38	1,772	1,185	15	
Contraction of the second s	17	11							1,486	521	10	103
WEST SOUTH CENTRAL	17	11	258	255 16	1	1	73	56	4,917	2,850	284	102
Louisiana	9	6	39	76	1	-1-1-2-2	12	9	959 540	227	ī	
Oklahoma	1. in .	-	9	25	1.2	1.1.25	8	1	337	322	5	3
Texas	8	5	206	138	-	1	46	40	3,081	2,122	278	94
MOUNTAIN	0.040	31 <u>2</u> 9	12	36	1	1	88	129	3,887	2,964	234	199
Montana	1.00	-	2	3	2	1	3	123	336	175	120	25
Idaho	1	1.000	Criticant	11	1.440	10.00	3	6	303	318	9	40
Wyoming			4	5	1.15.		1		157	36	1	39
New Mexico	100		5	3	1.5	1,000	29	44	1,345	1,079	19	-
Arizona	1.21	125-	1 -	3	1	1	6 19	25	318	335	NN 21	53
Utah	1.2-1	17-	11.5	7		- in i	18	7	626	279	54	24
Nevada	8 <b>-</b>	I	1	1.0	1.0	0-04	9	22	145	124	10	18
PACIFIC	3	to really	11	2	8	11	191	177	9,026	6,981	573	383
Washington	3	1.00	5	1255	1	-	21	31	1,038	933	209	131
Oregon	1 1201	10 250	-	1000	이 가는 것	1	32	23	1,431	1,085	67	60
California	1.00	-	2	1	7	10	135	112	6,071	4,636	255	190
Hawaii	5	-	4	1	-	-	2	7	415	242	37	1.11
	No.	1000	1 2 100	1-21	-	1.0	1	4	71	85	5	
and the second												


4

NN-NOT NOTIFIABLE

# Table 2. CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES, EACH DIVISION AND STATE, AND PUERTO RICO, FOR WEEKS ENDED DECEMBER 10, 1960 AND DECEMBER 9, 1961 - Continued

(By place of occurrence. Numbers under diseases are category numbers of the Seventh Revision of the International Lists, 1955)

00034	Malaria	Meningo infec	ccocal tions	Psitta- cosis	Strepto- coccal sore throat,		yphoid :	fever 040	1	Typhus fever, endemic		es in
Агев	110-117	05	57	096.2	etc. 050,051	49th	49th Week		ative, 9 weeks	101	animals	
A-1	1961	1961	1960	1961	1961	1961	1960	1961	1960	1961	1961	1960
UNITED STATES	1	52	53	25	7,454	10	7	785	776	1	53	64
- CO 3/501 - 3 /						-1 17	- 7	1	247		T SAUSA	51.
NEW ENGLAND	A1.	6	2		314		1750	20	11	1 - 1		
Maine	1.15	<u>_</u>		-	13	1.20	1	1	2	-	-	-
Vermont	-	1	1		1	100			1	-	-	-
Massachusetts	-	2	1	1000	83	-	-	14	5	-	000	9.41
Rhode Island		1		-	42	-	-	2	-	-		100
Connecticut		1.1		1.1	174		-	3	4	- 1	1.3	-
MIDDLE ATLANTIC	-	8	9	1	217	2		98	58		1	4
New York	-	3	7	-	134	1		54	34	- T	1	3
New Jersey		1 4	-		27	-		17	7	-		-
Pennsylvania			2	1	56	1	-	27	17	-	- 124	1
EAST NORTH CENTRAL	-	15	14	1	546	1	1	106	99	-	8	4
Ohio	31.11	4	3		69 98	-	i.	43	29	•	001	1
Illinois	1	5	4		98		1	24 29	25		1	1
Michigan	100	5	3		110	1	COL.	7	21 16	124	15	2
Wisconsin	-		2	1	170			3	8		-	100
WEST NORTH CENTRAL	2.	1		-	1,029	-	-	35	48	_	1.1	
Minnesota	Sant in	10.000			727	-	1.1.1	5	40	1. A 1.	11 2	8
Iowa	-	•	-	-	63	-		2	11		2	4
Missouri	-	1	-	-	19		12190	21	26	11-20-7283	3	
North Dakota	11.00			- 11 <b>-</b> 1/25	92		301.44	1377-1	1	Cassin-2	-	1.11-
South Dakota	100 300	411 BA	101 201	1.16	4	avi ba	20, 090	3	4	and the state	4	1
Kansas	100-07	1.235	10.00		124	-		1 3	3	and south	1.5	-
SOUTH ATLANTIC	GRI MARLENNI	6	4	in the last of	455	1	strack	1.000		to and the second	12000	
Delavare	-	1	-	and the	455	-	-	133	117		3	16
Maryland	-	-	-	-	16	-	-	4	6	Ster Col	10.104	100
District of Columbia	1, 19921 1	10,201	10014	110950	1	10.0	2006_33	13	10	Notes its	199. 200	15-48
Virginia		1	2	211 -21	97	100	10000	20	24			9
West Virginia		1	10.000	10.1°	147	1	-	11	14	a the second	2	4
North Carolina	Citizen 1	2	al of	0.07	22		-	16	9		-	-
Georgia		-	1	-	7 2		-	8 39	12 27	10 - 10 - 10 - 10 - 10 - 10 - 10 -		1
Florida	o Treiligo	1	0.0010	-	158	an dens	1251.21	21	14	744 PRIT	1	- 2
EAST SOUTH CENTRAL	i anima	6	7		1,252	A HELL	1	84	120	aldv sn	12	
Kentucky	mar with	5	5	-	95			19	32	- Mr.	4	6
Tennessee	And and	m Bak	1	-	1,070			50	56	-	6	4
Alabama	-	-	1		42	-	1	11	24	-	2	2
Mississippi		1	-	-	45	-	-	4	8		-	-
EST SOUTH CENTRAL	-	6	3	22	825	3	2	159	205	1	14	25
Arkansas	record vi	1	100	in Éarr	4	1.000 100	BEFAI	30	52	-	3	15
Louisiana		1 2			5			27	59 12		1	•
Oklahoma	Mars at 1	2	3	22	811	3	2	90	82	1	10	10
		and the second second		- 1	1,493		1	64				10
MOUNTAIN	2 2330	1	6	100	1,493	1.2	-	20	46 13		-	-
Montana	E Lopa	in Land	-1 ha	100	143			1	3	2.1	0	-
Wyoming	berne	10 E 13	300	and the	7	-	-	3	4		-	
Colorado	11 Berl	1	10.00		551	1	10-1-	7	1	-	-	-
New Mexico			ACL		363	1000	1	17	13	- 249	-	-
Arizona			4	-	222		_	10	10	-	-	-
Utah			2	-	148 3	-		2 4	2			-
Nevada	1. 1.			12.000	The second	3	2	86	100 TH 2007	5040H (1)	1.0.0	0.000
PACIFIC	1	3	8	1	1,323	3	2	86	72		4	1
Washington	5	ī	1	1000	36			1	8		L. Baser	10
Oregon- California-	1	2	7	1	752	3	2	75	58	in the second	4	1
Alaska	1	1 2 2	i	-	21	101410	2)	This section in	1	fattil	60 x=1	5. 2. 4
Bavaii	-0.0	- 6.2	-	-	50	101-10	11	3			100	1
Puerto Rico		1.000	4			20122223	2	22	20	1 anna		2



The chart shows the number of deaths reported for 117 major cities of the United States by week for the current year, a 5-week moving average of these figures plotted at the central week, and an adjusted average for comparison. For each region the adjusted average was computed as follows: From the total deaths reported each week for the years 1956-1960, 3 central figures were selected by eliminating the highest and lowest figure reported for that week. A 5-week moving average of the arithmetic mean of the 3 central figures was then computed with adjustment to allow for population growth in each region. The average value of the regional increases was 2 percent which was incorporated in the adjusted average shown in the chart.

Table 4 shows the number of death certificates re-

ceived during the week indicated for deaths that occurred in selected cities. Figures compiled in this way, by week of receipt, usually approximate closely the number of deaths occurring during the week. However, differences are to be expected because of variations in the interval between death and receipt of the certificate and because of incomplete reporting due to holidays or vacations. If a report is not received from a city in time to be included in the total for the current week, an estimate is used.

The number of deaths in cities of the same size may also differ because of variations in the age, race, and sex composition of the populations and because some cities are hospital centers serving the surrounding areas. Changes from year to year in the number of deaths may be due in part to population increases or decreases.

				49th week	48th week	Adjusted average,	Percent change, adjusted	Cumulative, first 49 weeks			
	Ares	1 4 4 9 -	2 4 4 5	ended         ended           Dec.         Dec.           9,         2,           1961         1961		49th veek 1956-60	average to current week	1961	1960	Percent	
TOTAL, 117 REPO	ORTING CITIES			12,312	13,157	11,690	+5.3	561,380	562,959	-0.3	
New England			4 cities)	808*	838	701	+15.3	34,627	35,426	-2.3	
Middle Atlantic	************		O cities)	3,333*	3,601	3,161	+5.4	159,119	156,535	+1.7	
East North Central	**************		l cities)	2,518	2,766	2,487	+1.2	120,019	121,861	-1.5	
West North Central			9 cities)	811	954	845	-4.0	38,346	39,064	-1.8	
Bouth Atlantic			l cities)	1,080*	1,151	993	+8.8	48,481	48,211	+0.6	
East South Central			8 cities)	628	613	526	+19.4	25,510	25,498	+0.0	
lest South Central			3 cities)	1,015	1,216	1,076	-5.7	47,912	48,895	-2.0	
Mountain		(	8 cities)	411	403	370	+11.1	17,924	17,697	+1.3	
Pacific		1.	3 cities)	1,708*	1,615	1,531	+11.6	69,442	69,772	-0.5	

(By place of occurrence and week of filing certificate. Excludes fetal deaths. Data exclude figures shown in parentheses in table 4)

6

\*Includes estimate for missing reports.

### Table 4. DEATHS IN SELECTED CITIES

(By place of occurrence and week of filing certificate. Excludes fetal deaths)

Area	49th week ended Dec.	d ended		ative, 9 weeks	Area	49th week ended Dec.	48th week ended Dec.	Cumula first 49	weeks
	9, 1961	2, 1961	1961	1960	of Separation research barner	9, 1961	2, 1961	1961	1960
anotal estenation provide	1501	1501		1500					
NEW ENGLAND:	2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1				WEST NORTH CENTRALCon.:	11-12-1	1951 BO		
Boston, Mass	279	318	11,907	12,283	St. Louis, Mo	260	302	11,586	11,924
Bridgeport, Conn	46*	42	1,919	1,999	St. Paul, Minn	55	87	3,230	3,373
Cambridge, Mass	39	44	1,448	1,495	Wichita, Kans	51	48	2,299	2,272
Fall River, Mass	34	18	1,347	1,382	terra reprised from the second	12311	inerviti.	and states	
Hartford, Conn	51	51	2,390	2,435	SOUTH ATLANTIC:	ACCURATE ON TAXABLE	1		
Lowell, Mass	27	23	1,210	1,160	Atlanta, Ga	123	114	5,548	5,755
Lynn, Mass	35	22	1,087	1,192	Baltimore, Md	240	294	11,999	12,303
New Bedford, Mass	20 54	30 62	1,259	1,231	Charlotte, N.C	56 65	54 70	1,794 2,841	1,892
Providence, R.I.	75	62	2,210 3,060	2,193	Miami, Fla.	69	104	3,702	2,884
Somerville, Mass	19	15	647	3,147 652	Norfolk, Va	44*	50	2,411	3,476
Springfield, Mass	37	54	2,160	2,207	Richmond, Va	87	88	3,818	3,809
Waterbury, Conn	30	39	1,300	1,367	Savannah, Ga	38	50	1,604	1,641
Worcester, Mass	62	58	2,683	2,683	St. Petersburg, Fla	(61)	(76)	(3,273)	(3,469
southerness about the frame	12.1			_,	Tampa, Fla	103	63	3,201	3,169
MIDDLE ATLANTIC:			and the second second		Washington, D.C	217	210	9,623	9,452
Albany, N.Y	42	44	2,259	2,133	Wilmington, Del	38	54	1,940	1,888
Allentown, Pa	38	43	1,680	1,714	and the grant is a first				,
Buffalo, N.Y	156	150	7,095	7,074	EAST SOUTH CENTRAL:	0.028	patrols	- Creation	
Camden, N.J	51	42	2,071	2,056	Birmingham, Ala	102	102	4,216	4,173
Elizabeth, N.J	27	30	1,446	1,425	Chattanooga, Tenn	45	58	2,274	2,319
Erie, Pa	40*	56	1,913	1,902	Knoxville, Tenn	26	26	1,359	1,370
Jersey City, N.J	66	99	3,401	3,424	Louisville, Ky	165	131	5,591	5,546
Newark, N.J.	90	169	4,994	4,754	Memphis, Tenn	131	140	5,589	5,469
New York City, N.Y	1,718	1,799	81,244	79,940	Mobile, Ala	53	47	1,981	2,030
Paterson, N.J.	33	44	1,901	1,903	Montgomery, Ala	42	38	1,585	1,679
Philadelphia, Pa	515	447	24,599	23,701	Nashville, Tenn	64	71	2,915	2,912
Pittsburgh, Pa	200	274	9,376	9,468	WEST SOUTH CENTRAL:	199.2.2.2	21 7-0473	ibeau until T	1.4 500
Rochester, N.Y	26 101	28	1,144	1,164	Austin, Tex	42	47	1,657	1,639
Schenectady, N.Y	28	106 22	4,963	4,941	Baton Rouge, La.	25	64	1,379	1,406
Scranton, Pa	47	33	1,185 1,730	1,151	Corpus Christi, Tex	17	28	1,076	1,126
Syracuse, N.Y	60	68	3,019	1,878 3,063	Dallas, Tex	142	109	6,079	6,099
Trenton, N.J.	46	- 68	2,199	1,997	El Paso, Tex	46	55	1,738	1,853
Utica, N.Y	25	49	1,406	1,329	Fort Worth, Tex	62	88	3,171	3,243
Yonkers, N.Y	24	30	1,494	1,518	Houston, Tex	137	257	8,070	8,137
2 2 2 2 3 4 4	10		-,	-,510	Little Rock, Ark	44	72	2,743	2,789
EAST NORTH CENTRAL:	24		Allow The		New Orleans, La	185	157	8,217	8,742
Akron, Ohio	50	64	2,789	2,776	Oklahoma City, Okla	76	106	3,636	3,681
Canton, Ohio	49	32	1,565	1,692	San Antonio, Tex	117	122	4,985	4,888
Chicago, Ill	725	788	36,256	37,582	Shreveport, La	67	44	2,492	2,621
Cincinnati, Ohio	174	177	7,685	7,681	Tulsa, Okla	55	67	2,669	2,671
Cleveland, Ohio	180	252	9,902	10,314					
Columbus, Ohio	119	143	5,633	5,800	MOUNTAIN:	had st	diam's	and the second	
Dayton, Ohio	94	100	3,921	3,730	Albuquerque, N. Mex	28	45	1,558	1,577
Detroit, Mich	313	373	16,255	16,426	Colorado Springs, Colo	20	20	808	818
Evansville, Ind	34	46	1,776	1,821	Denver, Colo	140	115	5,644	5,831
Flint, Mich	48	48	2,089	1,970	Ogden, Utah	10	18	826	803
Fort Wayne, Ind	47	33	1,857	1,795	Phoenix, Ariz	100	109	4,037	3,742
Gary, Ind.	24	54	1,516	1,516	Pueblo, Colo	19	18	825	794
Grand Rapids, Mich	53	33	2,209	2,017	Salt Lake City, Utah	52	52	2,359	2,374
Indianapolis, Ind	165	174	7,028	7,101	Tucson, Ariz	42	26	1,867	1,758
Madison, Wis Milwaukee, Wis	26	50	1,630	1,595	PACIFIC:	26, 1110		1	
Peoria, Ill.	145	117	6,010	6,125	Berkeley, Calif	18	19	827	0.11
Rockford, Ill	24 34	34 26	1,395 1,394	1,482	Fresno, Calif	(57)	(48)	(2,128)	831
South Bend, Ind	34	33	1,394	1,409	Glendale, Calif	(38)	(34)	(1,629)	(2,181)
Toledo, Ohio	114	126	4,858	4,881	Honolulu, Hawaii	50*	42	1,986	2,026
Youngstown, Ohio	69	63	2,850	2,707	Long Beach, Calif	65*	65	2,725	2,695
Tara (S. Paranese and and		2.5	1 5		Los Angeles, Calif	587	567	24,554	24,442
WEST NORTH CENTRAL:		C		1.1	Oakland, Calif	99	116	4,791	4,690
Des Moines, Iowa	78	81	2,683	2,703	Pasadena, Calif	41*	35	1,640	1,696
Duluth, Minn	20	36	1,255	1,241	Portland, Oreg	189	96	5,307	5,401
Kansas City, Kans	46	37	1,847	1,724	Sacramento, Calif	68*	55	3,029	2,872
Kansas City, Mo	128	137	6,278	6,135	San Diego, Calif	108*	118	4,402	4,444
Lincoln, Nebr	(29)	(42)	(1,376)	(1,261)	San Francisco, Calif	235*	236	9,541	9,665
Minneapolis, Minn	109	144	5,792	6,105	San Jose, Calif	()	(33)	()	(1,695
Omaha, Nebr	64	82	3,376	3,587	Seattle, Wash	148	160	6,421	6,670
Culdina, Mebr.			· ·		Spokane, Wash	140	100	0,721	

San Juan, P. R .-----

(25)

(26)

(1,611)

(1,700)

() Figures shown in parenthesis are from cities which have been reporting less than five years and hence are not included in Table 3. **Polio** – Fifteen cases of poliomyelitis, 10 paralytic, were reported for the 49th week ending December 8. This is essentially unchanged from the 15 cases, 11 paralytic, reported the preceding week.

**Hepatitis** – The 1,367 cases of hepatitis reported for the 49th week are 74 more than reported for the 48th week. This represents 208 more cases than were reported for the 49th week of 1960. This continued increase in reported cases indicates the seasonal rise characteristic of hepatitis. Significant rises were reported from the Mountain and East North Central States. Only slight increases were noted in all other regions with the exception of the East South Central States which reported a slight decrease.

## International Notes – Quarantine Measures Cholera and Cholera-like Disease

Unofficially reported are over two thousand cases and three hundred deaths due to a cholera-like disease which have occurred to date in the Philippine Islands. The disease appears to be spreading.

All persons entering the United States within five days after leaving infested areas and the Philippine Islands will be required to present a valid certificate of immunization against cholera.

The Division of Foreign Quarantine recommends that all persons traveling to the Far East receive cholera vaccine.

#### Yellow Fever Vaccination Centers

The following name should be added to the list of Yellow Fever Vaccination Centers in Section 6:

City	Center	Clinic Hours Fee
Virginia	Bureau of Public Health	Friday, 1-4:30 pm Yes
Lynchburg	701 Hollins Street	by appointment
	Tel: Victor 6-5273	and another with the second

### Immunization Information for International Travel

Correction of Public Health Service Publication #384, Section 5, page 53.

France - under recommendations by USPHS, delete "T" which refers to Typhus.

#### SOURCE AND NATURE OF MORBIDITY DATA

These provisional data are based on reports to the Public Health Service from the health departments of each State and Puerto Rico. They give the total number of cases of certain communicable diseases reported during the week usually ended the preceding Saturday. Total figures for the United States and the Pacific Division include data for the States of Alaska and Hawaii. Cases of anthrax, botulism, and rabies in man are not shown in table 2, but a footnote to table 1 shows the States reporting these diseases. When diseases of rare occurrence are reported by a State (cholera, dengue, plague, louse-borne relapsing fever, smallpox, louseborne epidemic typhus, and yellow fever) this is noted below table 1.

