SUPPLEMENTARY TABLE. Notes from the Field: Long COVID Prevalence Among Adults — United States, 2022

SUPPLEMENTARY TABI	LE. Notes fr	om the Field: Long	COVID Prev	OVID Prevalence Among Adults — United States, 2022			
					Don't know, refused to		
	Ever had Long COVID		Never had Long COVID		answer, or missing		
Jurisdiction	No.*	Row % (95% CI)	No.*	Row % (95% CI)	No.*	Row % (95% CI)	
Total no. of cases	26,783	_	415,415	_	2,934	_	
National	26,783	6.4 (6.2–6.5)	415,415	93.0 (92.8–93.2)	2,934	0.6 (0.6–0.7)	
Alabama	372	9.8 (8.6–11.2)	4,109	89.4 (88.0–90.7)	25	0.7 (0.5–1.2)	
Alaska	345	6.6 (5.7–7.5)	5,488	92.8 (91.8–93.7)	32	0.6 (0.4–1)	
Arizona	638	7.1 (6.3–8.0)	9,483	92.2 (91.2–93.0)	64	0.7 (0.5–1)	
Arkansas	337	6.8 (6.0–7.7)	4,939	92.5 (91.5–93.4)	33	0.7 (0.5–1)	
California	574	5.5 (4.8–6.1)	10,305	94.0 (93.3–94.6)	73	0.6 (0.4–0.8)	
Colorado	583	6.4 (5.8–7.0)	8,721	92.9 (92.2–93.5)	61	0.7 (0.5-1)	
Connecticut	519	5.2 (4.6–5.9)	9,198	94.0 (93.2–94.6)	67	0.8 (0.5-1.2)	
Delaware	210	5.9 (4.8-7.2)	3,749	93.3 (91.9–94.5)	28	0.8 (0.4-1.3)	
District of Columbia	93	3.5 (2.7-4.4)	3,131	96.2 (95.2–97.0)	13	0.3 (0.2-0.6)	
Florida	805	5.9 (5.1-6.8)	12,529	93.7 (92.8–94.5)	59	0.4 (0.2-0.6)	
Georgia	567	6.2 (5.5-7.0)	8,618	93.2 (92.4–93.9)	51	0.6 (0.4-0.8)	
Guam	83	3.5 (2.6-4.9)	2,175	96.1 (94.7–97.2)	8	0.3 (0.1–0.9)	
Hawaii	274	3.8 (3.3–4.5)	7,449	95.9 (95.2–96.5)	24	0.3 (0.2–0.5)	
Idaho	511	8.3 (7.5–9.2)	5,714	90.9 (89.9–91.7)	55	0.8 (0.6–1.1)	
Illinois	261	6.4 (5.6–7.4)	3,776	93.2 (92.2–94.1)	19	0.4 (0.2–0.6)	
Indiana	639	6.5 (5.9–7.1)	9,765	93.0 (92.3–93.5)	62	0.6 (0.4–0.8)	
Iowa	597	7.1 (6.4–7.9)	8,308	92.3 (91.5–93.0)	44	0.6 (0.4–0.9)	
Kansas	755	7.4 (6.7–8.1)	10,399	91.9 (91.1–92.5)	93	0.8 (0.6–1.0)	
Kentucky	303	7.5 (6.4–8.7)	3,685	91.6 (90.3–92.8)	35	0.9 (0.5–1.5)	
Louisiana	432	7.9 (7.0–8.9)	5,167	91.6 (90.5–92.5)	30	0.5 (0.3–0.9)	
Maine	472	5.3 (4.7–6.0)	10,102	94.1 (93.3–94.7)	72	0.6 (0.5–0.9)	
Maryland	678	4.3 (3.8–4.9)	15,639	95.1 (94.5–95.6)	101	0.6 (0.4–0.8)	
Massachusetts	584	5.3 (4.8–5.9)	10,353	93.7 (93.1–94.3)	92	1.0 (0.7–1.3)	
Michigan	533	6.3 (5.6–6.9)	9,461	93.1 (92.4–93.8)	64	0.6 (0.4–0.8)	
Minnesota	960	6.2 (5.7–6.8)	15,731	93.0 (92.5–93.6)	130	0.7 (0.6–0.9)	
Mississippi	375	8.7 (7.7–9.8)	3,836	90.6 (89.4–91.7)	28	0.7 (0.4–1.2)	
Missouri	560	7.9 (7.1–8.7)	6,818	91.5 (90.6–92.3)	60	0.7 (0.5–1.0)	
Montana	581	9.8 (8.8–10.8)	6,402	89.3 (88.3–90.3)	65	0.9 (0.7–1.2)	
Nebraska	514	7.5 (6.7–8.4)	6,933	92.1 (91.2–92.9)	26	0.4 (0.2–0.7)	
Nevada	203	7.3 (6.0–8.8)	2,964	92.3 (90.9–93.6)	21	0.4 (0.2–0.6)	
New Hampshire	253	5.0 (4.2–5.9)	6,467	94.5 (93.6–95.3)	37	0.5 (0.3–0.8)	
New Jersey	438	5.7 (5.0–6.4)	7,726	93.9 (93.1–94.6)	45	0.4 (0.3–0.6)	
New Mexico	307	7.4 (6.4–8.7)	4,419	92.1 (90.8–93.1)	32	0.5 (0.3–0.8)	
New York	945	5.4 (4.9–5.9)	16,751	94.1 (93.5–94.6)	104	0.5 (0.4–0.6)	
North Carolina	313	6.5 (5.6–7.4)	4,170	93.1 (92.1–93.9)	22	0.5 (0.4–0.8)	
North Dakota	335		3,786	90.0 (88.8–91.1)	32	0.7 (0.4–1.0)	
Ohio		9.3 (8.3–10.5)	1	92.1 (91.5–92.7)		` '	
	1,194	7.1 (6.6–7.7)	15,170	,	123	0.7 (0.6–1.0)	
Oklahoma	525 271	9.1 (8.3–10.0)	5,192	90.1 (89.1–90.9)	58 52	0.8 (0.6–1.1)	
Oregon	271	5.3 (4.6–6.1)	5,433	93.8 (92.9–94.5)	52	0.9 (0.7–1.3)	
Pennsylvania	308	6.9 (5.9–8.1)	4,239	92.4 (91.2–93.5)	35	0.6 (0.4–0.9)	
Puerto Rico	356	6.5 (5.6–7.5)	5,129	92.9 (91.9–93.8)	24	0.6 (0.3–1.0)	
Rhode Island	272	4.8 (4.1–5.7)	5,575	94.5 (93.6–95.3)	46	0.7 (0.5–1.0)	
South Carolina	582	6.2 (5.5–6.8)	9,402	93.4 (92.7–94.1)	53	0.4 (0.3–0.6)	
South Dakota	618	8.3 (6.1–11.2)	6,738	90.3 (87.3–92.6)	68	1.4 (0.8–2.6)	
Tennessee	458	8.9 (7.9–9.9)	4,763	90.2 (89.1–91.3)	45	0.9 (0.6–1.3)	

Texas	904	6.5 (5.8–7.2)	13,256	92.8 (92.0–93.6)	85	0.7 (0.4–1.2)
U.S. Virgin Islands	39	1.9 (0.9-4.1)	1,487	97.8 (95.6–99.0)	5	0.3 (0.0-1.7)
Utah	784	8.3 (7.6–9.0)	8,976	91.1 (90.3–91.7)	66	0.6 (0.5-0.9)
Vermont	313	4.2 (3.5-4.9)	8,434	95.2 (94.4–95.8)	64	0.7 (0.5-0.9)
Virginia	547	5.9 (5.2-6.6)	9,876	93.4 (92.6–94.0)	84	0.7 (0.5-1.0)
Washington	1,238	5.2 (4.9-5.6)	24,753	94.1 (93.7–94.5)	161	0.6 (0.5–0.8)
West Virginia	455	10.6 (9.5-11.8)	4,471	88.4 (87.1–89.5)	55	1.0 (0.7–1.5)
Wisconsin	649	6.0 (5.5–6.7)	10,548	93.3 (92.6–93.9)	79	0.7 (0.5–0.9)
Wyoming	321	9.0 (7.9-10.3)	3,797	90.4 (89.1–91.5)	24	0.6 (0.3-1.0)

^{*} Numbers are unweighted. Prevalence estimates are standardized to the 2020 U.S. Census population, weighted, and account for complex sampling. Estimates were standardized to the U.S civilian, noninstitutionalized population. Sex-specific weights by age group were applied for ages 18–44 years, 45–64 years, and ≥65 years.