

**SUPPLEMENTARY TABLE. Incidence rate\* ratios of coccidioidomycosis, histoplasmosis, and blastomycosis, by selected patient characteristics — United States, 2019–2021**

Characteristic	Coccidioidomycosis <sup>§</sup>			Histoplasmosis <sup>¶</sup>			Blastomycosis <sup>**</sup>		
	2019	2020	2021	2019	2020	2021	2019	2020	2021
<b>Sex</b>									
Female	ref	ref	ref	ref	ref	ref	ref	ref	ref
Male	1.1 <sup>†</sup>	1.2 <sup>†</sup>	1.2 <sup>†</sup>	1.3 <sup>†</sup>	1.4 <sup>†</sup>	1.4 <sup>†</sup>	2.4 <sup>†</sup>	1.8 <sup>†</sup>	2.7 <sup>†</sup>
<b>Age group (yrs)</b>									
<5	<0.1 <sup>†</sup>	<0.1 <sup>†</sup>	<0.1 <sup>†</sup>	0.1 <sup>†</sup>	0.1 <sup>†</sup>	0.1 <sup>†</sup>	0.1 <sup>†</sup>	0.0	<0.1 <sup>†</sup>
5–19	0.3 <sup>†</sup>	0.2 <sup>†</sup>	0.2 <sup>†</sup>	0.4 <sup>†</sup>	0.5 <sup>†</sup>	0.5 <sup>†</sup>	0.4 <sup>†</sup>	0.3 <sup>†</sup>	0.3 <sup>†</sup>
20–39	0.7 <sup>†</sup>	0.6 <sup>†</sup>	0.6 <sup>†</sup>	0.7 <sup>†</sup>	0.8 <sup>†</sup>	0.8 <sup>†</sup>	1.0	0.6	0.6 <sup>†</sup>
<b>40–64</b>	<b>ref</b>	<b>ref</b>	<b>ref</b>	<b>ref</b>	<b>ref</b>	<b>ref</b>	<b>ref</b>	<b>ref</b>	<b>ref</b>
65–79	1.4 <sup>†</sup>	1.5 <sup>†</sup>	1.5 <sup>†</sup>	1.1	1.1	1.2 <sup>†</sup>	1.1	1.2	1.1
≥80	1.3 <sup>†</sup>	1.6 <sup>†</sup>	1.5 <sup>†</sup>	0.6 <sup>†</sup>	0.6 <sup>†</sup>	0.5 <sup>†</sup>	0.5	0.6	0.6
<b>Race and ethnicity</b>									
<b>White, NH</b>	<b>ref</b>	<b>ref</b>	<b>ref</b>	<b>ref</b>	<b>ref</b>	<b>ref</b>	<b>ref</b>	<b>ref</b>	<b>ref</b>
AI/AN, NH	4.2 <sup>†</sup>	4.4 <sup>†</sup>	4.2 <sup>†</sup>	0.9	1.2	1.5	6.4 <sup>†</sup>	7.8 <sup>†</sup>	7.4 <sup>†</sup>
Asian and NH/OPI, NH	1.1 <sup>†</sup>	1.0	0.9 <sup>†</sup>	0.8	0.5 <sup>†</sup>	0.5 <sup>†</sup>	2.3	1.0	1.9 <sup>†</sup>
Black or African American, NH	1.0	1.0	1.0	0.7 <sup>†</sup>	0.8 <sup>†</sup>	0.8 <sup>†</sup>	0.9	0.7	0.8
Hispanic or Latino (all races)	2.7 <sup>†</sup>	2.1 <sup>†</sup>	2.2 <sup>†</sup>	0.9	0.6 <sup>†</sup>	0.6 <sup>†</sup>	1.3	1.0	1.3
Other <sup>††</sup>	1.1	1.5 <sup>†</sup>	1.4 <sup>†</sup>	1.5 <sup>†</sup>	0.7 <sup>†</sup>	0.8	0.3	0.1	0.1
<b>Total</b>	<b>NA</b>	<b>0.9<sup>†</sup></b>	<b>1.1<sup>†</sup></b>	<b>NA</b>	<b>0.9</b>	<b>1.4<sup>†</sup></b>	<b>NA</b>	<b>1.0</b>	<b>1.0</b>

**Abbreviations:** AI/AN = American Indian or Alaska Native; NH = non-Hispanic; NH/OPI = Native Hawaiian or other Pacific Islander; ref = reference group; NA = not applicable.

\*Cases per 100,000 population. Summed state-specific denominators from 2019–2021 U.S. Census Bureau data were used to calculate incidence for the total grouping for each year.

<sup>†</sup>Statistically significant rate ratios based on a *p*-value of <0.05.

<sup>§</sup>Sex was missing for 68 cases, age was missing for 65 cases, and race and ethnicity were missing for 20,749 cases.

<sup>¶</sup>Sex was missing for five cases, age was missing for five cases, race and ethnicity were missing for 497 cases.

\*\*Race and ethnicity were missing for 48 cases.

<sup>††</sup>Calculated on basis of two or more reported race categories.