



Influenza (Flu) (</flu/index.htm>)

Weekly U.S. Influenza Surveillance Report

Updated March 1, 2024



A Weekly Influenza Surveillance Report Prepared by the Influenza Division

Key Updates for Week 8, ending February 24, 2024

Seasonal influenza activity remains elevated nationally with increases in some parts of the country.

Viruses

Clinical Lab

14.2% (Trend ↓)
positive for influenza
this week

[\(/flu/weekly/index.htm#ClinicalLaboratories\)](/flu/weekly/index.htm#ClinicalLaboratories)

Public Health Lab

The most frequently reported
influenza viruses this week were
influenza B/Victoria.

[\(/flu/weekly/index.htm#PublicHealthLaboratories\)](/flu/weekly/index.htm#PublicHealthLaboratories)

Virus Characterization

Genetic and antigenic
characterization and antiviral
susceptibility are summarized in
this report.

[\(/flu/weekly/index.htm#VirusCharacterization\)](/flu/weekly/index.htm#VirusCharacterization)

Illness

Outpatient Respiratory Illness

4.4% (Trend →)
of visits to a health care provider this week were for respiratory illness
(above baseline).

[\(/flu/weekly/index.htm#ILINet\)](/flu/weekly/index.htm#ILINet)

Outpatient Respiratory Illness: Activity Map

This week 12 jurisdictions experienced moderate activity and 27 jurisdictions experienced high or very high activity.

[\(/flu/weekly/index.htm#ORIAM\)](/flu/weekly/index.htm#ORIAM)

FluSurv-NET

61.5 per 100,000
cumulative hospitalization rate.

[\(/flu/weekly/index.htm#FluSurvNet\)](/flu/weekly/index.htm#FluSurvNet)

NHSN Hospitalizations

10,148 (Trend →)
patients admitted to hospitals with influenza this week.

[\(/flu/weekly/index.htm#NHSN\)](/flu/weekly/index.htm#NHSN)

NCHS Mortality

0.7% (Trend →)
of deaths attributed to influenza this week.

[\(/flu/weekly/index.htm#NCHSMortality\)](/flu/weekly/index.htm#NCHSMortality)

Pediatric Deaths

2
influenza-associated deaths were reported this week for a total of 93 deaths this season.

[\(/flu/weekly/index.htm#PedMortality\)](/flu/weekly/index.htm#PedMortality)

All data are preliminary and may change as more reports are received.

Directional arrows indicate changes between the current week and the previous week. [Additional information \(/flu/weekly/index.htm#Trends\)](/flu/weekly/index.htm#Trends) on the arrows can be found at the bottom of this page.

A description of the CDC influenza surveillance system, including methodology and detailed descriptions of each data component is available on the [surveillance methods \(/flu/weekly/overview.htm\)](/flu/weekly/overview.htm) page.

Additional information on the current and previous influenza seasons for each surveillance component are available on [FluView Interactive \(/flu/weekly/fluviewinteractive.htm\)](/flu/weekly/fluviewinteractive.htm).

Key Points

- Seasonal influenza activity remains elevated nationally with increases in some parts of the country.
- Nationally, percent positivity for influenza decreased slightly overall with percent positivity for influenza A decreasing and percent positivity for influenza B remaining stable. Trends in percent positivity for influenza A and B varied by region.
- Outpatient respiratory illness has been above baseline¹ nationally since November and is above baseline in all 10 HHS regions.
- The number of weekly flu hospital admissions remained stable compared to last week.
- During Week 8, of the 773 viruses reported by public health laboratories, 508 (65.7%) were influenza A and 265 (34.3%) were influenza B. Of the 338 influenza A viruses subtyped during Week 8, 186 (55.0%) were influenza A(H1N1) and 152 (45.0%) were A(H3N2).
- Two influenza-associated pediatric deaths were reported during Week 8, bringing the 2023-2024 season total to 93 pediatric deaths.
- CDC estimates that there have been at least 26 million illnesses, 290,000 hospitalizations, and 18,000 deaths from flu so far this season.
- CDC recommends that everyone 6 months and older get an annual flu vaccine as long as influenza viruses are spreading.² Vaccination can still provide benefit this season.
- There also are prescription flu antiviral drugs that can treat flu illness; those should be started as early as possible and are especially important for higher risk patients.³
- Flu viruses are among several viruses contributing to respiratory disease activity. CDC is providing updated, integrated information (<https://www.cdc.gov/respiratory-viruses/index.html>) about COVID-19, flu, and RSV activity on a weekly basis.

U.S. Virologic Surveillance

(<https://www.cdc.gov/flu/weekly/overview.htm#LabSurveillance>)

Nationally, the percentage of respiratory specimens testing positive for influenza in clinical laboratories decreased (change of ≤ 0.5 percentage points) compared to the previous week, but trends varied by region. Region 7 reported an increase, regions 1, 3, 5, 6, 8 and 9 reported a decrease, and regions 2, 4, and 10 remained stable during Week 8 compared to Week 7. The regions with the highest percent positivity were regions 7 (28.4%), 5 (19.4%), and 6 (18.3). Since Week 40, influenza A(H1N1)pdm09 has been the predominant virus circulating in all regions. However, the distribution of circulating viruses varies by region. For regional and state level data and age group distribution, please visit [FluView Interactive](https://gis.cdc.gov/grasp/fluview/fluportaldashboard.html) (<https://gis.cdc.gov/grasp/fluview/fluportaldashboard.html>). Viruses known to be associated with recent receipt of live attenuated influenza vaccine (LAIV) or found upon further testing to be a vaccine virus are not included, as they are not circulating influenza viruses.

Clinical Laboratories

The results of tests performed by clinical laboratories nationwide are summarized below. Data from clinical laboratories (the percentage of specimens tested that are positive for influenza virus) are used to monitor whether influenza activity is increasing or decreasing.

	Week 8	Data Cumulative since October 1, 2023 (Week 40)
No. of specimens tested	87,299	2,214,809
No. of positive specimens (%)	12,376 (14.2%)	249,106 (11.2%)
<i>Positive specimens by type</i>		
Influenza A	7,298 (59.0%)	185,263 (74.4%)
Influenza B	5,078 (41.0%)	63,833 (25.6%)

<http://gis.cdc.gov/grasp/fluview/fluportaldashboard.html>

[View Chart Data \(/flu/weekly/weeklyarchives2023-2024/data/whoAllregt_cl08.html\)](/flu/weekly/weeklyarchives2023-2024/data/whoAllregt_cl08.html) | [View Full Screen \(/flu/weekly/WeeklyArchives2023-2024/WHONPHL08.html\)](/flu/weekly/WeeklyArchives2023-2024/WHONPHL08.html)

Public Health Laboratories

The results of tests performed by public health laboratories nationwide are summarized below. Data from public health laboratories are used to monitor the proportion of circulating influenza viruses that belong to each influenza subtype/lineage.

	Week 8	Data Cumulative since October 1, 2023 (Week 40)
No. of specimens tested	2,532	80,259
No. of positive specimens	773	25,806

<i>Positive specimens by type/subtype</i>		
Influenza A	508 (65.7%)	20,783 (80.5%)
Subtyping Performed	338 (66.5%)	16,978 (81.7%)
(H1N1)pdm09	186 (55.0%)	12,790 (75.3%)
H3N2	152 (45.0%)	4,188 (24.7%)
H3N2v	0 (0.0%)	0 (0.0%)
Subtyping not performed	170 (33.5%)	3,805 (18.3%)
Influenza B	265 (34.3%)	5,023 (19.5%)
Lineage testing performed	187 (70.6%)	4,108 (81.8%)
Yamagata lineage	0 (0.0%)	0 (0.0%)
Victoria lineage	187 (100%)	4,108 (100%)
Lineage not performed	78 (29.4%)	915 (18.2%)

(<http://gis.cdc.gov/grasp/fluview/fluportaldashboard.html>)

[View Chart Data \(/flu/weekly/weeklyarchives2023-2024/data/whoAllregt_phl08.html\)](/flu/weekly/weeklyarchives2023-2024/data/whoAllregt_phl08.html) | [View Full Screen \(/flu/weekly/weeklyarchives2023-2024/WhoPHL08.html\)](/flu/weekly/weeklyarchives2023-2024/WhoPHL08.html)

Additional virologic surveillance information for current and past seasons:

[Surveillance Methods \(/flu/weekly/overview.htm#LabSurveillance\)](/flu/weekly/overview.htm#LabSurveillance) | [FluView Interactive: National, Regional, and State Data](#)
(<http://gis.cdc.gov/grasp/fluview/fluportaldashboard.html>) or [Age Data \(https://gis.cdc.gov/grasp/fluview/flu_by_age_virus.html\)](https://gis.cdc.gov/grasp/fluview/flu_by_age_virus.html)

Influenza Virus Characterization

(</flu/weekly/overview.htm#VirusCharacterization>)

CDC performs [genetic \(https://www.cdc.gov/flu/about/professionals/genetic-characterization.htm\)](https://www.cdc.gov/flu/about/professionals/genetic-characterization.htm) and [antigenic \(https://www.cdc.gov/flu/about/professionals/antigenic.htm\)](https://www.cdc.gov/flu/about/professionals/antigenic.htm) characterization of U.S. viruses submitted from state and local public health laboratories according to the Right Size Roadmap submission guidance. These data are used to compare how

antibodies raised against a particular virus can often recognize small changes in the surface proteins of other viruses. In HI assays, viruses with similar antigenic properties have antibody titer differences of less than or equal to 4-fold when compared to the reference (vaccine) virus. In HINT, viruses with similar antigenic properties have antibody neutralization titer differences of less than or equal to 8-fold. Viruses selected for antigenic characterization are a subset representing the genetic changes in the surface proteins seen in genetically characterized viruses.

Influenza A Viruses

- **A (H1N1)pdm09:** 172 A(H1N1)pdm09 viruses were antigenically characterized by HI, and all were well-recognized (reacting at titers that were within 4-fold of the homologous virus titer) by ferret antisera to cell-grown A/Wisconsin/67/2022-like reference viruses representing the A(H1N1)pdm09 component for the cell- and recombinant-based influenza vaccines.
- **A (H3N2):** 161 A(H3N2) viruses were antigenically characterized by HI or HINT, and 159 (99%) were well-recognized (reacting at titers that were within 4-fold of the homologous virus titer in HI or reacting at titers that were less than or equal to 8-fold of the homologous virus in HINT) by ferret antisera to cell-grown A/Darwin/6/2021-like reference viruses representing the A(H3N2) component for the cell- and recombinant-based influenza vaccines.

Influenza B Viruses

- **B/Victoria:** One hundred influenza B/Victoria-lineage virus were antigenically characterized by HI, and all were well-recognized (reacting at titers that were within 4-fold of the homologous virus titer) by ferret antisera to cell-grown B/Austria/1359417/2021-like reference viruses representing the B/Victoria component for the cell- and recombinant-based influenza vaccines.
- **B/Yamagata:** No influenza B/Yamagata-lineage viruses were available for antigenic characterization.

Assessment of Virus Susceptibility to Antiviral Medications

CDC assesses susceptibility of influenza viruses to the antiviral medications including the neuraminidase inhibitors (oseltamivir, zanamivir, and peramivir) and the PA endonuclease inhibitor baloxavir using next generation sequence analysis supplemented by laboratory assays. Information about antiviral susceptibility test methods can be found at [U.S. Influenza Surveillance: Purpose and \(https://www.cdc.gov/flu/weekly/overview.htm\) Methods | CDC \(https://www.cdc.gov/flu/weekly/overview.htm\)](https://www.cdc.gov/flu/weekly/overview.htm).

Viruses collected in the U.S. since October 01, 2023, were tested for antiviral susceptibility as follows:

Antiviral Medication			Total Viruses	A/H1	A/H3	B/Victoria
Neuraminidase Inhibitors	Oseltamivir	Viruses Tested	2,229	998	640	591
		Reduced Inhibition	1 (0.4%)	1 (0.1%)	0 (0.0%)	0 (0.0%)
		Highly Reduced Inhibition	1 (0.4%)	1 (0.1%)	0 (0.0%)	0 (0.0%)
	Peramivir	Viruses Tested	2,229	998	640	591
		Reduced Inhibition	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
		Highly Reduced Inhibition	1 (0.4%)	1 (0.1%)	0 (0.0%)	0 (0.0%)
	Zanamivir	Viruses Tested	2,229	998	640	591
		Reduced Inhibition	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
		Highly Reduced Inhibition	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
PA Cap-Dependent Endonuclease Inhibitor	Baloxavir	Viruses Tested	2,162	970	623	569
		Decreased Susceptibility	1 (0.5%)	0 (0.0%)	1 (0.2%)	0 (0.0%)

One A(H1N1)pdm09 virus had NA-H275Y amino acid substitution and showed highly reduced inhibition by oseltamivir and peramivir. One (H1N1)pdm09 virus had NA-I223V and NA-S247N amino acid substitutions and showed reduced inhibition by oseltamivir.

One A(H3N2) virus had PA-I38T amino acid substitution and showed reduced susceptibility to baloxavir.

High levels of resistance to the adamantanes (amantadine and rimantadine) persist among influenza A(H1N1)pdm09 and influenza A(H3N2) viruses (the adamantanes are not effective against influenza B viruses). Therefore, use of these antivirals for treatment and prevention of influenza A virus infection is not recommended and data from adamantane resistance testing are not presented.

Outpatient Respiratory Illness Surveillance

(<https://www.cdc.gov/flu/weekly/overview.htm#ILINet>)

The U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet) monitors outpatient visits for respiratory illness referred to as influenza-like illness [ILI (fever plus cough or sore throat)], not laboratory-confirmed influenza and will therefore capture respiratory illness visits due to infection with pathogens that can present with similar symptoms, including influenza viruses, SARS-CoV-2, and RSV. It is important to evaluate syndromic surveillance data, including that from ILINet, in the context of other sources of surveillance data to obtain a more complete and accurate picture of influenza, SARS-CoV-2, and other respiratory virus activity. CDC is providing integrated information about COVID-19, influenza, and RSV activity on a [website \(https://www.cdc.gov/respiratory-viruses/index.html\)](https://www.cdc.gov/respiratory-viruses/index.html) that is updated weekly. Information about other respiratory virus activity can be found on [CDC's National Respiratory and Enteric Virus Surveillance System \(NREVSS\) website \(https://www.cdc.gov/surveillance/nrevss/index.html\)](https://www.cdc.gov/surveillance/nrevss/index.html).

Outpatient Respiratory Illness Visits

Nationwide, during Week 8, 4.4% of patient visits reported through ILINet were due to respiratory illness that included fever plus a cough or sore throat, also referred to as ILI. This has remained stable (change of ≤ 0.1 percentage points) compared to Week 7. The percentage of visits for ILI remained stable in regions 1, 2, 3, and 4, increased in regions 5, 7, and 9, and decreased in regions 6, 8, and 10 in Week 8 compared to Week 7. All regions remain above their region-specific baselines this week. Multiple respiratory viruses are co-circulating, and the relative contribution of influenza virus infection to ILI varies by location.

Outpatient Respiratory Illness Visits by Age Group

About 70% of ILINet participants provide both the number of patient visits for respiratory illness and the total number of patient visits for the week broken out by age group. Data from this subset of providers are used to calculate the percentages of patient visits for respiratory illness by age group.

The percentage of visits for respiratory illness reported in ILINet remained stable for all age groups during Week 8 compared to Week 7.

Outpatient Respiratory Illness Activity Map

Data collected in ILINet are used to produce a measure of ILI activity* (</flu/weekly/overview.htm#ILINet>) by state/jurisdiction and Core Based Statistical Areas (CBSA).

	Number of Jurisdictions	Number of CBSAs
--	--------------------------------	------------------------

Activity Level	Week 8 (Week ending Feb. 24, 2024)	Week 7 (Week ending Feb. 17, 2024)	Week 8 (Week ending Feb. 24, 2024)	Week 7 (Week ending Feb. 17, 2024)
Very High	5	7	16	14
High	22	20	109	108
Moderate	12	11	116	123
Low	7	10	210	227
Minimal	8	7	250	235
Insufficient Data	1	0	228	222

*Data collected in ILINet may disproportionately represent certain populations within a jurisdiction or CBSA, and therefore, may not accurately depict the full picture of influenza activity for the entire jurisdiction or CBSA. Differences in the data presented here by CDC and independently by some health departments likely represent differing levels of data completeness with data presented by the health department likely being the more complete.

Additional information about medically attended visits for ILI for current and past seasons:
[Surveillance Methods \(/flu/weekly/overview.htm#ILINet\)](#) | [FluView Interactive: National, Regional, and State Data \(http://gis.cdc.gov/grasp/fluview/fluportaldashboard.html\)](#) or [ILI Activity Map \(https://gis.cdc.gov/grasp/fluview/main.html\)](#)

Hospitalization Surveillance

Hospitalization Surveillance

(<http://www.cdc.gov/flu/weekly/overview.htm#HospitalizationSurv>)

FluSurv-NET

The Influenza Hospitalization Surveillance Network (FluSurv-NET) conducts population-based surveillance for laboratory-confirmed influenza-related hospitalizations in select counties in 14 states and represents approximately 9% of the U.S. population. FluSurv-NET hospitalization data are preliminary. As data are received each week, prior case counts and rates are updated accordingly.

A total of 18,790 laboratory-confirmed influenza-associated hospitalizations were reported by FluSurv-NET sites between October 1, 2023, and February 24, 2024. The weekly hospitalization rate observed in Week 8 was 2.6 per 100,000 population. The weekly hospitalization rate observed during Week 52 is the third highest peak weekly rate observed during all seasons going back to 2010-2011 following the 2014-2015 and 2017-2018 seasons. The overall cumulative hospitalization rate was 61.5 per 100,000 population. This cumulative hospitalization rate is the second highest cumulative hospitalization rate when compared against previous end-of-season rates for Week 8, and it is the second highest cumulative in-season hospitalization rate observed in Week 8, following the 2017-2018 season (81.7). Cumulative in-season hospitalization rates observed in Week 8 from 2010-2011 through 2021-2022 ranged from 0.7 to 59.9.

When examining rates by age, the highest cumulative hospitalization rate per 100,000 population was among adults aged 65 years and older (166.1), followed by adults aged 50-64 years (74.8) and children aged 0-4 years (63.6). When examining age-adjusted rates by race and ethnicity, the highest rate of hospitalization per 100,000 population was among non-Hispanic Black persons (119.1), followed by non-Hispanic American Indian or Alaska Native persons (86.5), Hispanic persons (58.8), non-Hispanic White persons (46.2), and non-Hispanic Asian/Pacific Islander persons (33.5).

Among 18,790 hospitalizations, 16,351 (87.0%) were associated with influenza A virus, 2,307 (12.3%) with influenza B virus, 35 (0.2%) with influenza A virus and influenza B virus co-infection, and 97 (0.5%) with influenza virus for which the type was not determined. Among those with influenza A subtype information, 2,763 (74.3%) were A(H1N1) pdm09 and 954 (25.7%) were A(H3N2).

Among 2,052 hospitalized adults with information on underlying medical conditions, 95.6% had at least one reported underlying medical condition; the most commonly reported were hypertension, cardiovascular disease, obesity, and metabolic disease. Among 1,079 hospitalized women of childbearing age (15-49 years) with information on pregnancy status, 23.8% were pregnant. Among 571 hospitalized children with information on underlying medical conditions, 69.3% had at least one reported underlying medical condition; the most commonly reported was asthma, followed by obesity and neurologic disease.

<https://gis.cdc.gov/grasp/fluview/FluHospRates.html>

[View Full Screen \(/flu/weekly/weeklyarchives2023-2024/EIPRates08.html\)](/flu/weekly/weeklyarchives2023-2024/EIPRates08.html)

In this figure, cumulative rates for all seasons prior to the 2023-2024 season reflect end-of-season rates. For the 2023-2024 season, rates for recent hospital admissions are subject to reporting delays. As hospitalization data are reviewed each week, prior case counts and rates are updated accordingly.

(<https://gis.cdc.gov/grasp/fluview/FluHospChars.html>)

[View Full Screen \(/flu/weekly/weeklyarchives2023-2024/EIPConditions08.html\)](/flu/weekly/weeklyarchives2023-2024/EIPConditions08.html)

In this figure, weekly rates for all seasons prior to the 2023-24 season reflect end-of-season rates. For the 2023-24 season, rates for recent hospital admissions are subject to reporting delays and are shown as a dashed line for the current season. As hospitalization data are received each week, prior case counts and rates are updated accordingly.

Additional FluSurv-NET hospitalization surveillance information for current and past seasons and additional age groups:

Surveillance Methods (<https://www.cdc.gov/flu/weekly/overview.htm#FluSurvNet>) | FluView Interactive: Rates by Age, Sex, and Race/Ethnicity (<http://gis.cdc.gov/GRASP/Fluview/FluHospRates.html>) or Data on Patient Characteristics (<http://gis.cdc.gov/grasp/fluview/FluHospChars.html>) | RESP-NET Interactive (<https://www.cdc.gov/surveillance/resp-net/dashboard.html>)

National Healthcare Safety Network (NHSN) Hospitalization Surveillance

Hospitals report to NHSN the weekly number of patients admitted with laboratory-confirmed influenza. During Week 8, 10,148 patients with laboratory-confirmed influenza were admitted to a hospital. Nationally, the number of patients admitted to a hospital with laboratory-confirmed influenza for Week 8 remained stable (change of <5%) compared to Week 7. The number of hospitalizations increased in regions 5, 7, and 10, remained stable in regions 1 and 3, and decreased in regions 2, 4, 6, 8, and 9 this week compared to Week 7.

Additional NHSN Hospitalization Surveillance information:

Surveillance Methods (<https://www.cdc.gov/flu/weekly/overview.htm#NHSN>) | Additional Data (<https://data.cdc.gov/Public-Health-Surveillance/Respiratory-Virus-Response-RVR-United-States-Hospi/9t9r-e5a3>) | FluView Interactive (<http://gis.cdc.gov/grasp/fluview/FluView12.html>)

Mortality Surveillance

(<https://www.cdc.gov/flu/weekly/overview.htm#MortalitySurveillance>)


National Center for Health Statistics (NCHS) Mortality Surveillance

Based on NCHS mortality surveillance data available on February 29, 2024, 0.7% of the deaths that occurred during the week ending February 24, 2024 (Week 8), were due to influenza. This percentage remained stable (< 0.1 percentage point change) compared to Week 7. The data presented are preliminary and may change as more data are received and processed.

Season: Surveillance Area: Age:

**Pneumonia and Influenza Mortality from
the National Center for Health Statics Mortality Surveillance System**
National Summary data through the week ending February 24, 2024

Show Number of Influenza Deaths



[View Regional and State Level Data \(https://gis.cdc.gov/grasp/fluview/Mortality.html\)](https://gis.cdc.gov/grasp/fluview/Mortality.html) | [Download Chart Data](#) | [Download PowerPoint Presentation](#)

<

Additional pneumonia, influenza and COVID-19 mortality surveillance information for current and past seasons:


[Surveillance Methods \(https://www.cdc.gov/flu/weekly/overview.htm#NCHSMortality\)](https://www.cdc.gov/flu/weekly/overview.htm#NCHSMortality) | [FluView Interactive \(https://gis.cdc.gov/grasp/fluview/mortality.html\)](#)

Influenza-Associated Pediatric Mortality

Two influenza-associated pediatric deaths occurring during the 2023-2024 season were reported to CDC during Week 8. One death occurred during Week 6 (the week ending February 10, 2024) and one occurred during Week 7 (the week ending February 17, 2024). Both deaths were associated with an influenza B virus with no lineage determined.

A total of 93 influenza-associated pediatric deaths occurring during the 2023-2024 season have been reported to CDC.

**Influenza-Associated Pediatric Deaths
by Week of Death, 2020-21 season to 2023-24 season**



View FluView Interactive (<https://gis.cdc.gov/grasp/fluview/PedFluDeath.html>) | Download Chart Data | Download PowerPoint Presentation

Additional pediatric mortality surveillance information for current and past seasons:
[Surveillance Methods \(https://www.cdc.gov/flu/weekly/overview.htm#PediatricMortality\)](https://www.cdc.gov/flu/weekly/overview.htm#PediatricMortality) | [FluView Interactive \(https://gis.cdc.gov/GRASP/Fluview/PedFluDeath.html\)](https://gis.cdc.gov/GRASP/Fluview/PedFluDeath.html)

Trend Indicators

Increasing: ↑
 Decreasing: ↓
 Stable: →

Indicators Status by System

Clinical Labs: Up or down arrows indicate a change of greater than or equal to 0.5 percentage points in the percent of specimens positive for influenza compared to the previous week.

Outpatient Respiratory Illness (ILINet): Up or down arrows indicate a change of greater than 0.1 percentage points in the percent of visits due to respiratory illness (ILI) compared to the previous week.

NHSN Hospitalizations: Up or down arrows indicate change of greater than or equal to 5% of the number of patients admitted with laboratory-confirmed influenza compared to the previous week.

NCHS Mortality: Up or down arrows indicate change of greater than 0.1 percentage points of the percent of deaths due to influenza compared to the previous week.

Reference Footnotes

¹U.S. Influenza Surveillance: Purpose and Methods (2023 Oct). Centers for Disease Control and Prevention. <https://www.cdc.gov/flu/weekly/overview.htm#ILINet> (<https://www.cdc.gov/flu/weekly/overview.htm#ILINet>).

²Grohskopf LA, Blanton LH, Ferdinands JM, Chung JR, Broder KR, Talbot HK. Prevention and Control of Seasonal Influenza with Vaccines: Recommendations of the Advisory Committee on Immunization Practices — United States, 2023–24 Influenza Season. MMWR Recomm Rep 2023;72(No. RR-2):1–25. DOI: <http://dx.doi.org/10.15585/mmwr.rr7202a1> [↗](http://dx.doi.org/10.15585/mmwr.rr7202a1)

³Influenza Antiviral Medications: Summary for Clinicians (2023 Sept). Centers for Disease Control and Prevention. <https://www.cdc.gov/flu/professionals/antivirals/summary-clinicians.htm> (<https://www.cdc.gov/flu/professionals/antivirals/summary-clinicians.htm>).

Additional National and International Influenza Surveillance Information

FluView Interactive: FluView includes enhanced web-based interactive applications that can provide dynamic visuals of the influenza data collected and analyzed by CDC. These [FluView Interactive applications](http://www.cdc.gov/flu/weekly/fluviewinteractive.htm) (<http://www.cdc.gov/flu/weekly/fluviewinteractive.htm>) allow people to create customized, visual interpretations of influenza data, as well as make comparisons across flu seasons, regions, age groups and a variety of other demographics.

National Institute for Occupational Safety and Health: Monthly surveillance data on the prevalence of health-related workplace absenteeism among full-time workers in the United States are [available from NIOSH](https://www.cdc.gov/niosh/topics/absences/default.html) (<https://www.cdc.gov/niosh/topics/absences/default.html>).

U.S. State and local influenza surveillance: Select a jurisdiction below to access the latest local influenza information.

Alabama (<http://adph.org/influenza/>)

Alaska

(<http://dhss.alaska.gov/dph/Epi/id/Pages/influenza/fluinfo>)

Colorado (<https://www.mass.gov/info-details/influenza-reporting>)

Connecticut (<https://portal.ct.gov/DPH/Epidemiology-and-Emerging-Infections/Influenza-Surveillance-and-Statistics>)

Georgia (<https://dph.georgia.gov/flu-activity-georgia>)

Hawaii
(<http://health.hawaii.gov/docd/resources/reports/influenza-reports/>)

Iowa (<https://idph.iowa.gov/influenza/reports>)

Kansas (<http://www.kdheks.gov/flu/surveillance.htm>)

Maryland (<https://phpa.health.maryland.gov/influenza/fluwatch/>)

Massachusetts (<https://www.mass.gov/info-details/influenza-reporting>)

Missouri
(<http://health.mo.gov/living/healthcondiseases/communicable/influenza/reports.php>)

Montana
(<https://dphhs.mt.gov/publichealth/cdepi/diseases/influenza>)

New Jersey (<http://www.nj.gov/health/cd/topics/flu.shtml>)

New Mexico (<https://nmhealth.org/about/erd/ideb/isp/>)

Ohio (<http://www.flu.ohio.gov>)

Oklahoma (<https://oklahoma.gov/health/health-education/acute-disease-service/disease-information/influenza-home-page.html>)

South Carolina
(<http://www.scdhec.gov/Health/DiseasesandConditions/InfectiousDiseases/Flu/FluData/>)

South Dakota (<https://doh.sd.gov/health-data-reports/dashboards/influenza-dashboard/>)

Vermont (<http://www.healthvermont.gov/immunizations-infectious-disease/influenza/flu-activity-and-surveillance>)

Virginia (<http://www.vdh.virginia.gov/epidemiology/influenza-in-virginia/influenza-surveillance/>)

Wyoming (<https://health.wyo.gov/publichealth/infectious-disease-epidemiology-unit/disease/influenza/>)

New York City (<http://www1.nyc.gov/site/doh/provider-topics/flu-alerts.page>)

World Health Organization:

Additional influenza surveillance information from participating WHO member nations is available through FluNet [🔗](https://www.who.int/tools/flunet) (<https://www.who.int/tools/flunet>) and the Global Epidemiology Reports. [🔗](https://www.who.int/teams/global-influenza-programme/surveillance-and-monitoring/influenza-surveillance-outputs) (<https://www.who.int/teams/global-influenza-programme/surveillance-and-monitoring/influenza-surveillance-outputs>)


WHO Collaborating Centers for Influenza:

Australia [🔗](http://www.influenzacentre.org/Surveillance_Samples_Received.html) (http://www.influenzacentre.org/Surveillance_Samples_Received.html), China [🔗](http://www.chinaivdc.cn/cnic/) (<http://www.chinaivdc.cn/cnic/>), Japan [🔗](http://idsc.nih.go.jp/index.html) (<http://idsc.nih.go.jp/index.html>), the United Kingdom [🔗](https://www.crick.ac.uk/research/worldwide-influenza-centre) (<https://www.crick.ac.uk/research/worldwide-influenza-centre>), and the United States (<http://www.cdc.gov/flu/>) (CDC in Atlanta, Georgia)

Europe:

The most up-to-date influenza information from Europe is available from WHO/Europe and the European Centre for Disease Prevention and Control [🔗](http://www.flunewseurope.org) (<http://www.flunewseurope.org>).

Public Health Agency of Canada:

The most up-to-date influenza information from Canada is available in [Canada's weekly FluWatch report](#)  (<http://www.phac-aspc.gc.ca/fluwatch/>).

Public Health England:

The most up-to-date influenza information from the United Kingdom is available from [Public Health England](#)  (<http://www.hpa.org.uk/Topics/InfectiousDiseases/InfectionsAZ/SeasonalInfluenza/>).

Any links provided to non-Federal organizations are provided solely as a service to our users. These links do not constitute an endorsement of these organizations or their programs by CDC or the Federal Government, and none should be inferred. CDC is not responsible for the content of the individual organization web pages found at these links.

A description of the CDC influenza surveillance system, including methodology and detailed descriptions of each data component is available on the [surveillance methods](#) (<http://www.cdc.gov/flu/weekly/overview.htm>) page.

Last Reviewed: March 1, 2024, 11:00 AM

Source: Centers for Disease Control and Prevention (<https://www.cdc.gov/>), National Center for Immunization and Respiratory Diseases (NCIRD) (<https://www.cdc.gov/ncird/index.html>)