

Supplemental Table 1. Sport-related Concussions Across Varsity Sports
for Academic Years 2013-14 to 2019-20 (November 2013 through February 2020)

| Sport | Women (n=759, 36.5%) | Men (n=1318, 63.5%) | Total (n=2077, 100%) |
|---------------------------|-------------------------|------------------------|-------------------------|
| football | 0 (0.0) | 665 (50.5) | 665 (32.0) |
| soccer | 149 (19.6) | 118 (9.0) | 267 (12.9) |
| lacrosse | 88 (11.6) | 124 (9.4) | 212 (10.2) |
| basketball | 78 (10.3) | 70 (5.3) | 148 (7.1) |
| wrestling | 0 (0.0) | 127 (9.6) | 127 (6.1) |
| gymnastics | 73 (9.6) | 7 (0.5) | 80 (3.9) |
| volleyball | 69 (9.1) | 6 (0.5) | 75 (3.6) |
| field hockey | 71 (9.4) | 0 (0.0) | 71 (3.4) |
| water polo | 30 (4.0) | 40 (3.0) | 70 (3.4) |
| sprint football | 0 (0.0) | 59 (4.5) | 59 (2.8) |
| softball | 58 (7.6) | 0 (0.0) | 58 (2.8) |
| baseball | 0 (0.0) | 38 (2.9) | 38 (1.8) |
| swimming | 26 (3.4) | 8 (0.6) | 34 (1.6) |
| track&field/cross country | 17 (2.2) | 17 (1.3) | 34 (1.6) |
| diving | 19 (2.5) | 7 (0.5) | 26 (1.3) |
| rowing | 15 (2.0) | 6 (0.5) | 21 (1.0) |
| fencing | 7 (0.9) | 10 (0.8) | 17 (0.8) |
| squash | 12 (1.6) | 4 (0.3) | 16 (0.8) |
| sailing | 12 (1.6) | 3 (0.2) | 15 (0.7) |
| cheerleading | 10 (1.3) | 3 (0.2) | 13 (0.6) |
| skiing | 7 (0.9) | 2 (0.2) | 9 (0.4) |
| tennis | 7 (0.9) | 2 (0.2) | 9 (0.4) |
| equestrian | 8 (1.1) | 0 (0.0) | 8 (0.4) |
| polo | 2 (0.3) | 1 (0.1) | 3 (0.1) |
| golf | 1 (0.1) | 1 (0.1) | 2 (0.1) |

Supplemental Table 2. Linear regression

Outcome: days in RTP progression

| Variable | Women | | | | Men | | | |
|--|---------|--------------|---------|----------------|---------|--------------|---------|----------------|
| | β | 95% CI | p-value | R ² | β | 95% CI | p-value | R ² |
| Univariate linear regression | | | | | | | | |
| Affective domain | | | | | | | | |
| No affective | ref | ref | ref | | ref | ref | ref | |
| Co-occur affective | 10.82 | -2.90-24.54 | 0.12 | 0.02 | 4.00 | -3.80-11.79 | 0.31 | 0.002 |
| Only nervous-anxious | 38.41 | 7.15-69.67 | 0.02 | | -4.71 | -20.46-11.05 | 0.56 | |
| No nervous-anxious | 3.83 | -9.49-17.15 | 0.57 | | -0.10 | -6.93-6.74 | 0.98 | |
| Symptom prevalence (18) | 0.72 | -0.58-2.02 | 0.28 | 0.002 | 0.07 | -0.58-0.71 | 0.84 | 0.008 |
| Concussion history | 14.44 | 3.74-25.15 | <0.01 | 0.01 | 6.80 | 1.30-12.31 | 0.02 | 0.007 |
| Multivariable linear regression | | | | | | | | |
| Affective domain | | | | | | | | |
| No affective | ref | ref | ref | | ref | ref | ref | |
| Co-occur affective | 9.66 | -5.99-25.32 | 0.23 | | 4.90 | -4.05-13.86 | 0.28 | |
| Only nervous-anxious | 38.46 | 7.11-69.80 | 0.02 | 0.03 | -3.71 | -19.57-12.14 | 0.65 | 0.01 |
| No nervous-anxious | 3.06 | -10.80-16.91 | 0.67 | | -0.15 | -7.40-7.10 | 0.97 | |
| Symptom prevalence (18) | 0.21 | -1.28-1.70 | 0.78 | | -0.16 | -0.92-0.60 | 0.68 | |
| Concussion history | 14.60 | 3.94-23.29 | <0.01 | | 6.90 | 3.43-18.24 | 0.02 | |

Abbreviation: CI=confidence interval