

Iowa, 2007

Primary and secondary syphilis

- CASES: Iowa reported 21 cases in 2007.
- RATES: Iowa ranked 46 among 50 states, DC, and 3 territories with 0.7 cases per 100,000 population compared to the total rate of 3.8 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Iowa, the rate among males was 1.2 per 100,000 population compared to the total male rate of 6.6 per 100,000.
- The rate among females was 0.2 per 100,000 compared to the total female rate of 1.1 per 100,000.

Rates of primary and secondary syphilis, by race/ethnicity


In Iowa, the race/ethnicity adjusted rates per 100,000 population were 0.7 among whites, 2.5 among blacks, 0.0 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.

- The rate among blacks was 3.6 times that of whites.


Congenital syphilis

- CASES: Iowa reported 1 case in 2007.
- RATES: Iowa ranked 28 among 31 states, DC, and 2 territories reporting congenital syphilis cases with a rate of 2.6 cases per 100,000 live births compared to the total rate of 10.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Iowa, 1998-2007



Total P&S Syphilis Rates by County, Iowa, 2007



Rate (per 100,000 population)

