

Illinois, 2007

Primary and secondary syphilis

- CASES: Illinois reported 464 cases in 2007.
- RATES: Illinois ranked 18 among 50 states, DC, and 3 territories with 3.6 cases per 100,000 population compared to the total rate of 3.8 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Illinois, the rate among males was 6.7 per 100,000 population compared to the total male rate of 6.6 per 100,000.
- The rate among females was 0.6 per 100,000 compared to the total female rate of 1.1 per 100,000.

Rates of primary and secondary syphilis, by race/ethnicity


In Illinois, the race/ethnicity adjusted rates per 100,000 population were 2.1 among whites, 10.8 among blacks, 4.0 among Hispanics, 1.3 among Asian/Pacific Islanders, and 3.9 among American Indians/Alaska Natives.

- The rate among blacks was 5.1 times that of whites.
- The rate among Hispanics was 1.9 times that of whites.
- The rate among American Indians/Alaska Natives was 1.9 times that of whites.

Congenital syphilis

- CASES: Illinois reported 10 cases in 2007.
- RATES: Illinois ranked 20 among 31 states, DC, and 2 territories reporting congenital syphilis cases with a rate of 5.5 cases per 100,000 live births compared to the total rate of 10.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Illinois, 1998-2007



Total P&S Syphilis Rates by County, Illinois, 2007

