

# New York, 2007

## Primary and secondary syphilis

- **CASES:** New York reported 1,068 cases in 2007.
- **RATES:** New York ranked 8 among 50 states, DC, and 3 territories with 5.5 cases per 100,000 population compared to the total rate of 3.8 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In New York, the rate among males was 11.1 per 100,000 population compared to the total male rate of 6.6 per 100,000.
- The rate among females was 0.3 per 100,000 compared to the total female rate of 1.1 per 100,000.

## Rates of primary and secondary syphilis, by race/ethnicity


In New York, the race/ethnicity adjusted rates per 100,000 population were 2.9 among whites, 13.2 among blacks, 9.5 among Hispanics, 2.1 among Asian/Pacific Islanders, and 11.2 among American Indians/Alaska Natives.

- The rate among blacks was 4.6 times that of whites.
- The rate among Hispanics was 3.3 times that of whites.
- The rate among American Indians/Alaska Natives was 3.9 times that of whites.

## Congenital syphilis

- **CASES:** New York reported 16 cases in 2007.
- **RATES:** New York ranked 17 among 31 states, DC, and 2 territories reporting congenital syphilis cases with a rate of 6.4 cases per 100,000 live births compared to the total rate of 10.5 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, New York, 1998-2007



## Total P&S Syphilis Rates by County, New York, 2007

