

Wisconsin, 2007

Primary and secondary syphilis

- **CASES:** Wisconsin reported 66 cases in 2007.
- **RATES:** Wisconsin ranked 34 among 50 states, DC, and 3 territories with 1.2 cases per 100,000 population compared to the total rate of 3.8 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Wisconsin, the rate among males was 2.2 per 100,000 population compared to the total male rate of 6.6 per 100,000.
- The rate among females was 0.2 per 100,000 compared to the total female rate of 1.1 per 100,000.

Rates of primary and secondary syphilis, by race/ethnicity


In Wisconsin, the race/ethnicity adjusted rates per 100,000 population were 0.5 among whites, 9.1 among blacks, 3.1 among Hispanics, 0.9 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.

- The rate among blacks was 18.2 times that of whites.
- The rate among Hispanics was 6.2 times that of whites.
- The rate among Asian/Pacific Islanders was 1.8 times that of whites.


Congenital syphilis

- **CASES:** Wisconsin reported 1 case in 2007.
- **RATES:** Wisconsin ranked 31 among 31 states, DC, and 2 territories reporting congenital syphilis cases with a rate of 1.4 cases per 100,000 live births compared to the total rate of 10.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Wisconsin, 1998-2007



Total P&S Syphilis Rates by County, Wisconsin, 2007



Rate (per 100,000 population)

