

SUPPLEMENTARY TABLE. Prevalence of mammography use in past 2 years among women aged 50–74 years by state and number of reported adverse social determinants of health and health-related social needs – Behavioral Risk Factor Surveillance System, United States, 2022

State	Prevalence (95% CI) by no. of reported adverse social determinants of health and health-related social needs			
	0	1	2	3–11
Alabama	89.1 (84.7–92.4)	82.8 (75.9–88.1)	75.4 (61.3–85.6)	56.7 (47.3–65.5)
Alaska	76.8 (71.8–81.1)	61.4 (49.1–72.4)	66.9 (54.1–77.6)	53.8 (43.2–64.1)
Arizona	82.0 (77.7–85.6)	77.2 (69.1–83.8)	72.9 (62.6–81.3)	54.8 (45.3–64.0)
California	85.5 (81.4–88.9)	79.3 (71.1–85.6)	64.8 (53.2–75.0)	66.5 (55.5–75.9)
Connecticut	87.1 (83.3–90.1)	87.2 (81.2–91.5)	77.4 (68.9–84.1)	70.1 (61.2–77.7)
Delaware	85.4 (80.9–88.9)	81.1 (72.8–87.4)	75.1 (63.0–84.2)	72.4 (61.6–81.1)
District of Columbia	85.4 (79.1–90.1)	77.4 (66.6–85.5)	63.6 (43.5–79.8)	83.8 (73.9–90.4)
Florida	81.8 (76.3–86.2)	83.1 (74.4–89.3)	77.0 (66.1–85.3)	70.6 (59.5–79.8)
Georgia	83.2 (78.7–87.0)	78.0 (71.6–83.2)	76.4 (66.4–84.2)	60.2 (51.0–68.7)
Idaho	72.6 (68.4–76.4)	64.7 (57.4–71.4)	65.2 (54.4–74.7)	54.7 (44.6–64.4)
Indiana	83.1 (80.4–85.5)	75.0 (69.5–79.8)	78.9 (71.4–84.9)	69.4 (62.8–75.3)
Iowa	83.4 (79.4–86.7)	84.0 (79.0–88.0)	69.9 (58.5–79.2)	74.5 (65.0–82.1)
Kansas	81.1 (77.9–83.9)	73.0 (66.1–79.0)	67.1 (58.4–74.8)	52.4 (44.2–60.5)
Kentucky	82.2 (75.9–87.1)	76.6 (65.6–84.9)	71.4 (55.8–83.1)	61.5 (50.8–71.3)
Maine	84.3 (81.2–87.0)	89.5 (85.7–92.4)	75.3 (67.4–81.8)	74.1 (66.2–80.7)
Maryland	85.7 (81.9–88.9)	84.0 (78.4–88.4)	83.5 (75.0–89.5)	77.7 (70.6–83.4)
Massachusetts	88.2 (84.5–91.1)	87.2 (82.8–90.6)	83.1 (75.4–88.7)	80.9 (74.1–86.2)
Michigan	83.0 (79.3–86.2)	77.0 (70.7–82.4)	65.7 (53.0–76.5)	65.1 (56.9–72.6)
Minnesota	85.4 (82.9–87.6)	76.0 (70.6–80.7)	77.5 (68.6–84.4)	69.3 (62.7–75.3)
Mississippi	83.6 (77.2–88.5)	71.3 (59.4–80.8)	70.3 (57.7–80.4)	60.2 (50.1–69.4)
Missouri	80.3 (76.0–84.0)	76.0 (68.8–82.0)	79.3 (68.0–87.4)	60.1 (51.5–68.2)
Montana	78.0 (74.0–81.5)	75.3 (67.4–81.8)	69.2 (58.8–77.9)	59.1 (48.7–68.8)
Nebraska	82.4 (77.4–86.5)	77.4 (67.2–85.1)	78.9 (63.4–89.0)	68.8 (55.2–79.8)
Nevada	74.1 (64.5–81.8)	72.5 (59.2–82.7)	66.4 (48.1–80.8)	59.4 (45.7–71.7)
New Hampshire	85.8 (82.6–88.4)	81.3 (75.5–86.0)	68.9 (57.6–78.3)	69.6 (59.7–78.0)
New Jersey	84.6 (80.5–87.9)	74.2 (65.1–81.5)	68.6 (58.7–77.1)	66.2 (56.4–74.8)
New Mexico	72.9 (66.5–78.4)	67.7 (57.8–76.2)	65.6 (53.4–76.0)	56.3 (47.0–65.2)
North Carolina	86.7 (81.9–90.3)	84.5 (77.2–89.8)	82.7 (69.9–90.8)	71.5 (58.8–81.5)
Ohio	83.2 (79.4–86.4)	74.8 (68.4–80.3)	72.3 (62.1–80.6)	58.6 (50.2–66.6)
Oklahoma*	81.2 (74.4–86.5)	63.3 (51.4–73.8)	—	52.7 (39.6–65.3)
Rhode Island	91.0 (87.7–93.5)	84.8 (78.4–89.6)	79.8 (68.5–87.7)	78.0 (69.8–84.4)
South Carolina	85.4 (82.5–87.8)	80.0 (74.1–84.8)	75.8 (67.5–82.4)	67.3 (60.2–73.7)
Tennessee	81.0 (75.3–85.6)	67.6 (56.9–76.7)	78.6 (65.8–87.6)	66.2 (55.5–75.4)
Texas	82.1 (77.8–85.8)	76.3 (68.7–82.6)	74.7 (64.7–82.7)	62.0 (52.7–70.5)
Utah	80.6 (77.1–83.7)	68.1 (61.2–74.2)	74.5 (63.9–82.8)	61.8 (51.5–71.2)
Vermont	78.6 (74.9–82.0)	77.8 (72.0–82.7)	71.4 (61.5–79.6)	60.4 (51.4–68.8)
Washington	78.9 (76.5–81.0)	76.0 (72.4–79.3)	73.3 (68.1–77.9)	63.0 (57.9–67.8)
West Virginia	84.4 (80.3–87.7)	80.0 (72.8–85.7)	68.8 (58.9–77.2)	70.4 (62.3–77.4)
Wisconsin	86.0 (83.2–88.5)	85.7 (81.4–89.1)	77.6 (70.2–83.5)	74.7 (68.1–80.4)
Wyoming	69.6 (64.6–74.1)	57.9 (48.3–67.0)	70.2 (57.7–80.3)	44.8 (32.5–57.7)

*Prevalence for 2 adverse social determinants of health and health-related social needs suppressed because sample size <50.