


FoodNet Surveillance Report for 2000

(Preliminary Report)



Centers for Disease Control and Prevention
National Center for Infectious Diseases
Emerging Infections Program
Division of Bacterial and Mycotic Diseases
Foodborne and Diarrheal Diseases Branch
Foodborne Diseases Active Surveillance Network
September 2001

Table of Contents

Executive summary	3
Background	5
Objectives	5
Methods	5
Results	5
Cases reported	5
Seasonality	6
2000 Rates	7
Rates by site	8
Rates by age	9
Rates by sex	9
Rates by age and sex	10
Hospitalizations	10
Outbreaks	11
1996-2000 Rates	12
Comments.	18
Other ongoing projects	20
Burden of illness.	20
Routes of transmission of foodborne pathogens	20
Other FoodNet activities	21
Future activities	22
On-line Materials Available on Web Site.	23
2000 FoodNet Working Group	24
FoodNet Web-Site	25

Executive summary

The Foodborne Diseases Active Surveillance Network (FoodNet) is the principal foodborne-disease component of the Centers for Disease Control and Prevention's (CDC's) Emerging Infections Program (EIP). FoodNet is a collaborative project among CDC, the nine EIP sites, the Food Safety and Inspection Service (FSIS) of the United States Department of Agriculture (USDA), and the United States Food and Drug Administration (FDA). FoodNet augments, but does not replace, longstanding activities at CDC, FSIS, FDA, and in states to identify, control, and prevent foodborne disease hazards.

FoodNet is a sentinel network that is producing more stable and accurate national estimates of the burden and sources of specific foodborne diseases in the United States through active surveillance and additional studies. Enhanced surveillance and investigation are integral parts of developing and evaluating new prevention and control strategies that can improve the safety of our food and the public's health. Ongoing FoodNet surveillance is being used to document the effectiveness of new food safety control measures, such as the USDA, FSIS Pathogen Reduction and Hazard Analysis and Critical Control Point (HACCP) Systems, in decreasing the number of cases of foodborne diseases in the United States each year.

The following are key findings of FoodNet surveillance activities during 2000:

- A modest decline in incidence of *Campylobacter* and *Salmonella* infection indicates that further prevention efforts are needed to meet the Healthy People 2010 objectives for those pathogens.
- *Salmonella* serotype Enteritidis infections continued to decline. Since the major source of *Salmonella* Enteritidis infections is eggs, this decline may be related to improvements in hygiene on egg-laying hen farms, improvements in keeping eggs refrigerated during transport and distribution, increased use of pasteurized eggs and egg products, and better cooking and handling of eggs in the kitchen.
- There was no substantial change in the rate of *E. coli* O157 infections. However, the absence of a large, recognized outbreak highlights the importance of sporadic infections. Preventing *E. coli* O157 will not be a simple task because it can be transmitted through food, water, person-to-person contact, and direct animal exposure. FoodNet studies and recent outbreaks have shown that an important route of transmission is from direct contact with cattle or their environment. Strategies that reduce *E. coli* O157 on farms could decrease food contamination and direct contact infection, as well as entry into the water supply.
- The incidence of *Listeria* infections has decreased over the past twelve years. A previous surveillance system reported an annual *Listeria* infection rate of 1.6 per 100,000 persons in 1989 compared with the rate of 0.34 per 100,000 persons in FoodNet sites in 2000 (Tappero J, Schuchat A, Deavers K, et al. Reduction in the incidence of human listeriosis in the United States. JAMA 1995;273:1118-1122.) This decline over the past 12 years suggests that improvements by the food industry in sanitation have been effective. PulseNet, CDC's network of public health laboratories that subtype bacteria that can be transmitted by food, has led the way to improved outbreak identification in recent years (Information available at

<http://www.cdc.gov/ncidod/dbmd/pulsenet/pulsenet.htm>). Investigation of these outbreaks, such as one due to deli turkey meat in 2000, indicates that production processes still need improvement to reach the Healthy People 2010 objective of 0.25 per 100,000 persons (CDC. Multistate outbreak of listeriosis - United States, 2000. MMWR 2000;49:1129-1130. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm4950a1.htm>).

- There are important regional variations in the rates of specific bacterial foodborne infections. For example, *Campylobacter* infections are five times more common in California than in Georgia, Tennessee, or Maryland sites. *E. coli* O157 infections are more common in the northern states of Connecticut, Minnesota, New York, and Oregon than in other sites. *Salmonella* shows less variation, being relatively common in all the FoodNet sites. Focused research into the reasons for these local differences may provide information about prevention that is of general use.
- The FoodNet surveillance system will be useful in measuring progress towards the US Department of Health and Human Services' Healthy People 2010 Objectives; these objectives are designed to measure and address preventable health threats to the nation (US Department of Health and Human Services. Healthy People 2010: Understanding and Improving Health. 2nd ed. Washington, D.C: US Government Printing Office, November 2000. Available at <http://www.health.gov/healthypeople/Document/Word/uih/uih.doc>). The year 2000 rates in FoodNet sites meet the Healthy People 2000 Objectives for each of the four infections, *Campylobacter jejuni*, *Salmonella*, *E.coli* O157:H7, and *Listeria monocytogenes*, specifically targeted by the Department of Health and Human Services. However, further prevention efforts are needed to reach the 2010 objectives.

Background

Foodborne infections are an important public health challenge. The Centers for Disease Control and Prevention (CDC) estimates that in 1997, foodborne infections caused 76 million illnesses, 325,000 hospitalizations, and 5,000 deaths. CDC, the Food Safety and Inspection Service (FSIS) of the United States Department of Agriculture (USDA), the United States Food and Drug Administration (FDA), and the nine Emerging Infections Program (EIP) sites are actively involved in preventing foodborne diseases. In 1997, the interagency national Food Safety Initiative was established to meet the public health challenge of foodborne diseases. CDC's principal role in the Food Safety Initiative has been to enhance surveillance and investigation of infections that are usually foodborne. FoodNet has been instrumental in accomplishing this mission.

Objectives

The objectives of FoodNet are to determine the frequency and severity of foodborne diseases; determine the association of common foodborne diseases with eating specific foods; and describe the epidemiology of new and emerging bacterial, parasitic, and viral foodborne pathogens. To address these objectives, FoodNet uses active surveillance and conducts related epidemiologic studies. By monitoring the burden of foodborne diseases over time, FoodNet can document the effectiveness of new food safety initiatives, such as the USDA Hazard Analysis and Critical Control Points (HACCP) System, in decreasing the rate of foodborne diseases in the United States each year.

Methods

In 2000, FoodNet conducted population-based active surveillance for laboratory-confirmed cases of *Campylobacter*, *Cryptosporidium*, *Cyclospora*, Shiga toxin-producing *E. coli*, including *E. coli* O157, *Listeria*, *Salmonella*, *Shigella*, *Vibrio*, and *Yersinia* infections in Connecticut, Georgia, Minnesota, and Oregon and selected counties in California, Maryland, New York, and Tennessee (total population 29.5 million)¹. To identify cases, FoodNet personnel contact each of the more than 450 clinical laboratories serving the catchment areas, either weekly or monthly, depending on the size of the clinical laboratory. FoodNet also conducts surveillance for foodborne disease outbreaks and hemolytic uremic syndrome, the latter principally through pediatric nephrologists.

Results

Cases reported

The results contained in this report are preliminary. FoodNet attempts to finalize surveillance case-counts in late February; however, audits and additional case-finding may change the number of cases that will be published in the final 2000 Annual Report. In 2000, a total of 12,631 laboratory confirmed infections caused by the pathogens under surveillance were identified in eight sites. Of these, 12,125 were bacterial, including 4,640 *Campylobacter* infections, 4,237 *Salmonella* infections, 2,324 *Shigella* infections, 631 *E. coli* O157 infections², 131 *Yersinia* infections, 101 *Listeria*

1 A ninth site, selected counties in Colorado, was added in 2000 and will contribute active surveillance data in 2001.

2 This report only contains information on *E. coli* O157 infections. Information on other STEC infections will be included in the 2000 Final

infections, and 61 *Vibrio* infections (Table 1A). Of the 3,663 *Salmonella* isolates that were serotyped, the most commonly identified serotypes were Typhimurium (862 cases), Enteritidis (565), Newport (399), and Heidelberg (248). In addition, 506 cases of parasitic diseases were reported, including 484 cases of *Cryptosporidium* infection and 22 cases of *Cyclospora* infection (Table 1B).

Table 1A. Infections caused by specific bacterial pathogens, reported by FoodNet sites, 2000

Pathogen	Total
<i>Campylobacter</i>	4640
<i>Escherichia coli</i> O157	631
<i>Listeria</i>	101
<i>Salmonella</i>	4237
<i>Shigella</i>	2324
<i>Vibrio</i>	61
<i>Yersinia</i>	131
Total	12125

Table 1B. Infections caused by specific parasitic pathogens, reported by FoodNet sites, 2000



Pathogen	Total
<i>Cryptosporidium</i>	484
<i>Cyclospora</i>	22
Total	506

Seasonality

Isolation rates for pathogens showed seasonal variation; 45% of *E. coli* O157, 38% of *Salmonella*, 37% of *Campylobacter*, and 37% of *Shigella* were

isolated during June through August (Figure 1). *Yersinia* infections were more likely to have occurred in winter months, with 46% of cases being reported during January, February, or December (Figure 1).

Figure 1. Cases of foodborne disease caused by specific pathogens, by month, FoodNet sites, 2000



2000 Rates


To compare the number of cases among sites with different populations, preliminary annual incidence rates were calculated. Incidence is the number of cases divided by the population. All 2000 rates reported here use preliminary numbers of cases for 2000 and 1999 population estimates and are therefore considered preliminary. Final incidence data will be available once


2000 population counts are available in mid-2001. Overall incidence rates were highest for infections with *Campylobacter* (15.7/100,000 population), *Salmonella* (14.4/100,000), and *Shigella* (7.9/100,000). Lower overall incidence rates were reported for *E. coli* O157 (2.1/100,000), *Cryptosporidium* (1.5/100,000), *Yersinia* (0.44/100,000), *Listeria* (0.34/100,000), *Vibrio* (0.21/100,000), and *Cyclospora* (0.067/100,000).

Rates by site

Incidence rates for many of these pathogens varied substantially among the sites (Figure 2). The incidence rates for *Campylobacter* infection varied from 6.6/100,000 in Tennessee to 38.2/100,000 in California, and for *Shigella* infections from 1.1/100,000 in New York to 18.8/100,000 in Minnesota. Incidence rates for aggregate *Salmonella* infection also varied among the sites, from 8.9/100,000 in Oregon to 18.0/100,000 in Georgia. Among the two most common serotypes of *Salmonella*, *S. Typhimurium* ranged from 1.9/100,000 in California to 3.7/100,000 in Tennessee and *S. Enteritidis* ranged from 1.0/100,000 in New York and Tennessee to 5.1/100,000 in Maryland. Incidence rates for *E. coli* O157 infection varied from 0.53/100,000 in Maryland to 4.6/100,000 in Minnesota. Infection caused by *Yersinia* varied from 0.21/100,000 in Minnesota to 0.90/100,000 in California. Incidence rates of *Cryptosporidium* infection ranged from 0.24/100,000 in Maryland to 3.9/100,000 in Minnesota. Listeriosis ranged from 0.15/100,000 in Minnesota to 1.0/100,000 in New York, and *Vibrio* infections ranged from no detected cases in New York to 0.90/100,000 in California. Reasons for these regional differences in incidence rates are being investigated; for example, most laboratories do not test specimens routinely for all pathogens. However, regional differences in *E. coli* O157 incidence are only partially accounted for by differences in laboratory practices.

Figure 2. Cases per 100,000 population of foodborne disease caused by specific pathogens, FoodNet sites, 2000






Rates by age

Annual incidence rates of foodborne illness varied by age, especially for *Campylobacter* and *Salmonella* infections (Figure 3). For children <1 year of age, the rate of *Salmonella* infection was 88.4/100,000 and the rate of *Campylobacter* infection was 33.6/100,000, rates substantially higher than for other age groups.

Figure 3. Incidence of *Campylobacter* and *Salmonella* infections by age group, FoodNet sites, 2000



Rates by sex

Incidence rates also varied by sex (Table 2). Overall, males were more likely than females to be infected with every pathogen except *Cyclospora*, *E. coli* O157 and *Listeria*. Rates of *Cryptosporidium* infection were 67% higher among males, rates of *Campylobacter* infection were 28% higher among males, and rates of *Shigella* were 20% higher among males.

Table 2. Sex-specific incidence rates per 100,000 population, by pathogen, FoodNet sites, 2000

Pathogen	Male	Female
<i>Campylobacter</i>	17.6	13.8
<i>Cryptosporidium</i>	1.84	1.10
<i>Cyclospora</i>	0.04	0.09
<i>E. coli</i> O157	1.95	2.31
<i>Listeria</i>	0.30	0.38
<i>Salmonella</i>	14.5	14.0
<i>Shigella</i>	8.50	7.07
<i>Vibrio</i>	0.31	0.10
<i>Yersinia</i>	0.49	0.40


Rates by age and sex

The incidence rate of *Campylobacter* infection was higher for males than for females in all age groups, except for persons aged 20-29 years. Among persons more than 20 years of age, the incidence rate of *Salmonella* infection was higher among women than among men.

Hospitalizations

Information on hospitalization is still being collected. Preliminary data show that overall, 14.8% of persons with culture-confirmed infection were hospitalized; hospitalization rates differed markedly by pathogen (Figure 4). The percentage of hospitalizations was highest for persons infected with *Listeria* (90% of reported cases) followed by those infected with *E. coli* O157 (40%), *Yersinia* (27%), *Vibrio* (21%), *Cryptosporidium* (18%), *Salmonella* (17%), *Shigella* (10%), and *Campylobacter* (10%).

Figure 4. Percentage of persons hospitalized with infections caused by specific pathogens, FoodNet sites, 2000



Foodborne Outbreaks An outbreak is defined as two or more cases of a similar illness that are epidemiologically linked. The overall rate of reported foodborne outbreaks in FoodNet sites in which 10 or more persons become ill was 3.4 outbreaks per million population, ranging from 1.5 outbreaks per million in Connecticut to 5.9 outbreaks per million in Minnesota (Table 3A). The variation in rates may be in part explained by variation in resources and disease-surveillance activities of state and local public health agencies. Almost half of reported foodborne outbreaks were viral in etiology and nearly a quarter had unknown etiologies (Table 3B).

Table 3A: Outbreaks with 10 or more persons ill by FoodNet site, 2000

	Number of Outbreaks	Outbreak Rate / 1,000,000 persons	Median # Ill
California	10	3.2	21
Connecticut	5	1.5	13
Georgia	25	3.2	42
Maryland	8	3.3	21
Minnesota	28	5.9	21
New York	8	3.8	22
Oregon	11	3.3	37
Tennessee	6	2.2	73
Total	101	3.4	24

Table 3B: Reported outbreaks with 10 or more persons ill, by pathogen,

FoodNet Sites, 2000


Etiology	Number of Outbreaks	Median # Ill
<i>Bacillus cereus</i>	1	12
<i>Campylobacter</i>	1	13
<i>Clostridium</i>	4	49
<i>Cyclospora</i>	1	29
<i>E. coli</i> O157	4	40
Hepatitis A Virus	1	39
Norwalk-Like Virus	38	30
<i>Salmonella</i>	7	16
Scombroid Toxin	1	11
<i>Shigella</i>	2	42
<i>Staphylococcus</i>	8	15
Unknown, viral profile*	11	22
Unknown, other	22	22
Total	101	24

*Unconfirmed viral etiology largely based on symptoms

1996-2000 Rates

Because the population under surveillance expanded substantially from 1996 to 2000 (Figures 5, 6), examining the data only from the original five sites provides consistency (Table 4A). Comparing years 1996 to 2000, the incidence of laboratory-diagnosed campylobacteriosis declined in the original five sites combined, and in four of the five original sites considered individually (Figure 7A). The magnitude and pattern of decrease varied by site. The incidence of diagnosed salmonellosis declined in all five sites combined and in each of the five original sites (Figure 7B). Comparing 1996 to 2000, the incidence of infection with each of the two most common serotypes of *Salmonella* also declined, from 2.5 to 1.8 for *Salmonella* Enteritidis and from 3.9 to 2.7 for *Salmonella* Typhimurium (Table 4B). The incidence of listeriosis declined overall and in each of the sites (Figure 7C). In contrast, the overall incidence of shigellosis varied substantially from year to year and from site to site; the incidence increased in all sites combined and in four of the five individual sites; large increases occurred in California and Minnesota during 2000 (Figure 7D). The overall incidence of diagnosed *E. coli* O157 infections increased in the combined five sites and in four of the five original sites considered separately. There was marked year-to-year fluctuation in the rates of *E. coli* O157 infections in individual sites, and marked variation from site to site (Figure 7E). *Vibrio* rates were consistently higher in California than in the other four FoodNet sites (Figure 7F). Although *Yersinia* rates in FoodNet decreased from 1996-2000, Georgia still reported higher rates than the other four FoodNet sites (Figure 7G). The incidence of cryptosporidiosis and cyclosporiasis also declined after surveillance began in 1997.

Figure 5. 1996 FoodNet surveillance area (Sites indicated by black areas) (total population=14, 281, 096)



**Figure 6. 2000 FoodNet surveillance area (Sites indicated by black areas)
(total population=32, 919, 948)**




Table 4A. Rate* of selected pathogens detected by FoodNet at the five original sites and the 2000 sites, by year and pathogen, 1996-2000

Pathogen	Original Five Sites					2000 Sites
	1996	1997	1998	1999 [§]	2000 [§]	2000
<i>Campylobacter</i>	23.5	25.3	21.4	17.5	20.1	15.7
<i>Cryptosporidium</i>	NR ^α	3.7**	2.9**	1.8**	2.4**	1.5
<i>Cyclospora</i>	NR ^α	0.40**	0.06**	0.05**	0.07**	0.07
<i>E. coli</i> O157	2.7	2.3	2.8	2.1	2.9	2.1
<i>Listeria</i>	0.46	0.49	0.57	0.53	0.37	0.34
<i>Salmonella</i>	14.5	13.6	12.3	13.6	12.0	14.4
<i>Shigella</i>	8.9	7.5	8.5	5.0	11.6	7.9
<i>Vibrio</i>	0.15	0.33	0.25	0.20	0.28	0.21
<i>Yersinia</i>	1.04	0.92	1.01	0.83	0.55	0.44

* Per 100,000 population

[†]In 1996, active surveillance began for laboratory-confirmed cases of *Campylobacter*, *Escherichia coli* O157, *Listeria*, *Salmonella*, *Shigella*, *Vibrio*, and *Yersinia* infections in Minnesota, Oregon, and selected counties in California, Connecticut, and Georgia. In 1997, active surveillance began for laboratory-confirmed cases of *Cryptosporidium* and *Cyclospora* infections in Connecticut, Minnesota, and Oregon, and selected counties in California.

[§] Urine isolates excluded, because urine isolates were not reported before 1999.

^α Not Reported

**Rates from 1997-2000 for *Cyclospora* and *Cryptosporidium* were calculated using the 1997 catchment area. Connecticut, Minnesota and selected counties in California began data collection at the beginning of 1997; Oregon and other selected counties in California began this process mid-year. Only full-year data (CA, CT, MN) are included in these rates.

Table 4B. Rate* of *S. Enteritidis* and *S. Typhimurium* detected by


FoodNet at the five original sites, by year and serotype, 1996-2000

Pathogen	1996	1997	1998	1999	2000
<i>Salmonella</i> Enteritidis	2.5	2.3	1.4	1.3	1.8
<i>Salmonella</i> Typhimurium	3.9	3.9	3.7	3.5	2.7


* Per 100,000 population.

Figure 7. Incidence of diagnosed infections for pathogens under surveillance in FoodNet at the five original sites, by year and organism – United States, 1996-2000


7A) *Campylobacter* infections




7B) *Salmonella* infections




7C) *Listeria* Infections




7D) *Shigella* Infections




7E) *E. coli* O157 infections



7F) *Vibrio* Infections



7G) *Yersinia* Infections



Comments

In all years, *Campylobacter* was the most frequently diagnosed pathogen, followed by *Salmonella*, *Shigella*, and *E. coli* O157; however, there was substantial regional and year-to-year variation. Differences in calendar year 2000 rates between the expanded and original (Table 4A) populations reflect regional differences in pathogen isolation rates. However, the rank order of isolation of pathogens is the same. Despite the year-to-year variation and regional fluctuations, the general magnitude of incidence and the relative order of pathogens has remained the same. This indicates that this expanded system will be useful for measuring progress towards the Healthy People 2010 Objectives for infections with *Campylobacter* (for which the 2010 objective is 12.3 per 100,000), *E. coli* O157 (2010 objective is 1.0 per 100,000), *Salmonella* (2010 objective is 6.8 per 100,000) and *Listeria* (2010 objective is 0.25 per 100,000) (Table 5) and for assessing attainment of the Healthy People 2000 Objectives. FoodNet data indicate a need to increase prevention efforts to reach the 2010 objectives.

Table 5. Comparing 2000 rates with the Healthy People 2000 and 2010 Objectives

Pathogen	2000 Rate*	2000 Objective*	2010 Objective*
<i>Campylobacter</i>	15.7	25.0	12.3
<i>Escherichia coli</i> O157	2.14	4.0	1.0
<i>Salmonella</i>	14.4	16.0	6.8
<i>Listeria</i>	0.34	0.50	0.25

*Per 100,000 population

The incidence of listeriosis (0.3 per 100,000 population in all sites combined) in 2000 was lower than in earlier reports from a previous surveillance system (1.6 in 1989) (Tappero J, Schuchat A, Deavers K, et al. Reduction in the incidence of human listeriosis in the United States. JAMA 1995;273:1118-1122.). The decline in *Listeria* infection over the past 12 years suggests that improvements by the food industry in sanitation have been effective. The 2000 decline may indicate further improvements resulting from PulseNet's contribution to outbreak identification. As PulseNet identifies outbreaks, particularly multistate outbreaks at a greater rate, public health officials are more quickly able to identify, respond, and contain outbreaks (<http://www.cdc.gov/ncidod/dbmd/pulsenet/pulsenet.htm>). However, the recent national *Listeria* outbreak due to deli turkey meat indicates the need for further improvements in the production process (CDC. Multistate outbreak of listeriosis - United States, 2000. MMWR 2000;49:1129-1130. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm4950a1.htm>).

The incidence of laboratory-diagnosed *Salmonella* and *Campylobacter* infections declined from 1996 to 2000. The year-to-year variations make overall trends difficult to measure with precision. However, *Salmonella* Enteritidis and *Salmonella* Typhimurium infections have modestly, but steadily, declined (Table 4B). The major source of *Salmonella* Enteritidis

infections is eggs; the continued decline in *Salmonella* Enteritidis infections may be related to improvements in hygiene on egg-laying farms, improvements in keeping eggs refrigerated during transport and distribution, increased use of pasteurized eggs and egg products, and better cooking and handling of eggs in the kitchen.

Although the incidence increased from 1999 to 2000 in the original five sites, a longer term trend in the incidence of diagnosed *E. coli* O157 cannot be discerned. The continued problem of *E. coli* O157 indicates that despite the absence of large, recognized outbreaks, this remains an important pathogen. Preventing *E. coli* O157 will not be a simple task because it can be transmitted through food, water, person-to-person contact, and direct animal exposure. FoodNet studies and recent outbreaks have shown that a growing part of the problem results from direct animal exposure, i.e., children visiting petting farms. Control of transmission through the food chain will not prevent these infections; strategies that reduce *E. coli* O157 on farms will alleviate both food contamination and direct contact infection.

The substantial overall increase in shigellosis (mainly caused by *Shigella sonnei*) was driven primarily by large increases in Minnesota and California. An estimated 80% of shigellosis is transmitted by non-foodborne routes (Mead PS, Slutsker L, Dietz V, et al. Food-related illness and death in the United States. *Emerging Infectious Disease* 1999;5:607-25. Available at <http://www.cdc.gov/ncidod/eid/vol5no5/mead.htm>). The increase in Minnesota appears to be the result of several community outbreaks, especially in day-care centers and elementary schools, and California's increase was primarily caused by two outbreaks, a multistate foodborne outbreak associated with a commercially prepared bean dip (CDC. Public health dispatch-outbreak of *Shigella sonnei* infections associated with eating a nationally distributed dip – California, Oregon, and Washington, January 2000. *MMWR* 2000;49:60-1. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm4903a4.htm>), and a non-foodborne outbreak in San Francisco among men who have sex with men (CDC. Surveillance for Foodborne Disease Outbreaks –United States, 1993-1997. *MMWR CDC Surveillance Summaries* 2000;49(SS-1):1-51. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/ss4901a1.htm>).

Overall, the marked regional variability in the incidence of some laboratory-confirmed infections may indicate the need for prevention measures targeted to high-incidence locations. A low incidence in one area may provide information on how to decrease the incidence in other areas.

The findings in this report are subject to several limitations. First, although FoodNet surveillance encompassed just over 10% of the U.S. population in 2000, these data are subject to substantial local variation and may not be a nationally representative sample, particularly in analyses restricted to the five original sites. Second, FoodNet data are limited to laboratory-confirmed

illnesses, and most foodborne illnesses are neither laboratory confirmed nor reported to state health departments. CDC estimates that for every *Salmonella* infection confirmed, a number more occur that are not diagnosed or reported. Although clinical laboratories in FoodNet sites routinely test stool specimens for *Salmonella* and *Shigella*, and almost always for *Campylobacter*, only about 50% routinely test for *E. coli* O157, and fewer test routinely for other pathogens; variations in testing for pathogens could account for some of the variations in incidence. Third, some laboratory-confirmed illnesses reported to FoodNet can be acquired through non-foodborne routes, e.g., through contaminated water, person-to-person contact, and direct animal exposure; therefore, the reported rates do not represent foodborne sources exclusively. Further surveillance and comparison on the expanded geographic base is necessary to determine which changes represent year-to-year variation and which are definitive trends.

Other Ongoing Projects

Burden of Illness

Cases reported through active surveillance represent only a fraction of the number of cases in the community. To estimate better the number of cases of foodborne disease in the community, FoodNet conducts surveys of laboratories, physicians, and the general population in the participating EIP sites. Using these data, we can determine the proportion of people in the general population with a diarrheal illness and from among those, the number who seek medical care for the illness. We can estimate the proportion of physicians who ordered a bacterial stool culture for patients with diarrhea, and we can evaluate how variations in laboratory testing for bacterial pathogens influence the number of culture-confirmed cases. Using FoodNet and other data, CDC estimates that 76 million foodborne illnesses, 325,000 hospitalizations, and 5,000 deaths occurred in 1997 in the United States (Mead PS, Slutsker L, Dietz V, et al. Food-related illness and death in the United States. *Emerging Infectious Disease* 1999;5:607-25. Available at <http://www.cdc.gov/ncidod/eid/vol5no5/mead.htm>).

This model can be used for developing estimates of the burden of illness caused by each foodborne pathogen. For example, data from this model suggest that in 1997 there were 1,400,000 *Salmonella* infections, resulting in 113,000 physician office visits and 37,200 culture-confirmed cases in this country. Laboratory-confirmed cases alone resulted in an estimated 8500 hospitalizations and 300 deaths; additional hospitalizations and deaths occur among persons whose illness is not laboratory-confirmed.

Routes of Transmission Of Foodborne Pathogens

FoodNet conducts case-control studies to determine the proportion of foodborne diseases that are caused by specific foods or food preparation and handling practices. To date, FoodNet has conducted case-control studies of *E. coli* O157, of *Salmonella* serotypes Enteritidis, Heidelberg, and Typhimurium DT104, of infant salmonellosis, and of *Campylobacter*. A *Listeria* case-control study and a *Cryptosporidium* case-control study are ongoing. By determining this proportion of foodborne diseases caused by

specific foods or food preparation and handling practices, prevention efforts can be made more specific and their effectiveness documented.

Other FoodNet Activities

- A complete analysis of the third survey of clinical laboratories in FoodNet sites conducted in 2000 to determine changes in laboratory practices is ongoing.
- The population under active surveillance was expanded by including additional counties in Tennessee in 2000. Five counties in Colorado and Prince George's County and Montgomery County in Maryland will be added in 2001.
- The third cycle of the FoodNet population survey began in February 2000 in the 8 FoodNet sites and will run for 12 months. The purpose of the survey is to estimate more precisely the burden of acute diarrheal illness in the United States. FoodNet population survey data help determine the prevalence and severity of self-reported diarrheal illness, common symptoms associated with diarrhea, the proportion of persons with diarrhea who seek care, and exposures that may be associated with foodborne illness.
- A physician survey was conducted to assess food safety education practices.
- Enrollment was continued for the *E. coli* O157 case-control study.
- Enrollment was continued for the *Cryptosporidium* case-control study.
- FoodNet collaborated with environmental health specialists to form the Environmental Health Specialists Network (EHS-Net) to strengthen relationships between epidemiology, laboratory and food protection programs and to better identify factors contributing to foodborne illness and foodborne disease outbreaks, particularly in retail establishments.

Future activities

- Continue population-based surveillance for *Campylobacter*, *Cryptosporidium*, *Cyclospora*, *Salmonella*, *Shigella*, Shiga toxin-producing *Escherichia coli* including *E. coli* O157, *Listeria*, *Yersinia*, and *Vibrio* infections and for hemolytic uremic syndrome.
- Conduct the fourth cycle of the FoodNet population survey. Scheduled to begin in late 2001 in the 9 FoodNet sites, it will run for 12 months.
- Conduct surveillance for foodborne disease outbreaks of any cause that occur within the FoodNet sites and pilot electronic reporting of outbreaks.
- Expand the population under active surveillance by including additional counties in Colorado for participation starting 2001.
- Conduct analysis of the *E. coli* O157 case-control study.
- Continue the *Cryptosporidium* case-control study.
- Continue the *Listeria* case-control study.
- Continue the physician survey on food safety education practices.
- Conduct retrospective study of infant illness within FoodNet sites.
- Continue collaboration with EHS-Net to better identify factors contributing to foodborne illness and foodborne disease outbreaks, particularly in retail establishments.
- Conduct pilot surveillance of reactive arthritis and pilot case-control studies to estimate the proportion of enteric infections that progress to reactive arthritis.

On-Line Materials Available on Web Site

The following reports are available at the FoodNet web site:

<http://www.cdc.gov/foodnet>

CDC. 1996 Final FoodNet Surveillance Report. Atlanta: Centers for Disease Control and Prevention; 1998.

CDC. 1997 Final FoodNet Surveillance Report. Atlanta: Centers for Disease Control and Prevention; 1998.

CDC. FoodNet Surveillance Report for 1998: Final Report. Atlanta: Centers for Disease Control and Prevention; 1998.

CDC. FoodNet Surveillance Report for 1999: Final Report. Atlanta: Centers for Disease Control and Prevention; 2000.

The following MMWR articles about FoodNet are available at this web site:

<http://www.cdc.gov/epo/mmwr/mmwr.html>

CDC. The Foodborne Diseases Active Surveillance Network, 1996. Morbidity and Mortality Weekly Report. 1997; 46:258-61.

CDC. Incidence of foodborne illness-FoodNet, 1997. Morbidity and Mortality Weekly Report. 1998; 47:782 -86.

CDC. Incidence of foodborne illness: Preliminary data from the Foodborne Diseases Active Surveillance Network (FoodNet) – United States, 1998. Morbidity and Mortality Weekly Report. 1999; 48:189-94.

CDC. Preliminary FoodNet data on the incidence of foodborne illnesses – selected sites, United States, 1999. Morbidity and Mortality Weekly Report. 2000; 49: 201-5.

CDC. Preliminary FoodNet data on the incidence of foodborne illnesses – selected sites, United States, 2000. Morbidity and Mortality Weekly Report. 2001; 50: 241-46.

The following FoodNet News newsletters are available at the FoodNet web site:

<http://www.cdc.gov/foodnet>

FoodNet News. Volume 1, No. 1, Fall 1998

FoodNet News. Volume 1, No. 3, Fall 1999

FoodNet News. Volume 1, No. 2, Winter 1999

FoodNet News. Volume 3, No. 1, Spring 2000

FoodNet News. Volume 3, No. 2, Winter 2000

Additional information about the pathogens under FoodNet surveillance is available at the following web sites:

http://www.cdc.gov/ncidod/dbmd/diseaseinfo/foodborneinfections_g.htm

<http://www.cdc.gov/health/diseases.htm>

2000 FoodNet Working Group

CDC

Frederick Angulo
Timothy Barrett
Michael Beach
Nancy Bean
Richard Bishop
Chris Braden
Laura Conn
Stephanie DeLong
Sara Ehlers
Cindy Friedman
Patricia Griffin
Peggy Hayes
Mike Hoekstra
Jeff Jones
Malinda Kennedy
Beth Imhoff
Jenny Lay
Deborah Levy
Kathleen Maloney
Paul Mead
Thomas Navin
Oshine Najarian
Robert Pinner
Laurence Slutsker
Karen Stamey
Bala Swaminathan
Robert Tauxe
Thomas Van Gilder
David Wallace
Stephanie Wong

California

Sharon Abbott
Mary Ann Davis
Pam Daily
Lisa Gelling
Alexander McNeas
Janet Mohle-Boetani
Nandeeni Mukerjee
Joelle Nadle
Jan O'Connell
Gretchen Rothrock
Michael Samuel
Sue Shallow
Ben Silk
Duc Vugia
Ben Werner

Colorado

Jim Beebe
Steve Burnite
Matt Finke
Ken Gershman
Lucinda Hammond
Sally Hauser
Sean Lieske
Sue Lynch
Ellen Mangione
Pam Shillam

Connecticut

Gary Budnick
Matthew Cartter
Terry Fiorentino
James Hadler
Robert Howard
Kati Kelley
Aristea Kinney
Ruthanne Marcus
Donald Mayo
Patricia Mshar
Quyen Phan
Robin Ryder
Charles Welles

Georgia

Wendy Baughman
Paul Blake
Laurel Boykin
Sabrina Burden
Shama Desai
Monica Farley
Katherine Gibbs-
McCombs
Laura Gilbert
Susan Lance-Parker
Susan Ray
Cathy Rebmman
Matthew Sattah
Suzanne Segler

Maryland

Alicia Bustamante
Michael Carter
Yvonne Deane-Hibbert

Diane Dwyer
Lora Gay
Althea Glenn
Charmaine Gregg
Marguerite Hawkins
Kim Holmes
Jackie Hunter
Kelly Henning
Tobi Karchmer
Meghan McGavern
J. Glenn Morris, Jr.
Lola Olabode
Peggy Pass
Jafar Razeq
Jeffery Roche
Dale Rohn
Christian Steiner
Alexander Sulakvelidze
Yongyu Wang
Frances Yarber

Minnesota

Jeff Bender
John Besser
Richard Danila
Valerie Deneen
Craig Hedberg
Heidi Kassenborg
Carlota Medus
Kirk Smith
Dana Soderlund
Ellen Swanson
Julie Wicklund

New York

Bridget Anderson
Dianna Bopp
Kathy Carlton
Hwa-Gan Chang
Barbara Damaske
Nellie Dumas
Marie Fitzgerald
Karim Hechemy
Jonathan Hibbs
Dale Morse
Candace Noonan
Brian Sauders
Perry Smith

Nancy Spina
Shelley Zansky

Oregon

Vijay Balan
Chris Biggs
Maureen Cassidy
Paul Cieslak
Emilio DeBess
David Fleming
Bill Keene
Lore Lee
Eileen Lorber
Steve Mauvais
Teresa McGivern
Yijun Pang
Beletshachew Shiferaw
Bob Sokolow

Tennessee

Effie Boothe
Allen Craig
Diane Eigsti Gerber
Timothy Jones
William Moore
William Schaffner
Pat Turri

USDA-FSIS

Noreen Hynes
Tamar Lasky
Denise Lewis
Phyllis Sparling
Kaye Wachsmuth

FDA-CFSAN

Ken Falci
Bing Garthright
Clifford Purdy

For more information on FoodNet, visit our web-page at:

www.cdc.gov/foodnet