

TABLE 2

Laboratory confirmed *Shigella* isolates reported to the CDC by serotype in 2006

Rank	Serotype	Reported	Percent
1	<i>S. sonnei</i>	7471	72.3
2	<i>S. flexneri</i> unspecified	746	7.2
3	<i>S. flexneri</i> 2 unspecified	152	1.5
4	<i>S. flexneri</i> 2a	107	1.0
5	<i>S. flexneri</i> 1 unspecified	102	1.0
6	<i>S. boydii</i> unspecified	75	0.7
7	<i>S. flexneri</i> 3 unspecified	67	0.7
8	<i>S. flexneri</i> 6	61	0.6
9	<i>S. flexneri</i> 4 unspecified	54	0.5
10	<i>S. flexneri</i> 1b	38	0.4
11	<i>S. flexneri</i> 3a	38	0.4
12	<i>S. flexneri</i> 4a	37	0.4
13	<i>S. dysenteriae</i> unspecified	31	0.3
14	<i>S. flexneri</i> 2b	22	0.2
15	<i>S. flexneri</i> variant y	22	0.2
16	<i>S. boydii</i> 2	15	0.2
17	<i>S. flexneri</i> 1a	12	0.1
18	<i>S. boydii</i> 1	9	0.1
19	<i>S. flexneri</i> 3b	7	0.1
20	<i>S. boydii</i> 14	6	0.1
21	<i>S. flexneri</i> 4b	5	0.1
22	<i>S. dysenteriae</i> 2	4	0.0
23	<i>S. boydii</i> 4	3	0.0
24	<i>S. dysenteriae</i> 4	3	0.0
25	<i>S. flexneri</i> variant x	3	0.0
26	<i>S. boydii</i> 12	2	0.0
27	<i>S. boydii</i> 18	2	0.0
28	<i>S. dysenteriae</i> 1	2	0.0
29	<i>S. dysenteriae</i> 12	2	0.0
30	<i>S. dysenteriae</i> 3	2	0.0
31	<i>S. flexneri</i> 5 unspecified	2	0.0
32	<i>S. boydii</i> 10	1	0.0
33	<i>S. boydii</i> 8	1	0.0
34	<i>S. dysenteriae</i> 13	1	0.0
35	<i>S. dysenteriae</i> 7	1	0.0
36	<i>S. flexneri</i> 5b	1	0.0
37	<i>S. flexneri</i> 88-893	1	0.0
	Sub Total	9108	88.2
	Unknown	1228	11.9
	Sub Total	1228	11.9
	Total	10336	100.0