



Published in final edited form as:

Exerc Sport Sci Rev. 2024 April 01; 52(2): 39–46. doi:10.1249/JES.0000000000000332.

A Novel Conceptual Model for Human Heat Tolerance

Thomas E. Bernard¹, S. Tony Wolf², W. Larry Kenney³

¹College of Public Health, University of South Florida, Tampa, FL 33612

²Department of Kinesiology, University of Georgia, Athens, GA 30602

³Department of Kinesiology and Integrative and Biomedical Physiology Program, The Pennsylvania State University, University Park, PA 16802

Abstract

Human “heat tolerance” has no accepted definition or physiological underpinnings; rather it is almost always discussed in *relative or comparative* terms. We propose to use environmental limits to heat balance accounting for metabolic rate and clothing, that is, the environments for which heat stress becomes uncompensable for a specified metabolic rate and clothing, as a novel metric for quantifying heat tolerance.

Summary

We propose a novel quantification of human heat tolerance based on empirical assessment of the environment, metabolic rate, and clothing.

Keywords

climate change; critical environmental limit; wet-bulb globe temperature; heat stress; occupational exposure limit

Introduction

Heat tolerance is an ambiguous term when applied to humans, with multiple physiological, psychological, pathological, and biophysical definitions and interpretations. Wyndham defined heat tolerance as the ability to “thermoregulate well enough to work under most conditions of heat stress” (1, 2). While *heat intolerance* is not a medical or pathophysiological condition, it is often defined as a hypersensitivity to heat that may or may not have pathological consequences.

As such, heat tolerance is almost always discussed in *relative or comparative* terms, e.g., improved physiological responses with heat acclimatization (3–5), performance differences between or among individuals or groups in stressful environments, or passing/failing a standardized test (e.g., a heat tolerance test (6)). As an example of the latter, the Israeli

Corresponding Author: S. Tony Wolf, 115G Ramsey Center, 330 River Road, Athens, GA 30602, Phone: 706-542-0636, stwolf@uga.edu.

Conflicts of Interest: None declared

Defense Forces (IDF) uses a treadmill-based heat-tolerance test (HTT) as a screening tool to assist in decision-making on a soldier's return to duty after an episode of exertional heat illness. However, core temperature and heart rate responses to such a test are not finite markers of heat tolerance, but rather serve as a useful adjunct to the full medical record. There is no *absolute* or *categorical* measure of an individual's heat tolerance and, concomitantly, no understanding of how heat tolerance is distributed within a sample or population.

Part of the problem of describing heat tolerance is that heat stress and the resulting heat strain are multidimensional. Heat stress involves a combination of environmental factors and net metabolic heat production that is further modified by clothing (7). Therefore, at a minimum, a heat stress metric needs to account for these three factors. The individual response to heat stress (i.e., heat strain) is also modified by personal factors such as body size, age, and sex at birth (8, 9). Therefore, an absolute measure of heat tolerance would, by necessity, incorporate how heat tolerance is (1) driven by multidimensional determinants of heat stress and (2) modified by personal factors.

The purpose of this *Perspectives for Progress* is to introduce a novel conceptual model of individual heat tolerance grounded in empirical human subject data from two laboratories. The model describes how the ability to tolerate heat can be operationalized both for an individual and for a population. Knowing an individual's heat tolerance can inform a decision about specific heat stress exposures. Knowing the population distribution of heat tolerance can inform exposures that are sustainable for various percentages of the population. While relative measures of heat tolerance (e.g., improved heat tolerance, individual X is more heat tolerant than individual Y, etc.) have utility and are pervasive in the literature, heat tolerance can be described as a continuous metric once the effects of environment, exercise or work intensity, clothing, and personal factors are known. This type of measure would enhance our understanding of contributors to heat strain and provide a basis for decision-making for sustainable exercise and occupational work performance in hot environments.

Conceptual Model of Individual Heat Tolerance

A simplified conceptual model for heat tolerance is shown in Figure 1. The central focus is the individual. **Heat stress** is a combination of environmental influences (ambient temperature, water vapor pressure, radiant heat load, effective air movement), metabolic rate (basal plus activity-based corrected for external work performed), and clothing (primarily area covered, insulation, and evaporative resistance). Heat stress also has a temporal spectrum of exposures both within a day and serial heat stress exposures over multiple days (e.g., carryover effects) (10–12). In defining heat tolerance, the exposure conditions are specified as constant with a short exposure time on the order of an hour to avoid drift over longer periods (13).

For a given heat exposure, the individual exhibits measurable physiological responses (i.e., **heat strain**). The magnitude of the strain response is moderated by personal factors. To categorize personal factors in the context of heat stress/strain, we propose

4 categories: physical characteristics, situational factors, modifiable factors, and inherent factors. Physical characteristics are regularly reported in describing the participants, and include anthropometrics such as height, weight, body surface area, and adiposity or body mass index (BMI), age, and sex. Situational factors are transient or short-term factors, lasting hours or days, and can often be managed by policies and practices of the supervising organization, or by the individual when self-directed. Modifiable factors are those that may be modified by the individual with some effort but are less amenable to organization oversight, such as fitness or wellness. Inherent personal factors are inherited or acquired, are chronic, and are not easily changed without extraordinary intervention, if at all. These groupings have flexible boundaries and are offered to focus research questions and develop interventions.

In the conceptual model, we consider heat strain as a generic descriptor of integrated physiological changes. Following the arrow pointing down, heat strain is evidenced by changes in heart rate, body temperatures, and sweating rate as well as other physiological adjustments (e.g., water turn-over, fluid balances, hormonal adjustments) and psychophysiological outcomes (e.g., perceived exertion, thermal sensation, and thermal comfort). At some point, the heat strain leads to deleterious outcomes including heat-related illness (11), acute injury (14, 15), and decreased performance with lost productivity (16, 17). The probability associated with the deleterious outcomes increases with heat stress, but the levels of *de minimus* or acceptable probability are not well defined.

Because of the complex nature of heat strain, the model shown in Figure 1 operationalizes heat strain by placing it into a Category. Categories are described on the right side of Figure 1 and are based on Lind's proposed upper limit of the prescriptive zone (ULPZ) (18). The ULPZ is the highest environmental heat stress in which the participant can maintain at relatively constant core temperature (T_c) at a given metabolic heat production and has more recently been redefined at the critical environmental limit (CEL), illustrated as the Critical Condition in the figure. The CEL forms the basis of our proposed metric for heat tolerance. The CEL is defined as the maximal heat stress at which the T_c depends solely on the metabolic rate and is relatively independent of the ambient environment (19). Above the CEL, heat loss (a function of integrated physiological responses) can no longer offset heat production plus heat gain, internal heat storage occurs, and core temperature increases. Lind's method for determining the CEL was modified by Belding and Kamon (1973), and refined in a series of papers by Kenney (20–23) and Bernard (24). In summary, the proposed heat tolerance metric is the CEL, which indicates a condition where heat stress can be sustained with no change in T_c . It is not a clear defining point for conditions safe from deleterious outcomes.

The lower end of the spectrum in the figure (green, compensable) represents a low level of heat stress, a level that the individual can tolerate for prolonged periods of time with minimal physiological adjustment. As the magnitude of heat stress increases, the collection of physiological adjustments (i.e., heat strain) increases, but the response remains compensable at a steady T_c driven only by the metabolic rate (19). That is, a sustainable thermal equilibrium can be established for at least the short term (25). This heat strain category is compensable. The upper limit of the compensable category is the CEL, which

is our proposed environmental metric for heat tolerance. As a note, there is a marginal zone above the CEL in which thermal equilibrium can be maintained for a short period of time, albeit with higher heat strain (26). Finally, increasing levels of heat stress move the individual out of the compensable category and into the category of uncompensable heat stress. The uncompensable category is marked by non-steady state responses and time-limited exposures (25).

The conceptual model also points to alternative descriptions for heat strain via the gray arrow from categories to deleterious effects. In epidemiological studies, the classification of exposure by these categories may help to parse out the health and wellbeing outcomes.

For the purpose of this model, heat tolerance for a specific metabolic rate and clothing ensemble is defined by the CEL -- the transition from compensable to uncompensable heat stress (18, 23, 24, 27–31) as shown in Figure 2.

Establishing Critical Environmental Limits

The ability to define the CEL is based on the historical work of Lind (18, 32–34) and Belding and Kamon (28). The protocols have been optimized by more recent projects in our respective laboratories at the University of South Florida (24, 35, 36) and Penn State University (20–23, 29, 30). In most applications, metabolic rate and clothing are fixed and a starting environment is selected in the compensable range based on experience. Following a 30- to 45-min equilibration period, the environmental heat stress is increased in small increments every 5 min in stepwise fashion. Heart rate begins to increase slowly as the CEL is approached (Cottle 2023). In environments beyond the CEL, T_c increases $\sim 0.5 - 1.5^\circ\text{C}$ per hour (37). As previously described, the CEL is determined as the environmental conditions immediately before the T_c inflection point during a progressive heat stress protocol. An example time course for T_c is shown in Figure 2. We have previously reported excellent reliability and validity of this protocol to identify CELs (38).

In practice, the PSU and USF progressive heat stress protocols differ slightly in how the environment is controlled. The essential element to the progressive heat stress protocol is systematically changing environmental heat stress in small steps that allow a steady state response within ~ 5 min. At PSU, there is a critical dry bulb temperature (T_{crit}) approach that starts with a fixed water vapor pressure and T_{db} is increased about 1°C per 5-min step. The critical water vapor pressure (P_{crit}) approach starts with a fixed T_{db} and P_a is increased about 1 mmHg per step. At USF, the critical environment is found by increasing T_{db} in 1°C steps at a fixed relative humidity. In this approach both T_{db} and P_a are increased concurrently, and the CEL is reported as the critical wet bulb globe temperature ($WBGT_{crit}$). Data from both approaches can be combined into multiple psychrometric parameters, including (psychrometric) wet-bulb temperature (T_{wb} or T_{pwb}). Cottle et al. (2021) demonstrated that the T_{crit} and P_{crit} approaches yield the same CEL.

Alternative approaches to finding critical environmental conditions have been proposed. For instance, Lind (18) and Kuhlmeier (39, 40) used a series of 1-h exposures to heat stress with a rest break between increasing levels of heat stress. The data can then be inspected to find the inflection point for T_c . Another example of a progressive heat stress protocol is to

increase the metabolic rate in a constant environment (41). In theory, the progressive heat stress protocol can be used for any condition that can be changed in small, monotonic steps in the direction of increasing heat stress. Miller and Bates used consecutive periods of 30 min in a constant environment in which they adjusted metabolic rate up or down in search of a steady-state T_{c} at the highest metabolic rate (42).

Quantifying Individual Heat Tolerance

Data at the CEL are important indicators of the transition from compensable to uncompensable heat stress at specified metabolic rates and clothing ensembles, as demonstrated by the PSU Heat project described below (29, 37). For the purposes of operationalizing a heat tolerance as a single metric or index, we start by recognizing that heat stress is a combination of environment and metabolic rate and thus needs a single metric to represent both. On a psychrometric chart, a constant wet bulb globe temperature (WBGT) index has a slope of $-0.18^{\circ}\text{C}\cdot\text{kPa}^{-1}$. Kenney reported WBGT slopes of -0.20 and $-0.16^{\circ}\text{C}\cdot\text{kPa}^{-1}$ for woven clothing (i.e., work clothes and coveralls) (43) and Bernard reported slopes of -0.16 and $-0.17^{\circ}\text{C}\cdot\text{kPa}^{-1}$ for similar clothing (36). Because the WBGT and woven clothing have similar slopes, we chose the WBGT as the best index to represent the environment. We therefore propose three methods using critical wet bulb globe temperature ($\text{WBGT}_{\text{crit}}$) and adjust it for metabolic rate and clothing.

The first method for using $\text{WBGT}_{\text{crit}}$ is an industry-based approach driven by investigators from the US National Institute for Occupational Safety and Health (NIOSH) (44). NIOSH guidelines use a relation between WBGT and metabolic rate that describes a limiting WBGT at a given metabolic rate. This relation is termed the Recommended Exposure Limit (REL) by NIOSH (45), the Threshold Limit Value® (TLV®) by the ACGIH (46) and simply a limit value by the International Organization for Standardization (47). The WBGT-based *Occupational Exposure Limit* (OEL) is a general description of these limits. The operational method is to relate the measured $\text{WBGT}_{\text{crit}}$ outcome to the OEL (27, 48). Referencing heat tolerance to the OEL has the advantage of immediate applicability in WBGT-based exposure assessments. Using the WBGT-based OEL, the difference between $\text{WBGT}_{\text{crit}}$ and the OEL can be defined as ΔOEL as a function of metabolic rate (M) in watts (27, 48) such that

$$\Delta\text{OEL} = \text{WBGT}_{\text{crit}} - \text{OEL} = \text{WBGT}_{\text{crit}} - [56.7 - 11.5\log_{10}M]$$

Eq 1

In this manner, a positive value for ΔOEL (> 0) means that the individual has a heat tolerance greater than that required to work at the OEL. In contrast, a negative ΔOEL (< 0) means that the individual does not have sufficient heat tolerance for a sustainable exposure at the OEL. We also demonstrated that this approach fully accounts for the relation between $\text{WBGT}_{\text{crit}}$ and metabolic rate (49). While this is called an occupational limit, it is based on Lind's ULPZ which is applicable to non-occupational exposures as well.

A second method (48) normalizes the heat stress exposure by adjusting the value of $\text{WBGT}_{\text{crit}}$ to a reference metabolic rate (48). Garzón and colleagues reported a slope in a linear relation between $\text{WBGT}_{\text{crit}}$ in $^{\circ}\text{C}$ and metabolic rate in watts as $-0.0201^{\circ}\text{C}\cdot\text{W}^{-1}$

(27). Based on this relation, WBGT referenced to a steady-state work intensity of 300 W (WBGT_{300W}) is

$$\text{WBGT}_{300\text{W}} = \text{WBGT}_{\text{crit}} + 0.0201 * (M - 300)$$

Eq 2

This approach to normalization means that WBGT_{300W} is greater than WBGT_{crit} when the metabolic rate is greater than 300 W and is less than WBGT_{crit} when the metabolic rate is less than 300 W. This method still allows an approximate comparison to the OEL by comparing WBGT_{300W} to the OEL at 300 W (i.e., 28 °C) (45, 46).

The third approach follows the same procedure for WBGT_{300W} but uses the metabolic rate divided by body surface area (BSA). Garzón and colleagues reported a slope in a linear relation between WBGT_{crit} in °C and metabolic rate in watts divided by BSA in m² as -0.0367 °C m²/W (27). Based on this relation, WBGT referenced to a steady-state work intensity of 150 W/m² (WBGT_{150W/m2}) is

$$\text{WBGT}_{150\text{W/m}^2} = \text{WBGT}_{\text{crit}} + 0.0367 * (M/\text{BSA} - 150)$$

Eq 3

The advantage of this method is that it accounts for the relation between WBGT_{crit} and BSA (49). The downside of this method is many exposure assessment methods do not use metabolic rate adjusted for BSA.

Heat Tolerance Within a Population

The preceding sections described (1) methods to identify heat tolerance for an individual followed by (2) three methods that can be used to normalize heat tolerance data. Also of interest would be the distribution of heat tolerance within a population, e.g., a particular workforce. For instance, a heat tolerance profile for healthy, hydrated, and acclimatized participants wearing woven clothing over a range of metabolic rates and humidity conditions using the OEL metric is illustrated in Figure 3a. It is clear from the figure that there was a wide range of heat tolerance values from 0 at the OEL to 12°C above the OEL with a mean value at 6°C above the OEL. A similar outcome was found when WBGT_{300W} was used (see Figure 3b) where 28°C was equivalent to the OEL. Finally, the distribution standardized to BSA is offered (Figure 3c). These figures represent heat tolerance profiles for the selected population of young, healthy, acclimatized, and hydrated individuals wearing a single layer of woven clothing over shorts and tee shirt/sports bra.

To develop a heat tolerance profile for a specific group of interest, such as middle-distance track athletes or workers with diabetes, the OEL (or WBGT_{300W} or WBGT_{150W/m2}) data are rank ordered from the lowest to highest (48, 50). The probability (p) for each observation is the rank (i, starting at 1) divided by the number of observations (n) plus 1; that is $p_i = \frac{i}{n+1}$.

The odds for each observation is $odds_i = \frac{p_i}{1 - p_i}$. Then a ln(odds) linear regression estimates

$$\ln(odds) = a + b \Delta OEL$$

Eq 4

for the intercept (a) and slope (b). Finally, the data are plotted as p versus OEL and the descriptive line is

$$p = \frac{e^{(a + b \Delta OEL)}}{1 + e^{(a + b \Delta OEL)}}.$$

Eq 5

Utility of Heat Tolerance Metrics

Heat tolerance metrics have been used (1) to describe the effects of clothing on heat stress (50), (2) to estimate the effect of heat acclimatization on heat tolerance (49), and (3) to prescribe heat stress limits for everyday life (PSU HEAT), as described in the following sections.

Clothing Adjustment Values

The OELs were originally developed with individuals wearing long-sleeved cotton shirt and trousers and this cotton work uniform was used as the reference value. $WBGT_{crit}$ was used to demonstrate that there is no difference between the cotton work uniform and cotton coveralls (35, 36), or compared to cotton versus flame resistant fabrics based on treated cotton or manmade fibers (51). The approach was also used to demonstrate that full-face negative pressure respirators do not add to the level of heat stress (52). In contrast, nonwoven, particle barrier, water barrier, and vapor barrier fabrics add the equivalent of 1, 3 and 10°C WBGT, respectively, to the critical exposure (35, 36). As another example, the effect of wearing a hood (head and neck covering) added 1°C to the WBGT (51). Additionally, $WBGT_{crit}$ was used to demonstrate that the effective air permeability of clothing explained the evaporative resistance better than moisture vapor transfer rate (41).

A Clothing Adjustment Value (CAV) is used to quantify the effective change in environmental WBGT with respect to the reference work clothes. CAVs (24, 35, 36, 53) adopted by the ACGIH in promulgating their Threshold Limit Values (TLV) (46) for work in hot environments as well as the ISO WBGT-based exposure assessment (47). While CAV values are based on averages, the exposure-response profiles for four different clothing ensembles described shift by the average amount for woven and vapor-permeable non-woven fabrics (50). In the context of the conceptual model, the personal risk factors were controlled by a balanced, crossover design so that the different contributions of clothing could be investigated independent of the person.

Further, CAVs can be used in Equations 1 to 3 to adjust data for different clothing ensembles (48). The CAV is subtracted from the observed $WBGT_{crit}$ to obtain an effective $WBGT_{crit}$. For example, Equation 4 is representative of Equations 1 for OEL with the addition of CAV.

$$\Delta OEL = WBGT_{crit} - CAV - OEL = WBGT_{crit} - CAV - [56.7 - 11.5 \log_{10} M]$$

Eq 6

Heat Acclimatization

Most occupational and sport-based guidelines and standards use the term “acclimatization” to encompass both the short-term, often artificial (acclimation) and longer-term, natural (acclimatization) physiological adjustments to heat stress. The conceptual model for quantifying heat tolerance can be used to investigate the contributions of acclimatization to heat tolerance. For example, occupational health and safety policy allows that acclimatization increases the acceptable WBGT exposure limit by ~2.5 to 3°C (44). The exposure response curve for 9 women and 9 men who were unacclimatized and seminude working at 30% of their individual maximum aerobic capacity based on OEL (48) helps demonstrate the shift. There were two exposures each at high and low humidity levels.

OEL allowed for adjusting the different metabolic rates. The effects of acclimation were explored further over a broader range of metabolic rates and environmental conditions (49). The unacclimatized curve shifted 3°C to the left from Figure 3a and demonstrated a wider variability due to individual differences than the difference assigned to acclimatization.

Demonstration of Safe Limits for Everyday Life

Using the single metric approach to defining heat tolerance described in the previous section is not the only way to quantitatively describe heat tolerance. The PSU HEAT project demonstrates the utility of the CEL directly. Identification of CEL values at metabolic rates associated with activities of daily living provides important information regarding heat tolerance limits for free-living individuals during heat waves. Recently, the PSU HEAT project has demonstrated the CEL for young adults (18–34 y) at two low rates of metabolic heat production across a wide range of environments (29, 37). Subjects were tested in eight distinct environments ranging from warm-humid (P_{crit} trials) to hot-dry (T_{crit} trials) during light ambulation on a treadmill (LightAmb; $M_{net} = \sim 130 \text{ W} \cdot \text{m}^{-2}$) and during minimal activity performed on a cycle ergometer (MinAct; $M_{net} = \sim 80 \text{ W} \cdot \text{m}^{-2}$). Those two metabolic rates were intended to approximate the demands of leisurely walking and activities of daily living performed in the home, respectively.

The mean CEL data with lower bounds of the 95% confidence intervals (CI; i.e., the one-tailed 5th percentile) were plotted as psychrometric limits (i.e., properties of atmospheric air that influence heat balance such as temperature and humidity) on a standard psychrometric chart (Figure 4)(29). Using this graphical approach, the compensability of any combination of temperature and humidity can be assessed. Environments that are below and to the left of the 95% CI lines are considered to be compensable, and therefore reasonably well-tolerated for prolonged exposures, for 95% of the population in this age range. Combinations that are above and to the right of the limit lines are uncompensable and therefore may be unsafe for prolonged exposures without any mitigating measures.

By examining the rate of rise in T_c below and above the CEL, projections can be made from and to any theoretical T_c to assess risk in real-world scenarios. In one investigation (37), heat

storage and the rate of rise in T_c were examined below and above the CEL in MinAct and LightAmb. The rate of rise in T_c below or above the CEL was not significantly influenced by the environment or metabolic rate. Using the slopes established in that study, during physical activity in environments below the CEL, it would take an average of ~36 h without any intervention to reach a $T_c > 40^\circ\text{C}$. In environments above the CEL, however, it would take as little as 3–7 h. These data may lend to development of recommendations for safe exposure times in environments that are above CEL.

Future Perspectives

The preceding sections summarized past and current applications of the heat tolerance model. The two sections below describe some future applications as examples of other possibilities to demonstrate how the conceptual model can be used to frame research and practice questions.

Role of Heat Stress Contributors in the Conceptual Model

From the basic premise that heat stress can be described as the combined effect of environment (i.e., WBGT), metabolic rate, and clothing, the discussion of CAVs demonstrates how the model is used for differentiating among clothing ensembles with a crossover design to control for personal factors. In a similar approach, the cost of lower evaporative resistance on productivity was demonstrated by adjusting metabolic rate rather than WBGT, controlling for personal factors in a crossover design (41).

There are few controlled studies on how well radiant heat is accounted for in WBGT. The question can be addressed by asking if OEL (or $WBGT_{300W}$ or $WBGT_{150/m^2}$) and its profile are the same with and without one or more levels of radiant heat and the incidence angle of that radiant heat on the person.

In the context of communicating heat risk to the general population, various heat metrics that consider T_{db} and other environmental characteristics including humidity, wind velocity, and radiative heat gain have been proposed. Along with the WBGT, these indices include the Heat Index (HI), the Universal Thermal Climate Index (UTCI), and the Humidex. Charts have been constructed for each of those metrics to denote danger thresholds at different combinations of temperature and humidity, although it is unclear how the thresholds were established.

Empirical CEL data from the PSU HEAT project were used to examine the effectiveness of WBGT, the HI, and the UTCI danger thresholds for predicting the temperature-humidity relation at the transition from compensable to uncompensable heat stress (54). For each metric, the two highest danger thresholds were used to compare against the mean and 95% CI CEL data. At very low metabolic rates, the HI was adequate for describing the temperature-humidity relation at the upper limit of compensability, whereas the WBGT was less suitable and the UTCI was not at all effective. At increasing metabolic rates, however, none of the indices adequately described the transition from compensable to uncompensable heat stress.

Role of Personal Factors in the Conceptual Model

Personal factors are a key feature of the conceptual model. Asking questions about the effects of individual or combined factors on heat tolerance is a straightforward process.

Anthropometrics.—Anthropometric factors help characterize a population and can be used as a treatment effect in a study design. For instance, comparing men versus women, young versus old, or lean versus obese. For these questions, the control of situational factors is common. More difficult is the control of modifiable and inherent factors. To the extent that contrasting populations with identified modifiable or inherent factors can be identified, the heat tolerance of each population can be compared. The unidentified factors are usually confined to the error term.

Situational Personal Factors.—The situational personal factors in the conceptual model are those factors that can be controlled in a study and where an intervention can be easily and obviously part of the supervising organization's policies and practices. It is widely recognized that acclimatization state and hydration state affect heat strain and thus heat tolerance. In addition, situational factors such as performance expectations, voluntary and forced overtime, work practices, and incentives can affect heat tolerance. Another situational factor is the pattern of exposures over days, in which high heat exposures on a previous day (12) or days (10, 11) influence the risk of heat disorders on the current day. Among personal factors, acclimatization and hydration are easily managed by policies and practices with some cooperation from the individual. Other situational factors may be less obvious but can be controlled by the organization. There may be times that the organization must include the formal cooperation of labor such as setting pay and overtime rules, or the athletic organization sets policies on the needed recovery time from a previous endurance event (e.g., distance running). In this regard, some considerations may include:

- Tracking heat tolerance following different protocols for recovery within a day or across days;
- Track progression of heat tolerance following different acclimation/acclimatization protocols;
- Compare the effects of prescribed versus self-directed recovery at work or during team practices.

Modifiable Personal Factors.—Modifiable personal factors may reduce heat tolerance but may be changed by the individual outside of the context of a research study or heat stress management. These factors include, among others, obesity, use of certain medications and drugs, lack of sleep, and low fitness level. This group of personal factors can be modified by the individual; however, there is a need to educate individuals about how these intrinsic factors can negatively impact their heat tolerance. For instance, many may not know that sleep deprivation may make them more susceptible to heat stroke (9). Although the organization may have no control over these personal factors, organizational policies and support can promote the risk reduction of these personal factors.

- Compare obese to lean with matching of fitness (using WBGT_{150/m2});

- Compare heat tolerance to use v. non-use of OTC supplements;
- Well rested and recovered from previous exposure due to lack of sleep.

Inherent Personal Factors.—The fourth group of personal factors are inherent to the individual and are generally refractory to interventions. These personal factors include genetics, physiological capacity, and chronic disease.

- Contrast healthy to specific diseases that may impair heat tolerance;
- Contrast normal genetic/epigenetic markers to those associated with increased heat susceptibility.

Personal factors collectively contribute to the inter-individual variability and may explain intra-individual variability. Further, there may be interactions between heat stress and personal factors that can be explored. The conceptual model provides a framework to see how they change heat tolerance.

Summary

Determination of the Critical Environmental Limit (CEL) at a specified metabolic rate and clothing condition using a progressive heat stress protocol is an efficient and effective method to assess a threshold environmental limit between compensable and uncompensable heat stress, i.e., heat tolerance. It is then a simple step to quantify heat tolerance with the addition of metabolic rate and clothing. The Conceptual Model provides a framework to assess the effects on heat tolerance due to personal factors. The model divides risk factors into four categories of demographics and how easily the risk factors can be a focus of intervention. The current paper demonstrates how the CEL and heat tolerance were used to demonstrate quantitatively the difference between unacclimatized and acclimatized participants, to assign clothing adjustment values to account for different clothing, and to map out psychrometric limits for defined activity and clothing.

Acknowledgements

Funding:

NIH Grant R01 AG067471 (WLK) and CDC/NIOSH Grant R01 OH03983 (TEB).

References

1. Bernard TE, Dukes-Dobos FN, Ramsey JD. Evaluation and control of hot working environments: Part II – The scientific basis (knowledge base) for the guide. In: Mital A, Kilbom Å, Kumar S, editors. Elsevier Ergonomics Book Series. 1: Elsevier; 2000. p. 337–46.
2. Wyndham CH. Research in the Human Sciences in the Gold Mining Industry. American Industrial Hygiene Association journal. 1974;35(3):113–36. doi: 10.1080/0002889748507014. [PubMed: 4595259]
3. Périard JD, Racinais S, Sawka MN. Adaptations and mechanisms of human heat acclimation: Applications for competitive athletes and sports. Scand J Med Sci Sports. 2015;25 Suppl 1:20–38. doi: 10.1111/sms.12408. [PubMed: 25943654]
4. Taylor NAS. Principles and practices of heat adaptation. Journal of the Human-Environment System. 2000;4(1):11–22.

5. Tyler CJ, Reeve T, Hodges GJ, Cheung SS. The Effects of Heat Adaptation on Physiology, Perception and Exercise Performance in the Heat: A Meta-Analysis. *Sports Med.* 2016;46(11):1699–724. doi: 10.1007/s40279-016-0538-5. [PubMed: 27106556]
6. Butler C, Dierickx E, Bruneau M, Stearns R, Casa DJ. Current Clinical Concepts: Heat Tolerance Testing. *J Athl Train.* 2023;58(2):84–90. doi: 10.4085/1062-6050-352-21. [PubMed: 34793593]
7. Havenith G Heat balance when wearing protective clothing. *Ann Occup Hyg.* 1999;43(5):289–96. [PubMed: 10481628]
8. Notley SR, Flouris AD, Kenny GP. Occupational heat stress management: Does one size fit all? *Am J Ind Med.* 2019;62(12):1017–23. [PubMed: 30791115]
9. Périard JD, DeGroot D, Jay O. Exertional heat stroke in sport and the military: epidemiology and mitigation. *Exp Physiol.* 2022;107(10):1111–21. [PubMed: 36039024]
10. Shire J, Vaidyanathan A, Lackovic M, Bunn T. Association Between Work-Related Hyperthermia Emergency Department Visits and Ambient Heat in Five Southeastern States, 2010–2012-A Case-Crossover Study. *Geohealth.* 2020;4(8):e2019GH000241.
11. Spector JT, Sampson L, Flunker JC, Adams D, Bonauto DK. Occupational heat-related illness in Washington State: A descriptive study of day of illness and prior day ambient temperatures among cases and clusters, 2006–2021. *Am J Ind Med.* 2023;66(8):623–36. [PubMed: 37291066]
12. Wallace RF, Kriebel D, Punnett L, Wegman DH, Wenger CB, Gardner JW, et al. The effects of continuous hot weather training on risk of exertional heat illness. *Med Sci Sports Exerc.* 2005;37(1):84–90. [PubMed: 15632673]
13. Smallcombe JW, Foster J, Hodder SG, Jay O, Flouris AD, Havenith G. Quantifying the impact of heat on human physical work capacity; part IV: interactions between work duration and heat stress severity. *Int J Biometeorol.* 2022. Epub 20221005. doi: 10.1007/s00484-022-02370-7.
14. Fogleman M, Fakhzadeh L, Bernard TE. The relationship between outdoor thermal conditions and acute injury in an aluminum smelter. *International Journal of Industrial Ergonomics.* 2005;35:47–55.
15. Spector JT, Masuda YJ, Wolff NH, Calkins M, Seixas N. Heat Exposure and Occupational Injuries: Review of the Literature and Implications. *Curr Environ Health Rep.* 2019;6(4):286–96. doi: 10.1007/s40572-019-00250-8. [PubMed: 31520291]
16. Foster J, Smallcombe JW, Hodder S, Jay O, Flouris AD, Nybo L, et al. An advanced empirical model for quantifying the impact of heat and climate change on human physical work capacity. *Int J Biometeorol.* 2021;65(7):1215–29. Epub 20210305. doi: 10.1007/s00484-021-02105-0. [PubMed: 33674931]
17. Sahu S, Sett M, Kjellstrom T. Heat exposure, cardiovascular stress and work productivity in rice harvesters in India: implications for a climate change future. *Industrial health.* 2013;51(4):424–31. Epub 20130520. doi: 10.2486/indhealth.2013-0006. [PubMed: 23685851]
18. Lind AR. A physiological criterion for setting thermal environmental limits for everyday work. *J Appl Physiol.* 1963;18:51–6. Epub 1963/01/01. [PubMed: 13930723]
19. Bernard TE, Ashley CD, Wolf ST, Kenney WL. Core temperature and heart rate at the upper limit of the prescriptive zone. *Physiol Rep.* 2023;11(17):e15812. doi: 10.14814/phy2.15812. [PubMed: 37688426]
20. Kenney WL, Lewis DA, Hyde DE, Dyksterhouse TS, Armstrong CG, Fowler SR, et al. Physiologically derived critical evaporative coefficients for protective clothing ensembles. *J Appl Physiol* (1985). 1987;63(3):1095–9. Epub 1987/09/01. doi: 10.1152/jappl.1987.63.3.1095. [PubMed: 3654457]
21. Kenney WL, Hyde DE, Bernard TE. Physiological evaluation of liquid-barrier, vapor-permeable protective clothing ensembles for work in hot environments. *American Industrial Hygiene Association journal.* 1993;54(7):397–402. Epub 1993/07/01. doi: 10.1080/15298669391354865. [PubMed: 8362761]
22. Kenney WL, Mikita DJ, Havenith G, Puhl SM, Crosby P. Simultaneous derivation of clothing-specific heat exchange coefficients. *Med Sci Sports Exerc.* 1993;25(2):283–9. Epub 1993/02/01. [PubMed: 8450734]
23. Kenney WL, Zeman MJ. Psychrometric limits and critical evaporative coefficients for unacclimated men and women. 2002;92(6):2256–63. doi: 10.1152/japplphysiol.01040.2001.

24. O'Connor DJ, Bernard TE. Continuing the search for WBGT clothing adjustment factors. *Applied occupational and environmental hygiene*. 1999;14(2):119–25. Epub 1999/08/24. [PubMed: 10457638]
25. Cottle RM, Lichter ZS, Vecellio DJ, Wolf ST, Kenney WL. Core temperature responses to compensable versus uncompensable heat stress in young adults (PSU HEAT Project). *J Appl Physiol* (1985). 2022;133(4):1011–8. Epub 2022/09/01. doi: 10.1152/jappphysiol.00388.2022. [PubMed: 36049058]
26. Pandolf KB, Burr RE. *Medical Aspects of Harsh Environments*. Washington DC: US Army Office of the Surgeon General; 2001.
27. Garzón-Villalba XP, Wu Y, Ashley CD, Bernard TE. Ability to Discriminate Between Sustainable and Unsustainable Heat Stress Exposures-Part 1: WBGT Exposure Limits. *Ann Work Expo Health*. 2017;61(6):611–20. Epub 2017/06/09. doi: 10.1093/annweh/wxx034. [PubMed: 28595332]
28. Belding HS, Kamon E. Evaporative coefficients for prediction of safe limits in prolonged exposures to work under hot conditions. *Fed Proc*. 1973;32(5):1598–601. Epub 1973/05/01. [PubMed: 4705206]
29. Wolf ST, Cottle RM, Vecellio DJ, Kenney WL. Critical environmental limits for young, healthy adults (PSU HEAT). *J Appl Physiol* (1985). 2021. Epub 2021/12/17. doi: 10.1152/jappphysiol.00737.2021.
30. Kenney WL. Psychrometric limits and critical evaporative coefficients for exercising older women. 2020;129(2):263–71. doi: 10.1152/jappphysiol.00345.2020.
31. Dougherty KA, Chow M, Larry Kenney W. Critical environmental limits for exercising heat-acclimated lean and obese boys. *European journal of applied physiology*. 2010;108(4):779–89. doi: 10.1007/s00421-009-1290-4. [PubMed: 20187283]
32. Lind AR. Effect of individual variation on upper limit of prescriptive zone of climates. *J Appl Physiol*. 1970;28(1):57–62. Epub 1970/01/01. [PubMed: 5409790]
33. Lind AR, Bass DE. Optimal exposure time for development of acclimatization to heat. *Federation Proceedings*. 1963;22:704–8. [PubMed: 13930724]
34. Lind AR, Humphreys PW, Collins KJ, Foster K, Sweetland KF. Influence of age and daily duration of exposure on responses of men to work in heat. *J Appl Physiol*. 1970;28(1):50–6. Epub 1970/01/01. [PubMed: 5409789]
35. Bernard TE, Caravello V, Schwartz SW, Ashley CD. WBGT clothing adjustment factors for four clothing ensembles and the effects of metabolic demands. *J Occup Environ Hyg*. 2008;5(1):1–5. Epub 2007/11/14. doi: 784715198 [pii]10.1080/15459620701732355 [doi]. [PubMed: 17999329]
36. Bernard TE, Luecke CL, Schwartz SW, Kirkland KS, Ashley CD. WBGT clothing adjustments for four clothing ensembles under three relative humidity levels. *Journal of occupational and environmental hygiene*. 2005;2(5):251–6. Epub 2005/04/05. doi: 10.1080/15459620590934224 [doi]. [PubMed: 15804982]
37. Cottle RM, Lichter ZS, Vecellio DJ, Wolf ST, Kenney WL. Core temperature responses to compensable vs. uncompensable heat stress in young adults (PSU Heat Project). *J Appl Physiol* (1985). 2022. Epub 2022/09/02. doi: 10.1152/jappphysiol.00388.2022.
38. Cottle RM, Wolf ST, Lichter ZS, Kenney WL. Validity and Reliability of a Protocol to Establish Human Critical Environmental Limits (PSU HEAT). *J Appl Physiol* (1985). 2021. Epub 2021/12/17. doi: 10.1152/jappphysiol.00736.2021.
39. Kuhlmeier KV, Miller JM, Dukes-Dobos FN, Jensen R. Determinants of the prescriptive zone of industrial workers. *J Appl Physiol*. 1977;43(2):347–51. Epub 1977/08/01. [PubMed: 893294]
40. Kuhlmeier KV, Wood TB. Laboratory evaluation of permissible exposure limits for men in hot environments. *Am Ind Hyg Assoc J*. 1979;40(12):1097–103. doi: 10.1080/15298667991430758. [PubMed: 539550]
41. Gonzalez NW, Bernard TE, Carroll NL, Bryner MA, Zeigler JP. Maximum sustainable work rate for five protective clothing ensembles with respect to moisture vapor transmission rate and air permeability. *Journal of occupational and environmental hygiene*. 2006;3(2):80–6. doi: 10.1080/15459620500498133. [PubMed: 16418081]

42. Miller V, Bates G. The thermal work limit is a simple reliable heat index for the protection of workers in thermally stressful environments. *Annals of Occupational Hygiene*. 2007;51(6):553–61. doi: 10.1093/annhyg/mem035. [PubMed: 17878259]
43. Kenney WL, Lewis DA, Armstrong CG, Hyde DE, Dyksterhouse TS, Fowler SR, et al. Psychrometric limits to prolonged work in protective clothing ensembles. *Am Ind Hyg Assoc J*. 1988;49(8):390–5. doi: 10.1080/15298668891379954. [PubMed: 3177217]
44. Dukes-Dobos FN, Henschel A. Development of permissible heat exposure limits for occupational work. *Am Soc Heating, Refrig Air Cond Eng J*. 1973;15:57–62.
45. NIOSH. NIOSH criteria for a recommended standard: Occupational exposure to heat and hot environments. By Jacklitsch B, Williams WJ, Musolin K, Coca A, Kim J-H, Turner N. Cincinnati, OH: Department of Health and Human Services, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health (DHHS (NIOSH) Publication No. 2016–106); 2016.
46. ACGIH®. TLV for Heat Stress and Strain. Threshold Limit Values and Biological Exposure Indices for Chemical Substances and Physical Agents. Cincinnati, OH: ACGIH®; 2023.
47. ISO. ISO 7243:2017 Ergonomics of the thermal environment — Assessment of heat stress using the WBGT (wet bulb globe temperature) index. Geneva: ISO; 2017.
48. Wolf ST, Bernard TE, Kenney WL. Heat exposure limits for young unacclimatized males and females at low and high humidity. *Journal of occupational and environmental hygiene*. 2022;1–15. Epub 20220510. doi: 10.1080/15459624.2022.2076859.
49. Bernard TE, Ashley CD, Wolf ST, Odera AM, Lopez RM, Kenney WL. Distribution of upper limit of the prescriptive zone values for acclimatized and unacclimatized individuals. *J Appl Physiol* (1985). 2023;135(3):601–8. Epub 20230727. doi: 10.1152/japplphysiol.00724.2022. [PubMed: 37498291]
50. Garzón-Villalba XP, Wu Y, Ashley CD, Bernard TE. Heat stress risk profiles for three non-woven coveralls. *Journal of occupational and environmental hygiene*. 2018;15(1):80–5. Epub 2017/10/21. doi: 10.1080/15459624.2017.1388514. [PubMed: 29053928]
51. Ashley CD, Bernard TE. Effects of hoods and flame-retardant fabrics on WBGT clothing adjustment factors. *J Occup Environ Hyg*. 2008;5(1):59–62. doi: 787378963 [pii]10.1080/15459620701764358 [doi]. [PubMed: 18041645]
52. Fletcher OM, Guerrina R, Ashley CD, Bernard TE. Heat stress evaluation of two-layer chemical demilitarization ensembles with a full face negative pressure respirator. *Industrial health*. 2014;52(4):304–12. Epub 2014/04/08. doi: 10.2486/indhealth.2012-0197. [PubMed: 24705801]
53. Kenney WL. WBGT adjustments for protective clothing. *American Industrial Hygiene Association Journal*. 1987;48:576–7.
54. Vecellio DJ, Wolf ST, Cottle RM, Kenney WL. Utility of the Heat Index in defining the upper limits of thermal balance during light physical activity (PSU HEAT Project). *International Journal of Biometeorology*. 2022;66(9):1759–69. doi: 10.1007/s00484-022-02316-z. [PubMed: 35778555]

Key points

- Heat tolerance is a nebulous term and is almost always discussed in *relative or comparative* terms.
- We propose a novel absolute metric for individual heat tolerance grounded in empirical human data, driven by multidimensional heat stress, and modified by personal factors.
- We make the case that individual heat tolerance based on critical environmental limits, the upper limit for heat balance across a wide range of ambient temperatures and humidity, is an appropriate metric.
- The effects of environment, exercise or work intensity, clothing, and personal factors can be quantified into a conceptual absolute heat tolerance category.




Figure 1.

Conceptual model of heat tolerance with heat stress factors and personal factors driving an individual's heat stress response (heat strain). Heat tolerance is the limiting sustainable response at the transition from compensable to uncompensable heat stress, also known as the critical environmental limit (CEL) at a specified metabolic rate and clothing ensemble. The distal outcomes of a heat stress exposure and the resulting heat strain are heat-related illnesses, decreased performance, and increased risk of injury. The response category (i.e., compensable vs. CEL vs. uncompensable) may provide an alternative classification for overall heat strain that modulates the risk of deleterious outcomes (illustrated by the gray arrow).




Figure 2.

Illustration of zones of compensable and uncompensable heat stress based on the core temperature response of an individual to increasing levels of environmental heat stress at a given metabolic rate. Heat tolerance is defined as the transition from compensable to uncompensable heat stress, i.e., the Critical Environmental Limit (CEL).




Figure 3.

Population distribution of heat tolerance in terms of the difference from the occupational exposure limit (OEL) (a), $\text{WBGT}_{300\text{W}}$ (b), and $\text{WBGT}_{150\text{W}/\text{m}^2}$ (c). The population was young, healthy, acclimatized, and hydrated individuals wearing a single layer of woven clothing over shorts and tee shirt/sports bra.




Figure 4.

A standard psychrometric chart demonstrated critical environmental limits for young, healthy adults during light ambulation on a treadmill (LightAmb; $M_{\text{net}} = \sim 130 \text{ W}\cdot\text{m}^{-2}$) and during minimal activity performed on a cycle ergometer (MinAct; $M_{\text{net}} = \sim 80 \text{ W}\cdot\text{m}^{-2}$). Psychrometric curves for LightAmb are shifted downward and to the left relative to MinAct as a result of increased metabolic heat production. Figure updated and redrawn from (29).